

Generalizing Riemann curvature to Regge metrics

Michael Neunteufel (Portland State University)

Jay Gopalakrishnan (Portland State University)


Joachim Schöberl (TU Wien)

Max Wardetzky (University of Göttingen)





ICOSAHOM 2025, Montreal, July 18, 2025



- N -dimensional Riemannian manifold (Ω, \bar{g})
with \bar{g} smooth metric tensor (spd bilinear form)





- N -dimensional Riemannian manifold (Ω, \bar{g})
with \bar{g} smooth metric tensor (spd bilinear form)
- g piecewise smooth metric on triangulation \mathcal{T} approximating \bar{g}



- N -dimensional Riemannian manifold (Ω, \bar{g})
with \bar{g} smooth metric tensor (spd bilinear form)
- g piecewise smooth metric on triangulation \mathcal{T} approximating \bar{g}
- Can we approximate **Riemann curvature** of (Ω, \bar{g}) with (\mathcal{T}, g) ?
Can we **generalize** curvatures to piecewise smooth metrics g ?



- N -dimensional Riemannian manifold (Ω, \bar{g})
with \bar{g} smooth metric tensor (spd bilinear form)
- g piecewise smooth metric on triangulation \mathcal{T} approximating \bar{g}
- Can we approximate **Riemann curvature** of (Ω, \bar{g}) with (\mathcal{T}, g) ?
Can we **generalize** curvatures to piecewise smooth metrics g ?



Yes, if g is in the Regge space!

- $\mathfrak{X}(\mathcal{T})$, $\Lambda^k(\mathcal{T})$, $\mathcal{T}_l^k(\mathcal{T})$ **piecewise smooth** vector, k -forms, and (k, l) -tensor fields
- $g \in \mathcal{S}(\mathcal{T}) = \{\sigma \in \mathcal{T}^2(\mathcal{T}) : \sigma(X, Y) = \sigma(Y, X) \text{ for all } X, Y \in \mathfrak{X}(\mathcal{T})\}$
- Elementwise Riemann curvature tensor $\mathcal{R} \in \mathcal{T}^4(\mathcal{T})$ by $(X, Y, Z, W \in \mathfrak{X}(\mathcal{T}))$

$$\mathcal{R}(X, Y, Z, W) = g(\mathcal{R}_{X,Y}Z, W) = g(\nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X,Y]}Z, W)$$

- $\mathfrak{X}(\mathcal{T})$, $\Lambda^k(\mathcal{T})$, $\mathcal{T}_l^k(\mathcal{T})$ **piecewise smooth** vector, k -forms, and (k, l) -tensor fields
- $g \in \mathcal{S}(\mathcal{T}) = \{\sigma \in \mathcal{T}^2(\mathcal{T}) : \sigma(X, Y) = \sigma(Y, X) \text{ for all } X, Y \in \mathfrak{X}(\mathcal{T})\}$
- Elementwise Riemann curvature tensor $\mathcal{R} \in \mathcal{T}^4(\mathcal{T})$ by $(X, Y, Z, W \in \mathfrak{X}(\mathcal{T}))$


$$\mathcal{R}(X, Y, Z, W) = g(\mathcal{R}_{X,Y}Z, W) = g(\nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X,Y]}Z, W)$$

- \mathcal{R} is a nonlinear **second-order** differential operator in g

$$\mathcal{R}_{ijkl} = \partial_i \Gamma_{jkl} - \partial_j \Gamma_{ikl} - \Gamma_{ilp} \Gamma_{jk}^p + \Gamma_{jlp} \Gamma_{ik}^p, \quad \Gamma_{jk}^p = g^{pq} \Gamma_{jkq}, \quad \Gamma_{ijk} = 1/2(\partial_i g_{jl} + \partial_j g_{il} - \partial_k g_{ij})$$

- “ $\mathcal{R}(g)$ ” is a **nonlinear distribution**. What are the curvature contributions on element interfaces?


- Regge's idea: Approximate metric by assigning squared lengths to edges



 REGGE: General relativity without coordinates, *Il Nuovo Cimento* (1955-1965), (1961).

- Regge's idea: Approximate metric by assigning squared lengths to edges
- With barycentric coordinates λ^i

$$g = - \sum_{i \neq j} l_{ij}^2 \nabla \lambda^i \odot \nabla \lambda^j$$




 REGGE: General relativity without coordinates, *Il Nuovo Cimento* (1955-1965), (1961).

 SORKIN: Time-evolution problem in Regge calculus, *Phys. Rev. D* 12 (1975).

- Regge's idea: Approximate metric by assigning squared lengths to edges
- With barycentric coordinates λ^i


$$g = - \sum_{i \neq j} l_{ij}^2 \nabla \lambda^i \odot \nabla \lambda^j$$



- g is piecewise constant and **tangential-tangential continuous**: for all interior facets F the value $g(X, Y)$ coincides from both elements for all tangential $X, Y \in \mathfrak{X}(F)$
- **Regge finite element space**


$$\mathcal{R}_h^k = \{g \in \mathcal{S}(\mathcal{T}) : g_{ij} \in \mathcal{P}^k(\mathcal{T}), g \text{ is tt-continuous}\}$$

 CHRISTIANSEN: On the linearization of Regge calculus, *Numerische Mathematik*, (2011).

 LI: Regge Finite Elements with Applications in Solid Mechanics and Relativity, *PhD thesis*, University of Minnesota (2018).

- Regge's idea: Approximate metric by assigning squared lengths to edges
- With barycentric coordinates λ^i


$$g = - \sum_{i \neq j} l_{ij}^2 \nabla \lambda^i \odot \nabla \lambda^j$$











- g is piecewise constant and **tangential-tangential continuous**: for all interior facets F the value $g(X, Y)$ coincides from both elements for all tangential $X, Y \in \mathfrak{X}(F)$
- **Regge finite element space**

$$\mathcal{R}_h^k = \{g \in \mathcal{S}(\mathcal{T}) : g_{ij} \in \mathcal{P}^k(\mathcal{T}), g \text{ is tt-continuous}\}$$

 CHRISTIANSEN: On the linearization of Regge calculus, *Numerische Mathematik*, (2011).

 N.: Mixed Finite Element Methods for Nonlinear Continuum Mechanics and Shells, *PhD thesis*, TU Wien (2021).










- In 2D, the angle defect Θ_V at vertex V is given by

$$\Theta_V = 2\pi - \sum_{T \ni V} \angle_V^T,$$


with the interior angle \angle_V^T is measured with $g|_T$





- In 2D, the angle defect Θ_V at vertex V is given by


$$\Theta_V = 2\pi - \sum_{T \supset V} \angle_V^T,$$




with the interior angle \angle_V^T is measured with $g|_T$

- N dimensions: generalized angle defect at $\mathring{e} = \{\text{interior subsimplices of codimension 2}\}$

$$\Theta_E = 2\pi - \sum_{T \supset E} \underbrace{\arccos(g(\mu_E^{F_+}, \mu_E^{F_-}))}_{\angle_E^T}$$



Plane g -perpendicular to E



- **Second fundamental form:** For hypersurface F with g -normal ν

$$\mathbb{I}^\nu(X, Y) = -g(\nabla_X \nu, Y), \quad X, Y \in \mathfrak{X}(F)$$

- Since the metric g and the g -normal ν jumps across interior facets $F \in \mathring{\mathcal{F}}$, the second fundamental form jumps as well
- **Facet contribution:** Jump of second fundamental form $[[\mathbb{I}]]$
- Motivation via Gauss-Bonnet theorem or mollification argument

- **Second fundamental form:** For hypersurface F with g -normal ν

$$\mathbb{I}^\nu(X, Y) = -g(\nabla_X \nu, Y), \quad X, Y \in \mathfrak{X}(F)$$

- Since the metric g and the g -normal ν jumps across interior facets $F \in \mathring{\mathcal{F}}$, the second fundamental form jumps as well
- **Facet contribution:** Jump of second fundamental form $[\![\mathbb{I}]\!]$
- Motivation via Gauss-Bonnet theorem or mollification argument

Summary: The generalized Riemann curvature tensor has the following contributions

$$\mathcal{R}(g) = \begin{cases} \mathcal{R}|_T & \text{on each } T \in \mathcal{T} \\ [\![\mathbb{I}]\!] & \text{on each } F \in \mathring{\mathcal{F}} \\ \Theta_E & \text{on each } E \in \mathring{\mathcal{E}} \end{cases}$$

Generalized Riemann curvature (Gopalakrishnan, N., Schöberl, Wardetzky)

$$\widetilde{\mathcal{R}\omega}(A) = \sum_{T \in \mathcal{T}} \int_T \langle \mathcal{R}, A \rangle \omega_T + 4 \sum_{F \in \mathring{\mathcal{F}}} \int_F \langle \llbracket II \rrbracket, A_{\cdot\nu\nu} |_F \rangle \omega_F + 4 \sum_{E \in \mathring{\mathcal{E}}} \int_E \Theta_E A_{\mu\nu\nu\mu} \omega_E$$

$\widetilde{\mathcal{R}\omega}$ is acting on $A \in \mathring{\mathcal{A}}$, ω_D volume form on D


$$A_{\cdot\nu\nu}(X, Y) = A(X, \nu, \nu, Y), \quad A_{\mu\nu\nu\mu} = A(\mu, \nu, \nu, \mu).$$

Test space $\mathring{\mathcal{A}}$ (has Riemann curvature tensor symmetries)

$$\mathcal{A} = \{A \in \mathcal{T}^4(\mathcal{T}) : A_{\cdot\nu\nu}|_F \text{ is single-valued for all } F \in \mathring{\mathcal{F}}, \text{ and}$$

$$A(X, Y, Z, W) = -A(Y, X, Z, W) = -A(X, Y, W, Z) = A(Z, W, X, Y)\}$$

$$\mathring{\mathcal{A}} = \{A \in \mathcal{A} : A_{\cdot\nu\nu}|_F = 0 \text{ on } \partial\Omega\}$$



$$\widetilde{\mathcal{R}\omega}(A) = \sum_{T \in \mathcal{T}} \int_T \langle \mathcal{R}, A \rangle \omega_T + 4 \sum_{F \in \mathring{\mathcal{F}}} \int_F \langle \llbracket II \rrbracket, A_{\cdot\nu\nu} \cdot |_F \rangle \omega_F + 4 \sum_{E \in \mathring{\mathcal{E}}} \int_E \Theta_E A_{\mu\nu\nu\mu} \omega_E$$

Set $A = -v \omega \otimes \omega$ for $v \in \mathring{\mathcal{V}} = \{u \in \Lambda^0(\mathcal{T}) : u \text{ continuous, } u|_{\partial\Omega} = 0\}$. Then

- On element T : $\langle \mathcal{R}|_T, A \rangle = 4K|_T v$, K Gauss curvature

Gauss curvature:

$$K = \frac{\mathcal{R}_{1221}}{\det g}$$

$$\widetilde{\mathcal{R}\omega}(A) = \sum_{T \in \mathcal{T}} \int_T \langle \mathcal{R}, A \rangle \omega_T + 4 \sum_{F \in \mathring{\mathcal{F}}} \int_F \langle \llbracket II \rrbracket, A_{\cdot \nu \nu \cdot} |_F \rangle \omega_F + 4 \sum_{E \in \mathring{\mathcal{E}}} \int_E \Theta_E A_{\mu \nu \nu \mu} \omega_E$$

Set $A = -v \omega \otimes \omega$ for $v \in \mathring{\mathcal{V}} = \{u \in \Lambda^0(\mathcal{T}) : u \text{ continuous, } u|_{\partial\Omega} = 0\}$. Then

- On element T : $\langle \mathcal{R}|_T, A \rangle = 4K|_T v$, K Gauss curvature
- On edge F : $\langle \llbracket II \rrbracket, A_{\cdot \nu \nu \cdot} |_F \rangle = \llbracket \kappa \rrbracket v$, κ geodesic curvature

Geodesic curvature: with g -unit tangent τ along edge F

$$\kappa = g(\nu, \nabla_\tau \tau) = II^\nu(\tau, \tau)$$


$$\widetilde{\mathcal{R}\omega}(A) = \sum_{T \in \mathcal{T}} \int_T \langle \mathcal{R}, A \rangle \omega_T + 4 \sum_{F \in \mathcal{F}} \int_F \langle \llbracket II \rrbracket, A_{\nu\nu} \cdot |_F \rangle \omega_F + 4 \sum_{E \in \mathcal{E}} \int_E \Theta_E A_{\mu\nu\nu\mu} \omega_E$$


Set $A = -v \omega \otimes \omega$ for $v \in \mathring{\mathcal{V}} = \{u \in \Lambda^0(\mathcal{T}) : u \text{ continuous, } u|_{\partial\Omega} = 0\}$. Then

- On element T : $\langle \mathcal{R}|_T, A \rangle = 4K|_T v$, K Gauss curvature
- On edge F : $\langle \llbracket II \rrbracket, A_{\nu\nu} \cdot |_F \rangle = \llbracket \kappa \rrbracket v$, κ geodesic curvature
- On vertex E : $\Theta_E A_{\mu\nu\nu\mu} = \Theta_E v$

Generalized Gauss curvature

$$\widetilde{K\omega}(v) = \frac{1}{4} \widetilde{\mathcal{R}\omega}(A) = \sum_{T \in \mathcal{T}} \int_T K|_T v \omega_T + \sum_{F \in \mathcal{F}} \int_F \llbracket \kappa \rrbracket v \omega_F + \sum_{E \in \mathcal{E}} \int_E \Theta_E v \omega_E$$

 BERCHENKO-KOGAN, GAWLIK: Finite element approximation of the Levi-Civita connection and its curvature in two dimensions, *Found Comput Math* (2022).

 GOPALAKRISHNAN, N., SCHÖBERL, WARDETSKY: Analysis of curvature approximations via covariant curl and incompatibility for Regge metrics, *SMAI J. Comput. Math.* (2023).

$$\widetilde{\mathcal{R}\omega}(A) = \sum_{T \in \mathcal{T}} \int_T \langle \mathcal{R}, A \rangle \omega_T + 4 \sum_{F \in \mathring{\mathcal{F}}} \int_F \langle \llbracket II \rrbracket, A_{\cdot\nu\nu\cdot}|_F \rangle \omega_F + 4 \sum_{E \in \mathring{\mathcal{E}}} \int_E \Theta_E A_{\mu\nu\nu\mu} \omega_E$$

Set $A = v g \oslash g$ for $v \in \mathring{\mathcal{V}} = \{u \in \Lambda^0(\mathcal{T}) : u \text{ continuous, } u|_{\partial\Omega} = 0\}$. Then

- On element T : $\langle \mathcal{R}|_T, A \rangle = 4S|_T v$, $S = g^{ij}g^{kl}\mathcal{R}_{kijl}$ scalar curvature

Kulkarni-Nomizu product \oslash : produces a 4-tensor from two symmetric 2-tensors with Riemann symmetries

$$(h \oslash k)(X, Y, Z, W) = h(X, W)k(Y, Z) + h(Y, Z)k(X, W) - h(X, Z)k(Y, W) - h(Y, W)k(X, Z)$$

$$\widetilde{\mathcal{R}\omega}(A) = \sum_{T \in \mathcal{T}} \int_T \langle \mathcal{R}, A \rangle \omega_T + 4 \sum_{F \in \mathring{\mathcal{F}}} \int_F \langle \llbracket H \rrbracket, A_{\cdot\nu\nu\cdot} |_F \rangle \omega_F + 4 \sum_{E \in \mathring{\mathcal{E}}} \int_E \Theta_E A_{\mu\nu\nu\mu} \omega_E$$

Set $A = v g \oplus g$ for $v \in \mathring{\mathcal{V}} = \{u \in \Lambda^0(\mathcal{T}) : u \text{ continuous, } u|_{\partial\Omega} = 0\}$. Then

- On element T : $\langle \mathcal{R}|_T, A \rangle = 4S|_T v$, $S = g^{ij} g^{kl} \mathcal{R}_{kijl}$ scalar curvature
- On facet F : $\langle \llbracket H \rrbracket, A_{\cdot\nu\nu\cdot} |_F \rangle = 2\llbracket H \rrbracket v$, H mean curvature

Mean curvature: for a facet F

$$H^\nu = \text{tr}(\Pi^\nu) = g^{ij} \Pi_{ij}^\nu$$

$$\widetilde{\mathcal{R}\omega}(A) = \sum_{T \in \mathcal{T}} \int_T \langle \mathcal{R}, A \rangle \omega_T + 4 \sum_{F \in \mathring{\mathcal{F}}} \int_F \langle \llbracket H \rrbracket, A_{\cdot \nu \nu} |_F \rangle \omega_F + 4 \sum_{E \in \mathring{\mathcal{E}}} \int_E \Theta_E A_{\mu \nu \nu \mu} \omega_E$$

Set $A = v g \oslash g$ for $v \in \mathring{\mathcal{V}} = \{u \in \Lambda^0(\mathcal{T}) : u \text{ continuous, } u|_{\partial\Omega} = 0\}$. Then

- On element T : $\langle \mathcal{R}|_T, A \rangle = 4S|_T v$, $S = g^{ij} g^{kl} \mathcal{R}_{kijl}$ scalar curvature
- On facet F : $\langle \llbracket H \rrbracket, A_{\cdot \nu \nu} |_F \rangle = 2\llbracket H \rrbracket v$, H mean curvature
- On E : $\Theta_E A_{\mu \nu \nu \mu} = 2\Theta_E v$

Generalized scalar curvature

$$\widetilde{S}\omega(v) = \frac{1}{4} \widetilde{\mathcal{R}\omega}(A) = \sum_{T \in \mathcal{T}} \int_T S|_T v \omega_T + 2 \sum_{F \in \mathring{\mathcal{F}}} \int_F \llbracket H \rrbracket v \omega_F + 2 \sum_{E \in \mathring{\mathcal{E}}} \int_E \Theta_E v \omega_E$$



$$\widetilde{\mathcal{R}\omega}(A) = \sum_{T \in \mathcal{T}} \int_T \langle \mathcal{R}, A \rangle \omega_T + 4 \sum_{F \in \mathring{\mathcal{F}}} \int_F \langle \llbracket II \rrbracket, A_{\cdot\nu\nu\cdot}|_F \rangle \omega_F + 4 \sum_{E \in \mathring{\mathcal{E}}} \int_E \Theta_E A_{\mu\nu\nu\mu} \omega_E$$

Set $A = g \oplus \sigma$ for $\sigma \in \mathring{\Sigma} = \{J\rho : \rho \in \mathring{\mathcal{R}}\}$. Then

- On element T : $\langle \mathcal{R}|_T, A \rangle = 4\langle \text{Ric}|_T, \sigma \rangle$, $\text{Ric} = g^{ij}\mathcal{R}_{kijl}$ Ricci curvature tensor

$J : \mathcal{S}(\mathcal{T}) \rightarrow \mathcal{S}(\mathcal{T})$ is a bijective algebraic operator

$$J\rho = \rho - \frac{1}{2}\text{tr}(\rho)g$$

$$\widetilde{\mathcal{R}\omega}(A) = \sum_{T \in \mathcal{T}} \int_T \langle \mathcal{R}, A \rangle \omega_T + 4 \sum_{F \in \mathcal{F}} \int_F \langle \llbracket II \rrbracket, A_{\nu\nu} \cdot |_F \rangle \omega_F + 4 \sum_{E \in \mathcal{E}} \int_E \Theta_E A_{\mu\nu\nu\mu} \omega_E$$

Set $A = g \oplus \sigma$ for $\sigma \in \mathring{\Sigma} = \{J\rho : \rho \in \mathring{\mathcal{R}}\}$. Then

- On element T : $\langle \mathcal{R}|_T, A \rangle = 4\langle \text{Ric}|_T, \sigma \rangle$, $\text{Ric} = g^{ij}\mathcal{R}_{kijl}$ Ricci curvature tensor
- On facet F : $\langle \llbracket II \rrbracket, A_{\nu\nu} \cdot |_F \rangle = \langle \llbracket II \rrbracket, \sigma|_F + \sigma(\nu, \nu)g|_F \rangle$
- On E : $\Theta_E A_{\mu\nu\nu\mu} = (\sigma(\nu, \nu) + \sigma(\mu, \mu)) \Theta_E$

Generalized Ricci curvature tensor

$$\widetilde{\text{Ric}\omega}(\sigma) = \sum_{T \in \mathcal{T}} \int_T \langle \text{Ric}|_T, \sigma \rangle \omega_T + \sum_{F \in \mathcal{F}} \int_F \langle \llbracket II \rrbracket, \sigma|_F + \sigma(\nu, \nu)g|_F \rangle \omega_F + \sum_{E \in \mathcal{E}} \int_E (\sigma(\nu, \nu) + \sigma(\mu, \mu)) \Theta_E \omega_E$$



$$\widetilde{\mathcal{R}\omega}(A) = \sum_{T \in \mathcal{T}} \int_T \langle \mathcal{R}, A \rangle \omega_T + 4 \sum_{F \in \mathcal{F}} \int_F \langle \llbracket II \rrbracket, A_{\nu\nu} \cdot |_F \rangle \omega_F + 4 \sum_{E \in \mathcal{E}} \int_E \Theta_E A_{\mu\nu\nu\mu} \omega_E$$

Use that $G = \text{Ric} - 1/2 S g = J \text{Ric}$. Set $A = g \oslash J\sigma$ for $\sigma \in \mathring{\mathcal{R}}$. Then

- On element T : $\langle \mathcal{R}|_T, A \rangle = 4 \langle G|_T, \sigma \rangle$
- On facet F : $\langle \llbracket II \rrbracket, A_{\nu\nu} \cdot |_F \rangle = \langle \llbracket II \rrbracket, \sigma|_F - \text{tr}(\sigma|_F)g|_F \rangle = \langle \llbracket II \rrbracket, \mathbb{S}_F \sigma|_F \rangle = \langle \llbracket \overline{II} \rrbracket, \sigma|_F \rangle$
where $\mathbb{S}_F \sigma = \sigma|_F - \text{tr}(\sigma|_F)g|_F$ and $\overline{II} = \mathbb{S}_F II = II - H g|_F$ the trace-reversed second fundamental form
- On E : $\Theta_E A_{\mu\nu\nu\mu} = -\text{tr}(\sigma|_E) \Theta_E$

Generalized Einstein tensor

$$\widetilde{G\omega}(\sigma) = \sum_{T \in \mathcal{T}} \int_T \langle G|_T, \sigma \rangle \omega_T + \sum_{F \in \mathcal{F}} \int_F \langle \llbracket \overline{II} \rrbracket, \sigma|_F \rangle \omega_F - \sum_{E \in \mathcal{E}} \int_E \text{tr}(\sigma|_E) \Theta_E \omega_E$$



- Assume that $g_h \in \mathcal{R}_h^k$ converges to a smooth metric \bar{g} for $h \rightarrow 0$. Does $\widetilde{\mathcal{R}\omega}_{g_h} \rightarrow (\mathcal{R}\omega)_{\bar{g}}$?
- **Approach:** Use its **integral representation**: For Gauss curvature

$$\widetilde{K\omega}_g(v) - (K\omega)_{\bar{g}}(v) = \int_0^1 \frac{d}{dt} \widetilde{K\omega}_{g(t)}(v) dt, \quad v \in \mathring{V},$$

where $g(t) = \bar{g} + t(g - \bar{g})$. Extend to N -dimensions

$$\widetilde{\mathcal{R}\omega}_g(A) - (\mathcal{R}\omega)_{\bar{g}}(A) = \int_0^1 \frac{d}{dt} \widetilde{\mathcal{R}\omega}_{g(t)}(A) dt, \quad A \in \mathring{A},$$



GAWLIK: High-order approximation of Gaussian curvature with Regge finite elements, *SIAM J. Numer. Anal.* (2020).

- Assume that $g_h \in \mathcal{R}_h^k$ converges to a smooth metric \bar{g} for $h \rightarrow 0$. Does $\widetilde{\mathcal{R}\omega}_{g_h} \rightarrow (\mathcal{R}\omega)_{\bar{g}}$?
- **Approach:** Use its **integral representation**: For Gauss curvature

$$\widetilde{\mathcal{K}\omega}_g(v) - (\mathcal{K}\omega)_{\bar{g}}(v) = \int_0^1 \frac{d}{dt} \widetilde{\mathcal{K}\omega}_{g(t)}(v) dt, \quad v \in \mathring{V},$$

where $g(t) = \bar{g} + t(g - \bar{g})$. Extend to N -dimensions

$$\widetilde{\mathcal{R}\omega}_g(A) - (\mathcal{R}\omega)_{\bar{g}}(A) = \int_0^1 \frac{d}{dt} \widetilde{\mathcal{R}\omega}_{g(t)}(A) dt, \quad A \in \mathring{A},$$

- **Problems:**
 1. Test function A depends on the metric tensor g
 2. Need to linearize curvature contributions



GAWLIK: High-order approximation of Gaussian curvature with Regge finite elements, *SIAM J. Numer. Anal.* (2020).

We use an approach inspired by the **Uhlenbeck trick**: Define the **metric independent** test space

$$\mathcal{U} = \{U \in \Lambda^{N-2}(\mathcal{T}) \odot \Lambda^{N-2}(\mathcal{T}) : U(X_1, \dots, X_{N-2}, Y_1, \dots, Y_{N-2}) \text{ is single-valued} \\ \text{on all } F \in \mathring{\mathcal{F}} \text{ for all } X_1, \dots, X_{N-2}, Y_1, \dots, Y_{N-2} \in \mathfrak{X}(F)\}$$

Lemma

The map $\mathbb{A}_g : \mathcal{U} \rightarrow \mathcal{A}$, $U \mapsto - \star^{\odot^2} U$ is a bijection.

Define $\widetilde{\mathcal{R}\omega\mathbb{A}}_g(U) = \widetilde{\mathcal{R}\omega}(\mathbb{A}_g(U))$ for g -independent $U \in \mathcal{U}$. Then we have

$$\widetilde{\mathcal{R}\omega\mathbb{A}}_{\mathbf{g}}(U) - (\mathcal{R}\omega\mathbb{A})_{\bar{\mathbf{g}}}(U) = \int_0^1 \frac{d}{dt} \widetilde{\mathcal{R}\omega\mathbb{A}}_{\mathbf{g}(t)}(U) dt.$$

We can proceed computing and estimating the right-hand side.

Theorem (Gopalakrishnan, N., Schöberl, Wardetzky)

Suppose g_h is a collection of Regge metrics such that $g_h \rightarrow \bar{g}$ in L^∞ and g_h is uniformly bounded in $W^{2,\infty}$. Then

$$\|\widetilde{\mathcal{R}\omega\mathbb{A}_{g_h}} - \mathcal{R}\omega\mathbb{A}_{\bar{g}}\|_{H^{-2}} \leq C_{\bar{g},g_h} \|g_h - \bar{g}\|_2.$$

Here,

$$\begin{aligned} \|\sigma\|_2^2 &= \sum_{T \in \mathcal{T}} \left(\|\sigma\|_{L^2(T)}^2 + h^2 \|\sigma\|_{H^1(T)}^2 + h^4 \|\sigma\|_{H^2(T)}^2 \right) \\ C_{\bar{g},g_h} &= C \left(1 + \max_{T \in \mathcal{T}} h_T^{-2+\delta_2^N} \|g_h - \bar{g}\|_{L^\infty(T)} + \max_{T \in \mathcal{T}} h_T^{-1} \|g_h - \bar{g}\|_{W^{1,\infty}(T)} \right) \end{aligned}$$

Corollary

If additionally $\|g_h - \bar{g}\|_{W^{t,\infty}} \lesssim h^{s-t} \|\bar{g}\|_{W^{s,\infty}}$ for $0 \leq t \leq s \leq k+1$ for some $k \geq 1 - \delta_2^N$, then

$$\|\widetilde{\mathcal{R}\omega\mathbb{A}_{g_h}} - \mathcal{R}\omega\mathbb{A}_{\bar{g}}\|_{H^{-2}} \lesssim \mathcal{O}(h^{k+1}).$$

Incompatibility operator

Define for smooth 2-tensor σ the **incompatibility operator** $\text{Inc} : \mathcal{T}^2 \rightarrow \mathcal{T}^4$ by

$$(\text{Inc } \sigma)(X, Y, Z, W) := \frac{1}{4} [(\nabla_{Y,Z}^2 \sigma)(X, W) + (\nabla_{X,W}^2 \sigma)(Y, Z) - (\nabla_{X,Y}^2 \sigma)(Z, W) - (\nabla_{Y,W}^2 \sigma)(X, Z)].$$

In 2D and 3D Inc can be related to the standard incompatibility operator $\text{inc} = \text{curl}^T \text{curl}$.

Lemma (linearization Riemann curvature tensor)

For t -independent vector fields $X, Y, Z, W \in \mathfrak{X}(T)$ there holds

$$\dot{\mathcal{R}}(X, Y, Z, W) = -\frac{1}{2}(\text{Inc } \dot{g})(X, Y, Z, W) + \frac{1}{2} [\dot{g}(\mathcal{R}_{X,Y} Z, W) - \dot{g}(\mathcal{R}_{X,Y} W, Z)].$$

Generalized incompatibility operator

For tt -continuous σ , a generalized Inc can be defined as

$$\widetilde{\text{Inc } \sigma}(A) = \sum_{T \in \mathcal{T}} \int_T \langle \text{Inc } \sigma, A \rangle \omega_T + \sum_{F \in \mathring{\mathcal{F}}} \cdots + \sum_{E \in \mathring{\mathcal{E}}} \cdots$$

$$1. \widetilde{\mathcal{R}\omega\mathbb{A}}_{g_h}(U) - (\mathcal{R}\omega\mathbb{A})_{\bar{g}}(U) = \int_0^1 \frac{d}{dt} \widetilde{\mathcal{R}\omega\mathbb{A}}_{g(t)}(U) dt \quad \text{with } g(t) = \bar{g} + t(g_h - \bar{g}).$$

$$2. \frac{d}{dt} \widetilde{\mathcal{R}\omega\mathbb{A}}_{g(t)}(U) = a(g; \dot{g}, U) + b(g; \dot{g}, U) \quad \text{sum of the bilinear forms } a \text{ and } b.$$

$$3. b(g; \dot{g}, U) = -2 \widetilde{\text{Inc}} \dot{g}(A), \quad \text{with } \dot{g} = g_h - \bar{g} \text{ and } A = \mathbb{A}_g(U).$$




- Analyze the adjoint: $\widetilde{\text{Inc}} \dot{g}(A) = (\widetilde{\text{Inc}}^* A)(\dot{g})$

Then all spatial derivatives are applied on the test function A , not \dot{g} .

- $b(g; \dot{g}, U) \leq C_{g_h, \bar{g}} \|g_h - \bar{g}\|_2 \|U\|_{H^2}$




- $$4. a(g; \dot{g}, U) \text{ has no spatial derivatives of } \dot{g}$$
- $a = 0$ in 2D, but $a \neq 0$ in higher dimensions
 - $a(g; \dot{g}, U) \leq C_{g_h, \bar{g}} \|g_h - \bar{g}\|_2 \|U\|_{H^2}$

$k = 0$



NGSolve

$k = 1$



all terms

no angle defect


no geodesic curvature

$$\widetilde{K}\omega(v) = \sum_{T \in \mathcal{T}} \int_T K|_T v \omega_T \quad + \quad \sum_{E \in \mathcal{E}} \int_E \Theta_E v \omega_E \quad + \quad \sum_{F \in \mathcal{F}} \int_F \llbracket \kappa \rrbracket v \omega_F$$

$$\begin{aligned}\Phi(x, y, z) &= (x, y, z, f(x, y, z)), \\ f(x, y, z) &= \frac{1}{2}(x^2 + y^2 + z^2) - \frac{1}{12}(x^4 + y^4 + z^4) \\ \bar{g} &= \nabla \Phi^T \nabla \Phi, \quad q(x) = x^2(x^2 - 3)^2\end{aligned}$$

$$\mathcal{R}_{ijkl} = \varepsilon_{ijr} \varepsilon_{kls} \delta^{rs} \frac{9 \prod_{m \neq r} (x_m^2 - 1)}{q(x) + q(y) + q(z) + 9}$$


- Confirms theory for $k \geq 1$
- For $k = 0$ **linear convergence** is observed?!



$$\begin{aligned}\Phi(x, y, z) &= (x, y, z, f(x, y, z)), \\ f(x, y, z) &= \frac{1}{2}(x^2 + y^2 + z^2) - \frac{1}{12}(x^4 + y^4 + z^4) \\ \bar{g} &= \nabla \Phi^T \nabla \Phi, \quad q(x) = x^2(x^2 - 3)^2\end{aligned}$$


$$\mathcal{R}_{ijkl} = \varepsilon_{ijr} \varepsilon_{kls} \delta^{rs} \frac{9 \prod_{m \neq r} (x_m^2 - 1)}{q(x) + q(y) + q(z) + 9}$$

- Confirms theory for $k \geq 1$
- For $k = 0$ **linear convergence** is observed?!
- Test only parts where theory indicates no convergence




Test convergence of theoretical sub-optimal terms (F_1 , F_2 , $F_1 + F_2 =: F_3$)

We observe rapid convergence, then stagnation of error \rightarrow pre-asymptotic



$k = 0$



$k = 1$

- Definition of generalized Riemann curvature tensor (Gauss, scalar, Ricci, Einstein)
- Numerical analysis with integral representation
- Uhlenbeck trick for test functions
- Generalized incompatibility operator and adjoint

- Extrinsic curvature of embedded submanifolds & connection 1-form
- Framework combining discrete differential geometry and distributional FEM
- NGSDiffGeo: <https://github.com/MichaelNeunteufel/NGSDiffGeo>
- Application to (nonlinear) shell analysis, geometric flows, and numerical relativity

- Extrinsic curvature of embedded submanifolds & connection 1-form
- Framework combining discrete differential geometry and distributional FEM
- NGSDiffGeo: <https://github.com/MichaelNeunteufel/NGSDiffGeo>
- Application to (nonlinear) shell analysis, geometric flows, and numerical relativity

Thank You for Your attention!



GOPALAKRISHNAN, N., SCHÖBERL, WARDETZKY: Analysis of curvature approximations via covariant curl and incompatibility for Regge metrics, *SMAI J. Comput. Math.* (2023).



GOPALAKRISHNAN, N., SCHÖBERL, WARDETZKY: On the improved convergence of lifted distributional Gauss curvature from Regge elements, *RINAM* (2024).



GAWLIK, N.: Finite element approximation of scalar curvature in arbitrary dimension, *Math. Comp.* (2024).



GAWLIK, N.: Finite element approximation of the Einstein tensor, *IMA J. Numer. Anal.* (2025).



GOPALAKRISHNAN, N., SCHÖBERL, WARDETZKY: Generalizing Riemann curvature to Regge metrics, *arXiv:2311.01603*.

Lemma

Let $\sigma := \dot{g}(t)$, $A \in \mathcal{A}$, $(SA)(X, Y, Z, W) = A(X, Z, Y, W)$ swaps second with third argument. There holds

$$\begin{aligned}\dot{A}(X, Y, Z, W) = & -\text{tr}(\sigma)A(X, Y, Z, W) + A(\sigma(X, \cdot)^\sharp, Y, Z, W) + A(X, \sigma(Y, \cdot)^\sharp, Z, W) \\ & + A(X, Y, \sigma(Z, \cdot)^\sharp, W) + A(X, Y, Z, \sigma(W, \cdot)^\sharp),\end{aligned}$$

Lemma

Let $\sigma := \dot{g}(t)$, $A \in \mathcal{A}$, $(SA)(X, Y, Z, W) = A(X, Z, Y, W)$ swaps second with third argument. There holds

$$\begin{aligned} \dot{A}(X, Y, Z, W) = & -\text{tr}(\sigma)A(X, Y, Z, W) + A(\sigma(X, \cdot)^\sharp, Y, Z, W) + A(X, \sigma(Y, \cdot)^\sharp, Z, W) \\ & + A(X, Y, \sigma(Z, \cdot)^\sharp, W) + A(X, Y, Z, \sigma(W, \cdot)^\sharp), \end{aligned}$$

$$\frac{d}{dt}(\langle \mathcal{R}, A \rangle \omega_T) = (2\langle \nabla^2 \sigma, S(A) \rangle + \langle \mathcal{R}(\sigma(\cdot, \cdot)^\sharp, \cdot, \cdot, \cdot), A \rangle - \frac{1}{2}\text{tr}(\sigma)\langle \mathcal{R}, A \rangle) \omega_T,$$

$$\frac{d}{dt}(\langle \llbracket \mathbb{I} \rrbracket, A_{\nu\nu\cdot|_F} \rangle \omega_F) = \frac{1}{2} \langle \llbracket (\sigma(\nu, \nu) - \text{tr}(\sigma|_F)) \mathbb{I} + 2(\nabla_F \sigma)(\nu, \cdot)|_F - (\nabla_\nu \sigma)|_F \rrbracket, A_{\nu\nu\cdot|_F} \rangle \omega_F,$$

$$\frac{d}{dt}(\Theta_E A_{\mu\nu\nu\mu} \omega_E) = -\frac{1}{2} \left(\sum_{F \supset E} \llbracket \sigma(\nu, \mu) \rrbracket_F^E + \text{tr}(\sigma|_E) \Theta_E \right) A_{\mu\nu\nu\mu} \omega_E.$$

Proposition

Let $\sigma := \dot{g}$ and $A \in \dot{\mathcal{A}}$ with corresponding $U = \mathbb{A}_g^{-1}A \in \dot{\mathcal{U}}$. Then there holds

$$\frac{d}{dt} \widetilde{\mathcal{R}\omega\mathbb{A}_g}(U) = a(g; \sigma, U) + b(g; \sigma, U),$$

$$\begin{aligned} a(g; \sigma, U) = & \sum_{T \in \mathcal{T}} \int_T (\langle \mathcal{R}(\sigma(\cdot, \cdot)^\sharp, \cdot, \cdot, \cdot), \mathbb{A}_g U \rangle - \frac{1}{2} \text{tr}(\sigma) \langle \mathcal{R}, \mathbb{A}_g U \rangle) \omega_T \\ & - 2 \sum_{F \in \mathcal{F}} \int_F (\text{tr}(\sigma|_F) \langle \llbracket II \rrbracket, (\mathbb{A}_g U)_{\cdot\nu\nu\cdot}|_F \rangle - \llbracket II \rrbracket : \sigma|_F : (\mathbb{A}_g U)_{\cdot\nu\nu\cdot}|_F) \omega_F \\ & - 2 \sum_{E \in \mathcal{E}} \int_E \text{tr}(\sigma|_E) \Theta_E(\mathbb{A}_g U)_{\mu\nu\nu\mu} \omega_E \end{aligned}$$

$$\llbracket II \rrbracket : \sigma|_F : (\mathbb{A}_g U)_{\cdot\nu\nu\cdot} = \llbracket II \rrbracket_{ij} (\sigma|_F)^{jk} ((\mathbb{A}_g U)_{\cdot\nu\nu\cdot})_k^i.$$

Proposition

Let $\sigma := \dot{g}$ and $A \in \dot{\mathcal{A}}$ with corresponding $U = \mathbb{A}_g^{-1}A \in \dot{\mathcal{U}}$. Then there holds

$$\frac{d}{dt} \widetilde{\mathcal{R}\omega_{\mathbb{A}_g}}(U) = a(g; \sigma, U) + b(g; \sigma, U),$$

$$\begin{aligned} b(g; \sigma, U) = & 2 \sum_{T \in \mathcal{T}} \int_T \langle \nabla^2 \sigma, S \mathbb{A}_g U \rangle \omega_T \\ & + 2 \sum_{F \in \mathcal{F}} \int_F \langle [\![\sigma(\nu, \nu)]\!] + (\nabla_F \sigma)(\nu, \cdot)|_F + \nabla_F(\sigma(\nu, \cdot))|_F - (\nabla_\nu \sigma)|_F, (\mathbb{A}_g U)_{\cdot \nu \nu} |_F \rangle \omega_F \\ & - 2 \sum_{E \in \mathcal{E}} \int_E \sum_{F \supset E} [\![\sigma(\nu, \mu)]\!]_F^E (\mathbb{A}_g U)_{\mu \nu \nu \mu} \omega_E. \end{aligned}$$