



Growth & mortality outcomes for different antiretroviral therapy initiation criteria in children aged 1-5 years:

– a causal modelling analysis from West and Southern Africa –

19th International Workshop on HIV Observational Databases
Catania, Italy

Background

Methodology

Results

Conclusions

Limitations

27 March 2015

The leDEA West & Southern Africa Paediatric Group



- Limited evidence from RCTs on when to start ART in children aged 1-5 years
- CHER study: Early ART initiation reduces mortality in infants (enrolled at < 3 months of age)

Violari (2008), NEJM

- PREDICT study, children aged 1-12 years:
 - No difference between starting ART immediately and deferring ART with respect to mortality and morbidity outcomes
 - However, better height gain for children who started ART immediately

Phutanakit (2012), Lancet Infectious Diseases

Background

Methodology


Results

Conclusions

Limitations

Background: PREDICT results

Michael Schomaker



Background

Methodology

Results

Conclusions

Limitations



To use observational data from children aged 1-5 years in the leDEA-WA and leDEA-SA collaborations to compare

- cumulative mortality
- growth

up to 3 years between 4 initiation strategies

- i) starting ART immediately, irrespective of CD4 criteria \approx WHO 2013
- ii) starting ART when CD4 drops below 750 cells/mm³ or 25% \approx WHO 2010
- iii) starting ART when CD4 drops below 350 cells/mm³ or 15% \approx WHO 2006
- iv) starting ART not at all

Background

Methodology

Results

Conclusions

Limitations

Inclusion criteria

We included 5826 ART naive children, with at least one follow-up visit, from 16 cohorts from West and Southern Africa

Michael Schomaker




Background

Methodology

Results

Conclusions

Limitations



Data from West Africa

Michael Schomaker



Background

Methodology

Results

Conclusions

Limitations



- The primary analysis used *g-computation* to estimate cumulative mortality and growth (mean HAZ of children who are alive) for different interventions strategies.
- We adjusted for time dependent confounding affected by prior treatment of CD4 count, CD4%, and Weight for age z-score (WAZ, proxy for WHO stage)
- missing baseline data was multiply imputed

Background

Methodology

Results

Conclusions

Limitations



This analysis emulates the following clinical trial:

- HIV positive and ART-naïve children, aged 1-5 years, presenting at a health care facility for the first time, are randomly assigned one of the four treatment strategies.
- Each of the four arms is therefore differing by the CD4 thresholds used to determine the timing of ART initiation.
- Assuming full adherence to the regime, no administrative censoring, no loss to follow-up, and regular CD4 monitoring we can estimate mortality/growth at time t ($t = 1, \dots, 36$ months after first visit).

[Background](#)

[Methodology](#)

[Results](#)


[Conclusions](#)

[Limitations](#)



	Southern Africa	West Africa
	Median (1 st ; 3 rd quartile)	
Age	2.6 (1.7; 3.7)	2.7 (1.8; 3.8)
CD4 count	646 (380; 984)	719 (433; 1081)
CD4%	16 (11; 23)	16 (10; 22)
WAZ	-1.5 (-2.6; -0.6)	-1.9 (-3.2; -0.9)
HAZ	-2.6 (-3.6; -1.6)	-2.1 (-3.1; -1.0)
started ART	75.6%	72.5%

[Background](#)[Methodology](#)[Results](#)[Conclusions](#)[Limitations](#)




[Background](#)

[Methodology](#)

[Results](#)

[Conclusions](#)

[Limitations](#)



Background

Methodology

Results

Conclusions

Limitations

Difference immediate ART to 750/25% at 3ys: -0.02 (-0.04; 0.01)

Difference immediate ART to 350/15% at 3ys: -0.08 (-0.10; -0.05)



- Lower mortality and better growth when starting ART earlier in children aged 1-5.
- These differences were small when comparing immediate ART initiation with deferring ART until the CD4 threshold of 750/25% is reached, but clearer when comparing it with the CD4 threshold of 350/15%.
- Our findings were consistent over age groups and regions, but mortality was estimated to be lower, and growth to be faster, in children aged 2-5 [data not shown].

[Background](#)

[Methodology](#)

[Results](#)

[Conclusions](#)

[Limitations](#)



- Missing HAZ data
[but sensitivity analyses reassuring]
- Competing event: death
[but sensitivity analyses reassuring]
- Other: Long term outcomes (toxicity, drug resistance),
WAZ as proxy for stage,

[Background](#)

[Methodology](#)

[Results](#)

[Conclusions](#)

[Limitations](#)



- All patients and staff from participating sites.
- Martina Penazzato (WHO), Lulu Muhe Mussa (WHO), Shaffiq Essajee (Clinton Foundation), and Thomas Gsponer (University of Bern)
- Funders: US National Institute of Allergy and Infectious Diseases (NIAID) through the International epidemiological Databases to Evaluate AIDS, Southern Africa (leDEA-SA).

Background

Methodology

Results

Conclusions

Limitations

Writing team: Michael Schomaker, Mary-Ann Davies, Karen Malateste, Lorna Renner, Shobna Sawry, Sylvie N'Gbeche, Karl-Günter Technau, François Eboua, Frank Tanser, Haby Sygnatè-Sy, Sam Phiri, Madeleine Amorissani-Folquet, Vivian Cox, Fla Koueta, Cleophas Chimbete, Annette Lawson-Evi, Janet Giddy, Clarisse Amani-Bosse, Robin Wood, Matthias Egger, Valeriane Leroy