

# Chapter 1

## Sensing

### 1.1 Quantum state distrimination with Bosonic channels aand Gaussian states

This PhD thesis is written by Sihui Tan, who is the one that first propose Quantum Illumination. So this thesis is a very valuable material to learn QI.

#### 1.1.1 Mathematical preliminaries

**Definition 1.1.1** (Direct sum of matrix).

#### 1.1.2 Physics preliminaries

**Lemma 1.1.1** (Eq. 2.23 in [T<sup>+</sup>10]). *A system made of multiple harmonic oscillators is described by their annihilation operators,*

$$\hat{a}_k, \quad k = 1, \dots, n. \quad (1.1.1)$$

*They satisfy the canonical commutation relation*

$$[\hat{a}_k, \hat{a}_l^\dagger] = \delta_{kl}. \quad (1.1.2)$$

**Lemma 1.1.2** (Eq. 2.27 in [T<sup>+</sup>10]). *The corresponding quadrature operators of a system made of multiple harmonic oscillators satisfy the canonical commutation relation*

$$[\hat{q}_k, \hat{p}_l] = \frac{i}{2} \delta_{kl}. \quad (1.1.3)$$

**Lemma 1.1.3.** *We describe the phase space of the single bosonic modes with the vector*

$$\mathcal{R} = (\hat{q}, \hat{p})^T, \quad (1.1.4)$$

so  $\mathcal{R}_1 = \hat{q}$  and  $\mathcal{R}_2 = \hat{p}$ . Then we have

$$[\mathcal{R}_k, \mathcal{R}_l] = \frac{i}{2} \omega_{kl}, \quad (1.1.5)$$

where

$$\omega = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \quad (1.1.6)$$

is the symplectic form of the phase space of single bosonic mode.

**Lemma 1.1.4.** We describe the phase space of the  $n$  bosonic modes with the vector

$$\mathcal{R} = (\hat{q}^1, \hat{p}^1, \dots, \hat{q}^n, \hat{p}^n)^T. \quad (1.1.7)$$

We denote  $\mathcal{R}_1^i = \hat{q}^i$ ,  $\mathcal{R}_2^i = \hat{p}^i$ , where  $i$  denotes  $i_{th}$  mode. Then we have

$$[\mathcal{R}_k^m, \mathcal{R}_l^n] = \frac{i}{2} \omega_{kl} \delta_{mn}, \quad (1.1.8)$$

where

$$\omega = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \quad (1.1.9)$$

is the symplectic form of the phase space of single bosonic mode. Then the symplectic of  $n$  bosonic modes is

$$\Omega = \bigoplus_{k=1}^n \omega, \quad (1.1.10)$$

where  $\bigoplus$  denotes the direct sum.

### 1.1.3 Introduction

### 1.1.4 Problem formulation

### 1.1.5 Protocol

### 1.1.6 Protocol Details

### 1.1.7 Performance evaluation

## 1.2 Joint Measurement of TFE via SFG [LH20]

### 1.2.1 Mathematical preliminaries

**Proposition 1.2.1** (The Fourier Transform of 1).

### 1.2.2 Physics preliminaries

**Definition 1.2.1** (Sum Frequency Generation (SFG)). *The SFG process in a  $\chi^{(2)}$  non-linear medium could be modeled as the following evolution operator:*

$$V = I + \varepsilon \left( \int d\omega_p d\omega_s d\omega_i a_p^\dagger(\omega_p) a_s(\omega_s) a_i(\omega_i) \delta(\omega_p - \omega_s - \omega_i) - H.C. \right), \quad (1.2.1)$$

where photons in the signal mode  $a_s(\omega_s)$  and ideler mode  $a_i(\omega_i)$  are annihilated to generate photons in the pump mode  $a_p(\omega_p)$  and  $\varepsilon$  characterizes the interaction strength. The SPDC process is the time-reversal process of SFG, which can also be described by the same evolution operator.

**Definition 1.2.2** (SPDC).

**Definition 1.2.3** (Evolution operator).

**Theorem 1** (Schmidt' decomposition [NC19]).

**Definition 1.2.4** (Schmidt's number [NC19]).

**Definition 1.2.5** (Heisenberg picture).

**Proposition 1.2.2** (The relationship between annihilation operator and frequency).

**Definition 1.2.6** (Time Correlation).

**Definition 1.2.7** (Frequency/Spectral/Energy Correlation).

### 1.2.3 Introduction

SPDC is widely adopted to generate TFE photons. SFG can be used to perform TFE joint-measurement.

[SZQ: 2023.06.03: TFE joint-measurement is just entangled measurement. SPDC is similart to unitary evolution for state preparation, in particular, the time-frequency entangled states. Thus, the reversed process SPDC, i.e., SFG process, is the entangled measurement. This is analogy to bell state preparation and bell measurements.]

### 1.2.4 Problem formulation

The frequency sum and time difference of two photons could be simultaneously measured through the sum-frequency generation process.

### 1.2.5 Protocol

Given the close connection between the spontaneous parametric down-conversion (SPDC) process and time-frequency entanglement (TFE), it's natural to utilize the time-reversal of the SPDC process, i.e., sum frequency generation (SFG) to obtain a TFE joint measurement based protocol.

**Definition 1.2.8** (Frequency sum (FS) operator). *The frequency sum operator  $P_{\delta\omega}(\omega)$  that selects states with the frequency sum  $\omega_s + \omega_i$  of the signal and idler photon being around  $\omega$  within uncertainty  $\delta\omega$  is defined as:*

$$P_{\delta\omega}(\omega) = \int \int d\omega_s d\omega_i a_s^\dagger(\omega_s) a_i^\dagger(\omega_i) a_s(\omega_s) a_i(\omega_i) \text{Gate} \left( \frac{\omega - \omega_s - \omega_i}{\delta\omega} \right), \quad (1.2.2)$$

where  $\text{Gate}(x) = 1$  for  $|x| \leq 1/2$  and  $\text{Gate}(x) = 0$  otherwise.

**Lemma 1.2.1** (Frequency sum operator is a projection operator). *The frequency sum operator  $P_{\delta\omega}(\omega)$  is a projection operator satisfying*

$$P_{\delta\omega}(\omega)^2 = P_{\delta\omega}(\omega). \quad (1.2.3)$$

**Definition 1.2.9** (Time difference (TD) operator). *The time difference operator  $P_{\delta t}(t)$  that selects states with the time difference  $t_s - t_i$  of the signal and idler photon being around  $t$  within uncertainty  $\delta t$  is defined as:*

$$P_{\delta t}(t) = \int \int dt_s dt_i \tilde{a}_s^\dagger(t_s) \tilde{a}_i^\dagger(t_i) \tilde{a}_s(t_s) \tilde{a}_i(t_i) \text{Gate} \left( \frac{t_s - t_i - t}{\delta t} \right), \quad (1.2.4)$$

where  $\text{Gate}(x) = 1$  for  $|x| \leq 1/2$  and  $\text{Gate}(x) = 0$  otherwise and

$$\tilde{a}_x = \frac{1}{\sqrt{2\pi}} \int d\omega \exp(-i\omega t) a_x(\omega) \quad (1.2.5)$$

**Lemma 1.2.2** (Time difference operator is a projection operator). *The time difference operator  $P_{\delta t}(t)$  is a projection operator satisfying*

$$P_{\delta t}(t)^2 = P_{\delta t}(t). \quad (1.2.6)$$

**Definition 1.2.10** (Joint projection operator). *The joint projection operator of the time difference and frequency sum is defined as*

$$P_{\delta\omega, \delta t}(\omega, t) = P_{\delta\omega}(\omega) P_{\delta t}(t), \quad (1.2.7)$$

which means selecting states of which the time difference between the signal and idler photon  $t_s - t_i$  is around  $t$  within uncertainty  $\delta t$  and frequency sum of the signal and idler photon being around  $\omega$  within uncertainty  $\delta\omega$ , simultaneously.

**Lemma 1.2.3** (Commutation relationship between frequency-time operators). *We have the commutation relationship*

$$[P_{\delta_\omega}(\omega), P_{\delta_t}(t)] = 0. \quad (1.2.8)$$

**Definition 1.2.11** (TD an FS probability density operator (PDF)). *We define*

$$P(\omega, t) = \lim_{\delta_t \rightarrow 0, \delta_\omega \rightarrow 0} \frac{1}{\delta_\omega \delta_t} P_{\delta_\omega}(\omega) P_{\delta_t}(t). \quad (1.2.9)$$

**Lemma 1.2.4.** *We have*

$$P(\omega, t) = \frac{1}{2\pi} B_p^\dagger B_p, \quad (1.2.10)$$

where

$$B_p = \int \int d\omega_s d\omega_i \delta(\omega_s + \omega_i - \omega) \exp[i\omega_i t] a_s(\omega_s) a_i(\omega_i). \quad (1.2.11)$$

**Lemma 1.2.5** (Connection between TD and FS PDF and SFG Process). *We have*

$$B^\dagger B = 2\pi P(\omega, 0). \quad (1.2.12)$$

This lemma did not show the SFG process

**Lemma 1.2.6** (Discrete sum of evolution operator of SFG process). *The discrete sum of the evolution operator of SFG process can be obtained by a **two-step Schmidt decomposition** as:*

$$v = I + \varepsilon \sum_m \left( \sqrt{\lambda_m^{(1)}} A_m^\dagger B_m - H.C. \right), \quad (1.2.13)$$

where

$$B_m = \sum_n \sqrt{\lambda_{m,n}^{(2)}} F_{m,n} G_{m,n}, \quad (1.2.14)$$

$$A_m = \int d\omega \psi_{A,m}(\omega) a_p(\omega), \quad (1.2.15)$$

$$B_m = \int d\omega_s d\omega_i \psi_{B,m}(\omega_s, \omega_i) a_s(\omega_s) a_i(\omega_i), \quad (1.2.16)$$

$$F_{m,n} = \int d\omega \psi_{F,m,n}(\omega) a_s(\omega), \quad (1.2.17)$$

$$G_{m,n} = \int d\omega \psi_{G,m,n}(\omega) a_i(\omega). \quad (1.2.18)$$

**Lemma 1.2.7** (Non-uniqueness of the first step Schmidt Decomposition). *If the function  $f_0$  can be written in the following form:*

$$\delta(\omega_p - \omega_s - \omega_i) f_0(\omega_p - \omega_s - \omega_i) = \delta(\omega_p - \omega_s - \Omega_i) f\left(\frac{\omega_s - \Omega_i}{\sqrt{2}}\right), \quad (1.2.19)$$

*then the first step Schmidt decomposition in the main text is not unique.*

**Lemma 1.2.8** (An useful commutation relationship). *An useful commutation relationship:*

$$[B_{m'}, B_{m''}^\dagger] = \delta_{m'm''} + \int d\omega'_s d\omega_i d\omega''_s \psi_{B,m''}^*(\omega'_s, \omega_i) a_s^\dagger(\omega''_s) a_s(\omega'_s) \quad (1.2.20)$$

$$+ \int d\omega_s d\omega'_i d\omega''_i \psi_{B,m''}^*(\omega_s, \omega'_i) \psi_{B,m'} a_i^\dagger(\omega''_i) a_i(\omega'_i). \quad (1.2.21)$$

**Corollary 1.2.1.** *By (1.2.8), we have*

$$[B_{m'}, B_{m''}^\dagger] = \delta_{m'm''} |0\rangle. \quad (1.2.22)$$

**Lemma 1.2.9.** *We have the commutation relation between time difference projection operator and frequency sum projection operator:*

$$[P_{\delta\omega}(\omega), P_{\delta t}(t)] = 0. \quad (1.2.23)$$

**Lemma 1.2.10.** *The frequency spectrum  $S(\omega)$  of the generated pump photon is given by the expectation value of the spectral density operator  $a_p^\dagger(\omega_p) a_p(\omega_p)$*

$$S(\omega_p) = \frac{\epsilon^2 \exp \left[ \frac{1}{8} \left( -4\Delta t^2 \sigma_-^2 - \frac{\Delta\omega^2}{\sigma_-^2} - \frac{4(\Delta\omega + \omega_0 - \Omega_p)^2}{\sigma_+^2} \right) \right]}{2\sqrt{\pi}\sigma_+}. \quad (1.2.24)$$

**Definition 1.2.12.** *The discrete mode operator  $F_{m,n}^{(b)}$  for the noise photons is defined as*

$$F_{m,n}^{(b)} = \int d\omega \psi_{F_{m,n}}(\omega) a_s^{(b)}(\omega). \quad (1.2.25)$$

**Definition 1.2.13.** *The virtual beam-splitter is modeledd as the following unitary transform:*

$$U_{loss} = \Pi_n \exp \left[ i \arccos(\eta) (F_{0,n}^\dagger F_{0,n}^{(b)} + H.C.) \right]. \quad (1.2.26)$$

**Definition 1.2.14.** *We use a density matrix  $\rho_b$  that satisfies the following conditions to describe the noise photons:*

$$\text{Tr}[F_{0,n''}^{(b)\dagger} F_{0,n'}^{(b)} \rho_b] = \delta_{n',n''} \mu_b, \quad (1.2.27)$$

$$\text{Tr}[F_{0,n'}^{(b)} \rho_b] = 0 \quad (1.2.28)$$

**Definition 1.2.15.** *The signal and idler photon pair source is described by the biphoton state  $|pair\rangle$ :*

$$|pair\rangle = B_0^\dagger |0\rangle \quad (1.2.29)$$

$$= \sum_n \sqrt{\lambda_{0,n}^{(2)}} F_{0,n}^\dagger G_{0,n} |0\rangle. \quad (1.2.30)$$

**Definition 1.2.16.** *The unitary transform of the SFG process is given by:*

$$V = I + \epsilon \sum_m \left[ \sqrt{\lambda_m^{(1)}} A_m^\dagger B_m - H.C. \right]. \quad (1.2.31)$$

**Definition 1.2.17.** *In the Heisenberg picture, the photon number operator of the generated pump photon in each pump mode  $A_m$  after the beam-splitter transform and the SFG process is given by:*

$$U_{loss}^\dagger V^\dagger A_m^\dagger V U_{loss}. \quad (1.2.32)$$

**Proposition 1.2.3.** *When the transmission of the signal photon is perfect ( $\eta = 1$ ), the pump photon can only generate in mode  $A_0 (m = 0)$ .*

**Lemma 1.2.11.** *We have*

$$\langle U_{loss}^\dagger V^\dagger A_0^\dagger A_0 V | U_{loss} \rangle = \epsilon^2 \lambda_0^{(1)} (\eta + \mu_b \sum_n \lambda_{0,n}^2). \quad (1.2.33)$$

**Definition 1.2.18.** *The generated SPDC state is given by:*

$$V = |0\rangle - \epsilon \sqrt{\lambda_0^{(1)}} \alpha B_0^\dagger |0\rangle. \quad (1.2.34)$$

**Lemma 1.2.12.** *The joint density operator of the noise-idler state  $\rho_j$  is given by the tensor product of  $\rho_i$  and  $\rho_b$ :*

$$\rho_j = \rho_i \otimes \rho_b \quad (1.2.35)$$

$$= \mu_b \int \int d\omega_s d\omega'_s \int \int d\omega'_i d\omega''_i \phi_0^*(\omega'_s, \omega'_i) \phi_0(\omega'_s, \omega''_i) a_i^\dagger(\omega'_i) a_s^\dagger(\omega_s) |0\rangle \langle 0| a_i(\omega''_i) a_s(\omega_s). \quad (1.2.36)$$

**Lemma 1.2.13.** *The spectral density  $S(\omega)$  of the upconverted photons is*

$$S(\omega) = \frac{\epsilon^2 \mu_b \exp \left[ -\frac{(\omega - \omega_0)^2}{8\sigma_-^2 - 2\sigma_+^2} \right]}{\sqrt{\pi} \sqrt{4\sigma_-^2 + \sigma_+^2}}. \quad (1.2.37)$$

**Theorem 2.** *The error exponent of the classical Chernoff bound of the TFE QI protocol is given by  $C_{QI}$  [NS09]:*

$$C_{QI} = -\log \min_{s \in [0,1]} \left\{ \sum_{b \in \{0,1\}} p_0(b)^s p_1(b)^{(1-s)} \right\}. \quad (1.2.38)$$

## 1.2.6 Performance evaluation

**Lemma 1.2.14.**

$$(1.2.39)$$

### 1.2.7 Ideas

## 1.3 Optimum Mixed-State Discrimination for Noisy Entanglement-Enhanced Sensing [ZZS17]

### 1.3.1 Mathematical preliminaries

### 1.3.2 Physics preliminaries

### 1.3.3 Introduction

### 1.3.4 Problem formulation

### 1.3.5 Protocol

### 1.3.6 Protocol Details

### 1.3.7 Performance evaluation

**Lemma 1.3.1.**

(1.3.1)

### 1.3.8 Ideas

## 1.4 Quantum Estimation Methods for Quantum Illumination [SLHGR<sup>+</sup>17]

### 1.4.1 Mathematical preliminaries

**Definition 1.4.1** (Quantum Fisher Information (QFI)). *Define a quantum state  $\rho_\eta$  parameterized by  $\eta$ . Then the QFI for  $\rho_\eta$  is*

$$H := 2 \sum_{mn} \frac{|\langle \phi_m | (\partial_\eta \rho_\eta|_{\eta=0} | \phi_n \rangle)|^2}{\lambda_m + \lambda_n}, \quad (1.4.1)$$

where  $\lambda_n$  is the eigenvalue of  $\rho_{\eta=0}$  corresponding to the eigenstate  $|\phi_n\rangle$ , and the derivative is evaluated as  $\eta = 0$ .

**Definition 1.4.2** (Cramér-Rao Bound (CROB) in Eq. (2) [SLHGR<sup>+</sup>17]). *The CROB asserts that the limits on the achievable precision of an unbiased estimator  $\tilde{\eta}$  is*

$$\Delta \tilde{\eta}^2 \geq \frac{1}{MH}, \quad (1.4.2)$$



where  $H$  is the QFI (??) of  $\rho_\eta$  and  $M$  is the number of copies.

**Theorem 3** (Cramér-Chernoff theorem [Hay02]).

**Definition 1.4.3** (Matrix exponential from wiki). Let  $X$  be an  $n \otimes n$  real or complex matrix. The exponential of  $X$ , denoted by  $e^X$  or  $\exp[X]$ , is the  $n \times n$  matrix given by the power series

$$e^X = \sum_{k=0}^{\infty} \frac{1}{k!} X^k, \quad (1.4.3)$$

where  $X^0$  is defined to be the identity matrix  $\text{id}$  with the same dimensions as  $X$ . The series always converges, so the exponential of  $X$  is well-defined.

**Lemma 1.4.1** (Properties of matrix exponential). Let  $X$  and  $Y$  be  $n \times n$  complex matrices and let  $a$  and  $b$  be arbitrary complex numbers. We denote the  $n \times n$  identity matrix by  $\text{id}$  and zero matrix by  $0$ . The matrix exponential satisfies the following properties:

**Elementary properties:**

- $\exp[0] = \text{id}$
- $\exp[X^\dagger] = \exp[X]^\dagger$
- If  $[X, Y] = 0$ , then  $\exp[X] \exp[Y] = \exp[X + Y]$ .

**The exponential map:** Define the map

$$t \mapsto \exp[tX], \quad t \in \mathbb{R}. \quad (1.4.4)$$

We have

•

$$\frac{d}{dt} e^{tX} = X e^{tX} = e^{tX} X. \quad (1.4.5)$$

**Lemma 1.4.2** (Geometric series). For  $|x| < 1$ , we have

$$\sum_{n=0}^{\infty} x^n = \frac{1}{1-x}. \quad (1.4.6)$$

**Lemma 1.4.3** (Generalized geometric series).

### 1.4.2 Physics preliminaries

#### 1.4.3 Introduction

#### 1.4.4 Problem formulation

The quantum illumination problem is modeled as a **Reflectivity Estimation Problem** as the following.

Let us consider a general bipartite pure state representation of signal-idler system written in the Schmidt decomposition form

$$|\psi\rangle_{SI} = \sum_{\alpha} \sqrt{p_{\alpha}} |w_{\alpha}\rangle_S |v_{\alpha}\rangle_I. \quad (1.4.7)$$

In the Quantum Illumination (QI) protocol, the signal modes of  $M$  copies of  $|\psi\rangle_{SI}$  are sent to the target region embedded in a bright thermal noise, in which there could possibly be an object. We receive  $M$  copies of the state

$$\rho_{\eta} = \text{Tr}_S \left[ U_{\eta} \psi \otimes \rho_B U_{\eta}^{\dagger} \right], \quad (1.4.8)$$

where

$$U_{\eta} := \exp[\sin^{-1}(\eta) - s^{\dagger}b - sb^{\dagger}] \simeq \exp[\eta(s^{\dagger}b - sb^{\dagger})] \quad (1.4.9)$$

is the signal-object interaction, modeled as a beamsplitter with amplitude reflectivity  $\eta \ll 1$ , and

$$\rho_B := \sum_n \frac{N_B^n}{(1 + N_B)^{1+n}} |n\rangle\langle n| \quad (1.4.10)$$

is the thermal state with mean photon number  $N_B$ .

In this framework, at the receiver side,  $\eta = 0$  corresponds to the absence of the object in the target region. We note  $\eta$  is unknown parameter. So we measurement  $\rho_{\eta}$  many times and get the estimation  $\tilde{\eta}$ . We then use  $\hat{\eta}$  to determine whether is target is absent or present. This lie in the Estimation problem and state discrimination problem!

#### 1.4.5 Protocol

The quantum illumination protocol has been presented the last subsection. Here, we focus on the estimation protocol.

The main idea of the estimation protocol is to use QFI (??) and CROB (??) for estimating  $\eta$ .

**Lemma 1.4.4.** *If a mode  $s$  sent into a beamsplitter that has a thermal state input  $b$  in the other port, then  $U_\eta = \exp[\eta(s^\dagger b - sb^\dagger)]$  is indeed the unitary transformation.*

*Proof.* By the properties of matrix exponential 1.4.3, we have

$$U_\eta^\dagger = \exp[\eta(s^\dagger b - sb^\dagger)]^\dagger \quad (1.4.11)$$

$$= \exp[\eta(b^\dagger s - bs^\dagger)]. \quad (1.4.12)$$

Then we have

$$U_\eta U_\eta^\dagger = \exp[\eta(s^\dagger b - sb^\dagger + b^\dagger s - bs^\dagger)] \quad (1.4.13)$$

$$= \exp[\eta([s^\dagger, b] + [b^\dagger, s])] \quad (1.4.14)$$

By lemma 1.1.3

$$= \exp[\eta 0] \quad (1.4.15)$$

$$= \text{id}. \quad (1.4.16)$$

□

**Lemma 1.4.5.** *The state  $\rho_\eta = \text{Tr}_S [U_\eta |\psi\rangle\langle\psi| \otimes \rho_B U_\eta^\dagger]$ , where  $U_\eta \simeq \exp[\eta(s^\dagger b - sb^\dagger)]$ .*

*We have that the derivative computed at  $\eta = 0$  is:*

$$\partial_\eta \rho_\eta = \text{Tr}_S [s^\dagger b - sb^\dagger, |\psi\rangle\langle\psi| \otimes \rho_B] = \sum_{\alpha\alpha'} \sqrt{p_\alpha p_{\alpha'}} |v_\alpha\rangle\langle v_\alpha| \otimes [\langle w_{\alpha'} | s^\dagger | w_\alpha \rangle b - \langle w_{\alpha'} | s | w_\alpha \rangle b^\dagger, \rho_B]. \quad (1.4.17)$$

*Proof.*

$$\partial_\eta \rho_\eta|_{\eta=0} = \text{Tr}_S [\partial_\eta U_\eta |\psi\rangle\langle\psi| \otimes \rho_B U_\eta^\dagger] |_{\eta=0} \quad (1.4.18)$$

$$= \text{Tr}_S [\partial_\eta \exp[\eta(s^\dagger b - sb^\dagger)] |\psi\rangle\langle\psi| \otimes \rho_B \exp[\eta(b^\dagger s - bs^\dagger)]] |_{\eta=0} \quad (1.4.19)$$

$$= \text{Tr}_S [(s^\dagger b - sb^\dagger) \exp[\eta(s^\dagger b - sb^\dagger)] |\psi\rangle\langle\psi| \otimes \rho_B \exp[\eta(b^\dagger s - bs^\dagger)] + \quad (1.4.20)$$

$$\exp[\eta(s^\dagger b - sb^\dagger)] |\psi\rangle\langle\psi| \otimes \rho_B \exp[\eta(b^\dagger s - bs^\dagger)] (b^\dagger s - bs^\dagger)] |_{\eta=0} \quad (1.4.21)$$

$$= \text{Tr}_S [(s^\dagger b - sb^\dagger) |\psi\rangle\langle\psi| \otimes \rho_B + |\psi\rangle\langle\psi| \otimes \rho_B (b^\dagger s - bs^\dagger)] \quad (1.4.22)$$

$$= \text{Tr}_S [(s^\dagger b - sb^\dagger) |\psi\rangle\langle\psi| \otimes \rho_B - |\psi\rangle\langle\psi| \otimes \rho_B (bs^\dagger - b^\dagger s)] \quad (1.4.23)$$

By lemma 1.1.3

$$= \text{Tr}_S [(s^\dagger b - sb^\dagger) |\psi\rangle\langle\psi| \otimes \rho_B - |\psi\rangle\langle\psi| \otimes \rho_B (s^\dagger b - sb^\dagger)] \quad (1.4.24)$$

$$= \text{Tr}_S [s^\dagger b - sb^\dagger, |\psi\rangle\langle\psi| \otimes \rho_B]. \quad (1.4.25)$$

Then we continue the derivation as

$$\text{Tr}_S \left[ |\psi\rangle\langle\psi| \otimes \rho_B (s^\dagger b - s b^\dagger) \right] \quad (1.4.26)$$

$$= \text{Tr} \left[ |\psi\rangle\langle\psi| s^\dagger \otimes \rho_B b - |\psi\rangle\langle\psi| s \otimes \rho_B b^\dagger \right] \quad (1.4.27)$$

$$= \sum_k \langle w_k | \psi \rangle \langle \psi | s^\dagger \otimes \rho_B b | w_k \rangle - \sum_k \langle w_k | \psi \rangle \langle \psi | s \otimes \rho_B b | w_k \rangle \quad (1.4.28)$$

$$= \sum_{kij} \sqrt{p_i p_j} \langle w_k | \otimes I | w_i \rangle \langle w_j | \otimes | v_i \rangle \langle v_j | s^\dagger \otimes \text{id} | w_k \rangle \otimes \text{id} \otimes \rho_B b - \quad (1.4.29)$$

$$\sum_{kij} \sqrt{p_i p_j} \langle w_k | \otimes \text{id} | w_i \rangle \langle w_j | \otimes | v_i \rangle \langle v_j | s \otimes \text{id} | w_k \rangle \otimes \text{id} \otimes \rho_B b^\dagger \quad (1.4.30)$$

$$= \sum_{kij} \sqrt{p_i p_j} \langle w_k | w_i \rangle \langle w_j | s^\dagger | w_k \rangle \otimes | v_i \rangle \langle v_j | \otimes \rho_B b - \quad (1.4.31)$$

$$\sum_{kij} \sqrt{p_i p_j} \langle w_k | w_i \rangle \langle w_j | s | w_k \rangle \otimes | v_i \rangle \langle v_j | \otimes \rho_B b^\dagger \quad (1.4.32)$$

$$= \sum_{ij} \sqrt{p_i p_j} \langle w_j | s^\dagger | w_i \rangle | v_i \rangle \langle v_j | \otimes \rho_B b - \sum_{ij} \sqrt{p_i p_j} \langle w_j | s | w_i \rangle | v_i \rangle \langle v_j | \otimes \rho_B b^\dagger \quad (1.4.33)$$

$$= \sum_{ij} \sqrt{p_i p_j} | v_i \rangle \langle v_j | \otimes \rho_B \langle w_j | s^\dagger | w_i \rangle b - \sum_{ij} \sqrt{p_i p_j} | v_i \rangle \langle v_j | \otimes \rho_B \langle w_j | s | w_i \rangle b^\dagger \quad (1.4.34)$$

$$= \sum_{ij} \sqrt{p_i p_j} | v_i \rangle \langle v_j | \otimes \rho_B \left[ \langle w_j | s | w_i \rangle b^\dagger - \langle w_j | s^\dagger | w_i \rangle b \right]. \quad (1.4.35)$$

Similarly, we can find that

$$\text{Tr}_S \left[ (s^\dagger b - s b^\dagger) |\psi\rangle\langle\psi| \otimes \rho_B \right] \quad (1.4.36)$$

$$= \sum_{ij} \left[ \langle w_j | s^\dagger | w_i \rangle b - \langle w_j | s | w_i \rangle b^\dagger \right] \sqrt{p_i p_j} | v_i \rangle \langle v_j | \otimes \rho_B \quad (1.4.37)$$

$$= \sum_{ij} \sqrt{p_i p_j} | v_i \rangle \langle v_j | \otimes \left[ \langle w_j | s^\dagger | w_i \rangle b - \langle w_j | s | w_i \rangle b^\dagger \right] \rho_B. \quad (1.4.38)$$

In summary, we have

$$\sum_{ij} \sqrt{p_i p_j} | v_i \rangle \langle v_j | \otimes \left[ \left( \langle w_j | s^\dagger | w_i \rangle b - \langle w_j | s | w_i \rangle b^\dagger \right) \rho_B - \left( \langle w_j | s | w_i \rangle b^\dagger - \langle w_j | s^\dagger | w_i \rangle b \right) \rho_B \right] \quad (1.4.39)$$

$$= \sum_{ij} \sqrt{p_i p_j} | v_i \rangle \langle v_j | \otimes \left[ \left( \langle w_j | s^\dagger | w_i \rangle b - \langle w_j | s | w_i \rangle b^\dagger \right), \rho_B \right]. \quad (1.4.40)$$

□

**Lemma 1.4.6** (The QFI for the returned state  $\rho_\eta$  in Eq. (3) [SLHGR<sup>+</sup>17]). *The QFI for  $\rho_\eta$  is*

$$H := \frac{4}{1 + N_B} \sum_{\alpha\alpha'} \frac{p_\alpha p_{\alpha'}}{p_{\alpha'} + p_\alpha \frac{N_B}{1+N_B}} |\langle w_{\alpha'} | s | w_\alpha \rangle|^2. \quad (1.4.41)$$

*Proof.* □

We choose the observable that optimize the QFI of the returned state (??) and measure  $\rho_\eta$ , the outcome is  $o_i$ . Then by the strong law of large numbers, we perform the sample mean as

$$\tilde{\eta} = \frac{1}{M} \sum_{i=1}^M o_i. \quad (1.4.42)$$

---

**Algorithm 1** QI estimation protocol

---

**Input:**  $|\psi_{SI}\rangle$ : the signal-idler state,

$\eta$ : the reflectivity parameter,

$\rho_B$ : the thermal state

$M$ : the number of copies of  $|\psi_{SI}\rangle$

**Output:** Estimation of the reflectivity  $\tilde{\eta}$ .

- 1: **for**  $i = 1, \dots, M$  **do**
  - 2:   Sent the signal mode of  $\psi_{SI}^{(1)}$  to the target region,
  - 3:   Rceive the state  $\rho_\eta^{(1)}$ ,
  - 4:   **Choose the observable that optimize the QFI,**
  - 5:   Measure the observable and get the outcome  $o_i$ .
  - 6: **end for**
  - 7: Calculate  $\tilde{\eta}$ (1.4.41).
  - 8: Output the estimated reflectivity parameter  $\tilde{\eta}$ .
- 

After that, we use the estimation  $\tilde{\eta}$  for state discrimination.

---

**Algorithm 2** QI discrimination protocol

---

**Input:**  $\tilde{\eta}$ : the estimation reflectivity,

$\xi$ : the scaling parameter.

**Output:** Absent or present.

- 1: **if**  $\tilde{\eta} > \xi\eta$  **then**
  - 2:   Output present.
  - 3: **else if**  $\tilde{\eta} \leq \xi\eta$  **then**
  - 4:   Output absent.
  - 5: **else**
  - 6:   Output 'Wrong input!'
  - 7: **end if**
-

### 1.4.6 Protocol Details

### 1.4.7 Performance evaluation

**Theorem 4** (Type I-II error probabilities Theorem in [SLHGR<sup>+</sup>17]). *Let  $|\psi\rangle_{SI} = \sqrt{p_\alpha}|n\rangle_S|v_n\rangle_I$  be the Schmidt decomposition of the signal-idler state, and denote  $\rho_S$  the state of the signal. Then  $P_{I,II} \sim \exp\left(-\frac{\eta_{I,II}^2 HM}{2}\right)$  provided that  $\exists C > 0$  s.t.  $\langle s^k s^{\dagger k} \rangle_{\rho_S} \leq k! C^k, \forall k \in \mathcal{N}$ .*

*Proof.*

□

### 1.4.8 Exapmles

**Example 1.4.1** (Gaussian states).

**Example 1.4.2** (Schrödinger's cat state).

### 1.4.9 Main ideas summarized

This subsection summarizes the main ideas in this paper.

Choose the observable that optimize the QFI.

### 1.4.10 Ideas

How many copies of  $\psi_{SI}$  is needed to achieve a given confidence interval? This is a problem about the quantum resources?

# Bibliography

- [Hay02] Masahito Hayashi. Two quantum analogues of fisher information from a large deviation viewpoint of quantum estimation. *Journal of Physics A: Mathematical and General*, 35(36):7689, 2002.
- [LH20] Han Liu and Amr S Helmy. Joint measurement of time–frequency entanglement via sum frequency generation. *npj Quantum Information*, 6(1):66, 2020.
- [NC19] Michael A. Nielsen and Isaac L. Chuang. *Quantum computation and quantum information*. Cambridge University Press, 2019.
- [NS09] Michael Nussbaum and Arleta Szkoła. The chernoff lower bound for symmetric quantum hypothesis testing. 2009.
- [SLHGR<sup>+</sup>17] Mikel Sanz, U Las Heras, Juan José García-Ripoll, Enrique Solano, and R Di Candia. Quantum estimation methods for quantum illumination. *Physical review letters*, 118(7):070803, 2017.
- [T<sup>+</sup>10] Si Hui Tan et al. *Quantum state discrimination with bosonic channels and gaussian states*. PhD thesis, Massachusetts Institute of Technology, 2010.
- [ZZS17] Quntao Zhuang, Zheshen Zhang, and Jeffrey H Shapiro. Optimum mixed-state discrimination for noisy entanglement-enhanced sensing. *Physical review letters*, 118(4):040801, 2017.