

Tutorial of ST5215

AY2020/2021 Semester 1

10 Nov 2020

Exercise 1. Let (X_1, \dots, X_n) be a random sample from the exponential distribution on (a, ∞) with scale parameter θ , where $a \in \mathcal{R}$ and $\theta > 0$ are unknown. Show that $T = (X_{(1)}, \sum_{i=1}^n X_i - nX_{(1)})$ is a complete statistic.
Hint: Use the Rényi representation (see the reading material “Representation_BlitzsteinMorris.pdf” on LumiNUS).

Exercise 2. Consider a linear model in matrix form $X_{n \times 1} = Z_{n \times p} \beta_{p \times 1} + \epsilon_{n \times 1}$ with $p \leq n$ and with the assumption that $\epsilon \sim N(\mathbf{0}_n, \sigma^2 \mathbf{I}_n)$. Show that if each coordinate of β is estimable, then the rank of Z is p .

Exercise 3. (James-Stein estimator) Suppose X is a p -random vector from $N(\theta, \mathbf{I}_p)$ with an unknown $\theta \in \mathcal{R}^p$. Consider the squared loss function for estimating θ :

$$L(\theta, a) = \|a - \theta\|^2 = \sum_{i=1}^p (a_i - \theta_i)^2,$$

where a_i and θ_i are the i th coordinates of the estimator and the estimand. Show that for any $p \geq 3$, the risk of the following estimator

$$\hat{\theta} = \left(1 - \frac{(p-2)}{\|X\|^2}\right) X$$

is strictly smaller than X .

Can you extend this result to the case where $X \sim N(\theta, D)$ with some known $p \times p$ positive definite matrix D ?

Hint: Use the Stein’s identity (Ex 1 in Tutorial 6).