

PatientID: HIVDR-1750-23

Sebuttemba 27, 2023

## Color Code

■ HR: High-Level Resistance    ■ PLR: Potential Low-Level Resistance  
■ LR: Low-Level Resistance    ■ IR: Intermediate Resistance  
■ S: Susceptible

DRUG.CLASS	DRUG	RESISTANCE.PROFILE	DRMS.above.20.percent.prevalence
PI	ATV	S	;L33F
	DRV	S	
	FPV	PLR	
	IDV	S	
	LPV	S	
	NFV	PLR	
	SQV	S	
	TPV	PLR	
NRTI	ABC	S	
	AZT	S	
	D4T	S	
	DDI	S	
	FTC	S	
	LMV	S	
	TDF	S	
NNRTI	DOR	S	
	EFV	S	
	ETR	S	
	NVP	LR	
	RPV	S	
INSTI	BIC	S	
	CAB	S	
	DTG	S	
	EVG	PLR	
	RAL	PLR	

## Appendix

### Drug abbreviations in full

DRUG.CLASS	ABBREVIATION	DRUG.NAME
<b>PI</b>	ATV	Atazanavir
	DRV	Darunavir
	FPV	Fosamprenavir
	IDV	Indinavir
	LPV	Lopinavir
	NFV	Nelfinavir
	SQV	Saquinavir
	TPV	Tipranavir
<b>NRTI</b>	ABC	Abacavir
	AZT	Azidothymidine
	DFT	Stavudine
	DDI	Didanosine
	FTC	Emtricitabine
	LMV	Lamivudine
	TDF	Tenofovir
<b>NNRTI</b>	DOR	Doravirine
	EFV	Efavirenz
	ETR	Etravirine
	NVP	Nevirapine
	RPV	Rilpivirine
<b>INSTI</b>	BIC	Bictegravir
	CAB	Cabotegravir
	DTG	Dolutegravir
	EVG	Elvitegravir
	RAL	Raltegravir

### Comments

DRUG.CLASS	COMMENTS
<b>PI</b>	L33F is a relatively non-polymorphic accessory mutation selected by each of the PIs. In combination with other PI-resistance mutations, it is associated with reduced susceptibility to LPV, ATV, and DRV.
<b>NRTI</b>	
<b>NNRTI</b>	
<b>INSTI</b>	T97A is a polymorphic INSTI-selected mutation that, depending on subtype, occurs in 1% to 5% of viruses from untreated persons. Alone, it has minimal effects on INSTI susceptibility but in combination with other major resistance mutations, it synergistically reduces susceptibility to each of the INSTIs.