

PI Major Mutations:None

PI Accessory Mutations:None

PR Other Mutations:**L10G** • **V11E** • T12V • I13S • G16E • L33V • N37T • R41K • R57K • L63T • I64V • E65D

Protease Inhibitors	
atazanavir/r (ATV/r)	Susceptible
darunavir/r (DRV/r)	Susceptible
fosamprenavir/r (FPV/r)	Susceptible
indinavir/r (IDV/r)	Susceptible
lopinavir/r (LPV/r)	Susceptible
nelfinavir (NFV)	Susceptible
saquinavir/r (SQV/r)	Susceptible
tipranavir/r (TPV/r)	Susceptible

PR comments

Other

- L10F is a common non-polymorphic, PI-selected accessory mutation associated with reduced in vitro susceptibility to LPV and DRV. L10I/V are polymorphic, PI-selected accessory mutations that increase the replication of viruses with other PI-resistance mutations. L10R/Y are rare, non-polymorphic PI-selected mutations. Their effects on PI susceptibility have not been well studied. **L10G** is a highly unusual mutation at this position.
- L33I/V** are minimally polymorphic mutations that do not appear to be selected by PIs or to reduce their susceptibility.

No drug resistance mutations were found for PI.

NRTI Mutations:**K65R** • **S68N** • **Y115F** • **M184V**

NNRTI Mutations:**K103N** • **Y181C** • **G190A**

RT Other Mutations:V35T • T39A • K46Q • K49R • V60I • K101Q • D123E • I135T • I142V • D177E • **T200X** • Q207R • R211K • **P217S** • P225L • E233D • **Δ234** • **H235S** • P236S • **D237*** • **K238*** • V245N • **L246A** • **P247E** • E248K • **D250*** • **S251L** • W252Y • **T253D** • **V254L** • **N255Q** • **D256S** • **I257S** • Q258R • K259N • **L260T** • V261G • **G262Q** • **K263S** • L264I • **N265Y** • **W266Q** • **K275*** • V276I • L279I • **K281N** • **L282A** • **Δ284** • T286S • **K287*** • **V292*** • **I293*** • P294L

Nucleoside Reverse Transcriptase Inhibitors	
abacavir (ABC)	High-Level Resistance
zidovudine (AZT)	Susceptible
stavudine (D4T)	Intermediate Resistance
didanosine (DDI)	High-Level Resistance
emtricitabine (FTC)	High-Level Resistance
lamivudine (3TC)	High-Level Resistance
tenofovir (TDF)	High-Level Resistance

Non-nucleoside Reverse Transcriptase Inhibitors	
doravirine (DOR)	Low-Level Resistance
efavirenz (EFV)	High-Level Resistance
etravirine (ETR)	Intermediate Resistance
nevirapine (NVP)	High-Level Resistance
rilpivirine (RPV)	High-Level Resistance

RT comments

NRTI

- **K65R** confers intermediate reductions in susceptibility to TDF, ABC, and 3TC/FTC. It increases AZT susceptibility. In NRTI-experienced, INSTI-naïve patients with **K65R**, TDF+3TC+DTG is usually highly effective and more effective than AZT/3TC/DTG. However, in patients receiving TDF+3TC+DTG, there is a risk of emergent DTG resistance that does not arise in NRTI-naïve patients receiving TDF+3TC+DTG.
- **Y115F** causes intermediate resistance to ABC and low-level resistance to TDF.
- **M184V/I** cause high-level in vitro resistance to 3TC and FTC and low/intermediate resistance to ABC (3-fold reduced susceptibility). **M184V/I** are not contraindications to continued treatment with 3TC or FTC because they increase susceptibility to AZT and TDF and are associated with clinically significant reductions in HIV-1 replication.

NNRTI

- **K103N** is a non-polymorphic mutation that confers high-level reductions in NVP and EFV susceptibility. It is the most commonly transmitted DRM.
- **Y181C** is a non-polymorphic mutation selected in persons receiving NVP, ETR and RPV. It confers high-level resistance to NVP, intermediate resistance to ETR and RPV, and low-level resistance to EFV. It does not significantly reduce DOR susceptibility.
- **G190A** is a non-polymorphic mutation that causes high-level resistance to NVP and intermediate resistance to EFV. It does not significantly reduce susceptibility to RPV, ETR, or DOR.

Other

- **K101Q** is a relatively non-polymorphic mutation that is weakly selected in persons receiving NVP and EFV. It is of uncertain phenotypic and clinical significance.
- P225H is a non-polymorphic EFV-selected mutation that usually occurs in combination with K103N. The combination of P225H and K103N synergistically reduces NVP, EFV and DOR susceptibility. **P225L** is a highly unusual mutation at this position.
- L234I is a nonpolymorphic mutation selected in persons receiving NVP and EFV. It is also selected in vitro by ETR and DOR. In combination with V106A, it is associated with high-level DOR resistance. Its effect on susceptibility when it occurs alone has not been well characterized. **L234del** is a highly unusual mutation at this position.
- P236L is a rare mutation selected commonly by DLV, which appears to have little if any effect on current NNRTIs. **P236S** is a highly unusual mutation at this position.

Mutation scoring: RT

HIVDB 9.5.1 (2023-11-05)

Drug resistance mutation scores of NRTI:

Download CSV



Rule	ABC ↕	AZT ↕	D4T ↕	DDI ↕	FTC ↕	3TC ↕	TDF ↕
K65R	45	-10	60	60	30	30	50
Y115F	30	0	0	0	0	0	15
Y115F + M184V	15	0	0	0	0	0	5
M184V	15	-10	-10	10	60	60	-10
K65R + S68N	0	0	0	0	0	0	5
Total	105	-20	50	70	90	90	65

Drug resistance mutation scores of NNRTI:

Download CSV



Rule	DOR ↕	EFV ↕	ETR ↕	NVP ↕	RPV ↕
K103N + Y181C	5	0	0	0	0
Y181C	10	30	30	60	45
Y181C + G190A	10	0	10	0	10
K103N	0	60	0	60	0
G190A	0	45	10	60	15
Total	25	135	50	180	70