

PatientID: NC602-1997

Okitobba 06, 2023

### Color Code

■ HR: High-Level Resistance    ■ PLR: Potential Low-Level Resistance  
■ LR: Low-Level Resistance    ■ IR: Intermediate Resistance  
■ S: Susceptible

DRUG.CLASS	DRUG	RESISTANCE.PROFILE	DRMS.above.20.percent.prevalence
PI	ATV	PLR	D30N;F53LF
	DRV	S	
	FPV	S	
	IDV	S	
	LPV	S	
	NFV	HR	
	SQV	LR	
	TPV	S	
NRTI	ABC	IR	D67N;K70R;M184V;K219Q
	AZT	HR	
	D4T	IR	
	DDI	IR	
	FTC	HR	
	LMV	HR	
	TDF	LR	
NNRTI	DOR	S	
	EFV	S	
	ETR	S	
	NVP	S	
	RPV	S	

## Appendix

### Drug abbreviations in full

DRUG.CLASS	ABBREVIATION	DRUG.NAME
<b>PI</b>	ATV	Atazanavir
	DRV	Darunavir
	FPV	Fosamprenavir
	IDV	Indinavir
	LPV	Lopinavir
	NFV	Nelfinavir
	SQV	Saquinavir
	TPV	Tipranavir
<b>NRTI</b>	ABC	Abacavir
	AZT	Azidothymidine
	DFT	Stavudine
	DDI	Didanosine
	FTC	Emtricitabine
	LMV	Lamivudine
	TDF	Tenofovir
<b>NNRTI</b>	DOR	Doravirine
	EFV	Efavirenz
	ETR	Etravirine
	NVP	Nevirapine
	RPV	Rilpivirine
<b>INSTI</b>	BIC	Bictegravir
	CAB	Cabotegravir
	DTG	Dolutegravir
	EVG	Elvitegravir
	RAL	Raltegravir

### Comments

DRUG.CLASS	COMMENTS
<b>PI</b>	D30N is a non-polymorphic mutation NFV-selected mutation that causes high-level resistance to NFV but not to other PIs.
	F53L is a nonpolymorphic accessory mutation selected primarily by SQV, IDV, ATV and LPV. In combination with other mutations, It is associated with reduced susceptibility to ATV and possibly LPV. F53Y is an uncommon nonpolymorphic accessory PI-selected mutation that has not been well studied.
<b>NRTI</b>	D67N is a non-polymorphic TAM associated with low-level resistance to AZT.
	K219E/Q/N/R are accessory TAMs that usually occur in combination with multiple other TAMs.
	K70R is a TAM that confers intermediate resistance to AZT and contributes to reduced ABC and TDF susceptibility in combination with other TAMs.
	M184V/I cause high-level in vitro resistance to 3TC and FTC and low/intermediate resistance to ABC (3-fold reduced susceptibility). M184V/I are not contraindications to continued treatment with 3TC or FTC because they increase susceptibility to AZT and TDF and are associated with clinically significant reductions in HIV-1 replication.
<b>NNRTI</b>	

