

PatientID: HDR97

Okitobba 06, 2023

Color Code

■ HR: High-Level Resistance ■ PLR: Potential Low-Level Resistance
■ LR: Low-Level Resistance ■ IR: Intermediate Resistance
■ S: Susceptible

DRUG.CLASS	DRUG	RESISTANCE.PROFILE	DRMS.above.20.percent.prevalence
PI	ATV	IR	L90M;F53L
	DRV	S	
	FPV	IR	
	IDV	IR	
	LPV	LR	
	NFV	HR	
	SQV	HR	
	TPV	S	
NRTI	ABC	IR	L74I;M184V
	AZT	S	
	D4T	S	
	DDI	HR	
	FTC	HR	
	LMV	HR	
	TDF	S	
NNRTI	DOR	IR	P225H;K103N
	EFV	HR	
	ETR	S	
	NVP	HR	
	RPV	S	
INSTI	BIC	S	
	CAB	S	
	DTG	S	
	EVG	S	
	RAL	S	

Appendix

Drug abbreviations in full

DRUG.CLASS	ABBREVIATION	DRUG.NAME
PI	ATV	Atazanavir
	DRV	Darunavir
	FPV	Fosamprenavir
	IDV	Indinavir
	LPV	Lopinavir
	NFV	Nelfinavir
	SQV	Saquinavir
	TPV	Tipranavir
NRTI	ABC	Abacavir
	AZT	Azidothymidine
	DFT	Stavudine
	DDI	Didanosine
	FTC	Emtricitabine
	LMV	Lamivudine
	TDF	Tenofovir
NNRTI	DOR	Doravirine
	EFV	Efavirenz
	ETR	Etravirine
	NVP	Nevirapine
	RPV	Rilpivirine
INSTI	BIC	Bictegravir
	CAB	Cabotegravir
	DTG	Dolutegravir
	EVG	Elvitegravir
	RAL	Raltegravir

Comments

DRUG.CLASS	COMMENTS
PI	F53L is a nonpolymorphic accessory mutation selected primarily by SQV, IDV, ATV and LPV. In combination with other mutations, It is associated with reduced susceptibility to ATV and possibly LPV. F53Y is an uncommon nonpolymorphic accessory PI-selected mutation that has not been well studied.
	L90M is a non-polymorphic PI-selected mutation that reduces susceptibility to ATV and to a lesser extent LPV.
NRTI	L74V causes intermediate ABC resistance. L74I causes low-level ABC resistance.
	M184V/I cause high-level in vitro resistance to 3TC and FTC and low/intermediate resistance to ABC (3-fold reduced susceptibility). M184V/I are not contraindications to continued treatment with 3TC or FTC because they increase susceptibility to AZT and TDF and are associated with clinically significant reductions in HIV-1 replication.
NNRTI	K103N is a non-polymorphic mutation that confers high-level reductions in NVP and EFV susceptibility. It is the most commonly transmitted DRM.
	P225H is a non-polymorphic EFV-selected mutation that usually occurs in combination with K103N. The combination of P225H and K103N synergistically reduces NVP, EFV and DOR susceptibility.

