

INSTI Major Mutations:

INSTI Accessory Mutations:

IN Other Mutations:

G140GS • Q148QH

None

S17N • I72V • A98AT • L101I • K156KN • V201I • I208L • D286DN

Integrase Strand Transfer Inhibitors	
bictegravir (BIC)	Intermediate Resistance
cabotegravir (CAB)	High-Level Resistance
dolutegravir (DTG)	Intermediate Resistance
elvitegravir (EVG)	High-Level Resistance
raltegravir (RAL)	High-Level Resistance

IN comments

Major

- G140S[A/C] are non-polymorphic mutations that usually occur with Q148 mutations. Alone, they have minimal effects on INSTI susceptibility. However, in combination with Q148 mutations they are associated with high-level resistance to RAL and EVG and intermediate reductions in DTG and BIC susceptibility.
- Q148H[K/R] are nonpolymorphic mutations reported in persons receiving RAL, EVG, CAB, and DTG. They nearly always occur in combination with G140A/S or E138K. In this setting they are associated with near complete resistance to RAL and EVG, high-levels of reduction in CAB susceptibility, and low-to-intermediate reductions in DTG and BIC susceptibility.

- There is evidence for intermediate **DTG** resistance. If **DTG** is used, it should be administered twice daily.

Drug resistance mutation scores of INSTI:

Download CSV

Rule	BIC ÷	CAB ÷	DTG ÷	EVG ÷	RAL ÷
G140GS	10	10	10	30	30
G140GS + Q148QH	10	20	10	0	0
Q148QH	25	30	25	60	60
Total	45	60	45	90	90