

INSTI Major Mutations:

Q148R • N155H

INSTI Accessory Mutations:

None

IN Other Mutations:

E11D • A21T • D25E • S119G • T122I • T125A • V201I • I203M • T218S • L234I • D253E • D278E

Integrase Strand Transfer Inhibitors	
bictegravir (BIC)	Intermediate Resistance
cabotegravir (CAB)	High-Level Resistance
dolutegravir (DTG)	Intermediate Resistance
elvitegravir (EVG)	High-Level Resistance
raltegravir (RAL)	High-Level Resistance

IN comments

Major

- Q148H/K/R are nonpolymorphic mutations reported in persons receiving RAL, EVG, CAB, and DTG. They nearly always occur in combination with G140A/S or E138K. In this setting they are associated with near complete resistance to RAL and EVG, high-levels of reduction in CAB susceptibility, and low-to-intermediate reductions in DTG and BIC susceptibility.
- N155H is a common nonpolymorphic INSTI-resistance mutations. It has been reported in a high proportion of persons developing VF and HIVDR while receiving RAL, EVG, DTG, and CAB. Alone, it reduces RAL and EVG susceptibility about 10 and 30-fold, respectively. It has minimal effect on susceptibility to DTG, BIC, and CAB.
- There is evidence for intermediate DTG resistance. If DTG is used, it should be administered twice daily.

Drug resistance mutation scores of INSTI:

Download CSV

Rule	BIC ⚡	CAB ⚡	DTG ⚡	EVG ⚡	RAL ⚡
Q148R	25	40	25	60	60
Q148R + N155H	20	20	20	0	0
N155H	10	25	10	60	60
Total	55	85	55	120	120