

Drug resistance interpretation: PR		HIVDB 9.5.1 (2023-11-05)
PI Major Mutations:	None	
PI Accessory Mutations:	None	
PR Other Mutations:	V11X • T12V • I13R • L19V • K20R • L33V • M36I • R41K • L63V • I64V • E65D • K70R	
Protease Inhibitors		
atazanavir/r (ATV/r)	Susceptible	
darunavir/r (DRV/r)	Susceptible	
fosamprenavir/r (FPV/r)	Susceptible	
indinavir/r (IDV/r)	Susceptible	
lopinavir/r (LPV/r)	Susceptible	
nelfinavir (NFV)	Susceptible	
saquinavir/r (SQV/r)	Susceptible	
tipranavir/r (TPV/r)	Susceptible	
PR comments		
Other		
<ul style="list-style-type: none">• K20R is a highly polymorphic PI-selected accessory mutation that increases replication fitness in viruses with PI-resistance mutations.• L33I/V are minimally polymorphic mutations that do not appear to be selected by PIs or to reduce their susceptibility.		

Mutation scoring: PR	HIVDB 9.5.1 (2023-11-05)
----------------------	--------------------------

No drug resistance mutations were found for PI.

Drug resistance interpretation: RT	HIVDB 9.5.1 (2023-11-05)
------------------------------------	--------------------------

NRTI Mutations:	K65R • S68N • M184V		
NNRTI Mutations:	K101H • Y181C • G190A		
RT Other Mutations:	K20R • V35T • K46Q • S48T • K49R • V60I • K122E • I135V • E138T • S162C • E169A • D177E • I178M • T200A • Q207K • R211Q • T216I • P217L • K219X • P225S • P226S • L228H • E233D • P243L • I244* • V245S • L246A • P247E • E248K • D250L • S251D • W252C • T253M • V254I • N255Y • D256R • I257V • Q258V		
Nucleoside Reverse Transcriptase Inhibitors		Non-nucleoside Reverse Transcriptase Inhibitors	
abacavir (ABC)	High-Level Resistance	doravirine (DOR)	Low-Level Resistance
zidovudine (AZT)	Susceptible	efavirenz (EFV)	High-Level Resistance
stavudine (D4T)	Intermediate Resistance	etravirine (ETR)	High-Level Resistance
didanosine (DDI)	High-Level Resistance	nevirapine (NVP)	High-Level Resistance
emtricitabine (FTC)	High-Level Resistance	rilpivirine (RPV)	High-Level Resistance
lamivudine (3TC)	High-Level Resistance		
tenofovir (TDF)	Intermediate Resistance		

RT comments

NRTI

- **K65R** confers intermediate reductions in susceptibility to TDF, ABC, and 3TC/FTC. It increases AZT susceptibility. In NRTI-experienced, INSTI-naïve patients with **K65R**, TDF+3TC+DTG is usually highly effective and more effective than AZT/3TC/DTG. However, in patients receiving TDF+3TC+DTG, there is a risk of emergent DTG resistance that does not arise in NRTI-naïve patients receiving TDF+3TC+DTG.
- **M184V/I** cause high-level in vitro resistance to 3TC and FTC and low/intermediate resistance to ABC (3-fold reduced susceptibility). **M184V/I** are not contraindications to continued treatment with 3TC or FTC because they increase susceptibility to AZT and TDF and are associated with clinically significant reductions in HIV-1 replication.

NNRTI

- **K101H** is a non-polymorphic accessory mutation selected by NVP, EFV and ETR. When present with other NNRTI-resistance mutations, it contributes reduces susceptibility to these NNRTIs.
- **Y181C** is a non-polymorphic mutation selected in persons receiving NVP, ETR and RPV. It confers high-level resistance to NVP, intermediate resistance to ETR and RPV, and low-level resistance to EFV. It does not significantly reduce DOR susceptibility.
- **G190A** is a non-polymorphic mutation that causes high-level resistance to NVP and intermediate resistance to EFV. It does not significantly reduce susceptibility to RPV, ETR, or DOR.

Other

- E138K is a non-polymorphic mutation selected in a high proportion of persons receiving RPV. It reduces RPV susceptibility 2 to 3-fold. In combination with K101E or the NRTI-resistance mutation M184I, it is sufficient to cause VF on a first-line RPV-containing regimen. E138K causes low-level cross-resistance to ETR. E138A is a common polymorphic accessory mutation weakly selected in persons receiving ETR and RPV. It reduces ETR and RPV susceptibility ~2-fold. Its effect on ETR- and RPV-containing regimens is likely to be minimal. E138Q/G are non-polymorphic accessory mutations selected by ETR occasionally NVP and EFV. They cause low-level reductions in susceptibility to NVP, RPV, and ETR. E138R is an extremely rare non-polymorphic accessory mutation selected in vitro by RPV. Its effect on NNRTI susceptibility has not been well studied. **E138T** is an unusual mutation at this position.
- P225H is a non-polymorphic EFV-selected mutation that usually occurs in combination with K103N. The combination of P225H and K103N synergistically reduces NVP, EFV and DOR susceptibility. **P225S** is a highly unusual mutation at this position.

Mutation scoring: RT

HIVDB 9.5.1 (2023-11-05)

Drug resistance mutation scores of NRTI:

Download CSV



Rule	ABC ↕	AZT ↕	D4T ↕	DDI ↕	FTC ↕	3TC ↕	TDF ↕
K65R	45	-10	60	60	30	30	50
M184V	15	-10	-10	10	60	60	-10
K65R + S68N	0	0	0	0	0	0	5
Total	60	-20	50	70	90	90	45

Drug resistance mutation scores of NNRTI:

Download CSV



Rule	DOR ↕	EFV ↕	ETR ↕	NVP ↕	RPV ↕
Y181C	10	30	30	60	45
Y181C + G190A	10	0	10	0	10
K101H	0	10	10	15	10
G190A	0	45	10	60	15
Total	20	85	60	135	80