

PatientID: GU121687

Okitobba 06, 2023

Color Code

■ HR: High-Level Resistance ■ PLR: Potential Low-Level Resistance
■ LR: Low-Level Resistance ■ IR: Intermediate Resistance
■ S: Susceptible

DRUG.CLASS	DRUG	RESISTANCE.PROFILE	DRMS.above.20.percent.prevalence
INSTI	BIC	PLR	N155H;D232N
	CAB	LR	
	DTG	PLR	
	EVG	HR	
	RAL	HR	

Appendix

Drug abbreviations in full

DRUG.CLASS	ABBREVIATION	DRUG.NAME
PI	ATV	Atazanavir
	DRV	Darunavir
	FPV	Fosamprenavir
	IDV	Indinavir
	LPV	Lopinavir
	NFV	Nelfinavir
	SQV	Saquinavir
	TPV	Tipranavir
NRTI	ABC	Abacavir
	AZT	Azidothymidine
	DFT	Stavudine
	DDI	Didanosine
	FTC	Emtricitabine
	LMV	Lamivudine
	TDF	Tenofovir
NNRTI	DOR	Doravirine
	EFV	Efavirenz
	ETR	Etravirine
	NVP	Nevirapine
	RPV	Rilpivirine
INSTI	BIC	Bictegravir
	CAB	Cabotegravir
	DTG	Dolutegravir
	EVG	Elvitegravir
	RAL	Raltegravir

Comments

DRUG.CLASS	COMMENTS
PI	
NRTI	
NNRTI	
INSTI	D232N is a common nonpolymorphic accessory mutation selected in persons receiving RAL and EVG. Alone, it has little effect on INSTI susceptibility.
	N155H is a common nonpolymorphic INSTI-resistance mutations. It has been reported in a high proportion of persons developing VF and HIVDR while receiving RAL, EVG, DTG, and CAB. Alone, it reduces RAL and EVG susceptibility about 10 and 30-fold, respectively. It has minimal effect on susceptibility to DTG, BIC, and CAB.