

Drug resistance interpretation: PR		HIVDB 9.5.1 (2023-11-05)
PI Major Mutations:	None	
PI Accessory Mutations:	None	
PR Other Mutations:	L10Q • V11S • I13V • K14R • I15S • G16E • K20R • E35D • M36I • N37D • R41K • K45R • R57K • L63P • H69K • K70R • L89M	
Protease Inhibitors		
atazanavir/r (ATV/r)	Susceptible	
darunavir/r (DRV/r)	Susceptible	
fosamprenavir/r (FPV/r)	Susceptible	
indinavir/r (IDV/r)	Susceptible	
lopinavir/r (LPV/r)	Susceptible	
nelfinavir (NFV)	Susceptible	
saquinavir/r (SQV/r)	Susceptible	
tipranavir/r (TPV/r)	Susceptible	
PR comments		
Other		
<ul style="list-style-type: none">L10F is a common non-polymorphic, PI-selected accessory mutation associated with reduced in vitro susceptibility to LPV and DRV. L10I/V are polymorphic, PI-selected accessory mutations that increase the replication of viruses with other PI-resistance mutations. L10R/Y are rare, non-polymorphic PI-selected mutations. Their effects on PI susceptibility have not been well studied. L10Q is a highly unusual mutation at this position.K20R is a highly polymorphic PI-selected accessory mutation that increases replication fitness in viruses with PI-resistance mutations.		

Mutation scoring: PR	HIVDB 9.5.1 (2023-11-05)
----------------------	--------------------------

No drug resistance mutations were found for PI.

Drug resistance interpretation: RT	HIVDB 9.5.1 (2023-11-05)
------------------------------------	--------------------------

NRTI Mutations:	K70E • M184V • K219Q		
NNRTI Mutations:	K103N • K238T		
RT Other Mutations:	K20R • V35T • K43R • K49R • P55L • V60I • V118A • K122E • D123G • I135T • E169D • K173A • Q174K • D177E • T200A • Q207A • R211K • P217S • P225I • P226H • F227S • L228C • Y232D • V245E • Δ250 • S251K • W252V • T253D • V254C • N255H • D256E • L264S • N265E • S268X • V276W		
Nucleoside Reverse Transcriptase Inhibitors		Non-nucleoside Reverse Transcriptase Inhibitors	
abacavir (ABC)	Intermediate Resistance	doravirine (DOR)	Susceptible
zidovudine (AZT)	Susceptible	efavirenz (EFV)	High-Level Resistance
stavudine (D4T)	Low-Level Resistance	etravirine (ETR)	Susceptible
didanosine (DDI)	Intermediate Resistance	nevirapine (NVP)	High-Level Resistance
emtricitabine (FTC)	High-Level Resistance	rilpivirine (RPV)	Susceptible
lamivudine (3TC)	High-Level Resistance		
tenofovir (TDF)	Low-Level Resistance		

RT comments

NRTI

- **K70/E/Q/N/T/S/G** cause low-level resistance to ABC and TDF.
- **M184V/I** cause high-level in vitro resistance to 3TC and FTC and low/intermediate resistance to ABC (3-fold reduced susceptibility). **M184V/I** are not contraindications to continued treatment with 3TC or FTC because they increase susceptibility to AZT and TDF and are associated with clinically significant reductions in HIV-1 replication.
- **K219E/Q/N/R** are accessory TAMS that usually occur in combination with multiple other TAMS.

NNRTI

- **K103N** is a non-polymorphic mutation that confers high-level reductions in NVP and EFV susceptibility. It is the most commonly transmitted DRM.
- **K238T/N** are uncommon non-polymorphic mutations selected in persons receiving NVP and EFV usually in combination with K103N. Alone, **K238T/N** appear to have minimal effects on NNRTI susceptibility.

Other

- P225H is a non-polymorphic EFV-selected mutation that usually occurs in combination with K103N. The combination of P225H and K103N synergistically reduces NVP, EFV and DOR susceptibility. **P225I** is a highly unusual mutation at this position.
- F227L is a non-polymorphic mutation that usually occurs in combination with V106A. It is selected in vivo and in vitro with both NVP and DOR. In this context it is associated with high-level reductions in NVP and DOR susceptibility and intermediate reductions in EFV susceptibility. F227I/V are extremely rare mutations that have been selected in vitro by DOR. F227C is a nonpolymorphic mutation selected in persons receiving DOR and rarely in persons receiving ETR and RPV. It usually occurs in combination with other DRMs and in this setting has consistently been associated with the highest possible levels of DOR resistance. It is also usually associated with intermediate or high-level reductions in susceptibility to NVP, EFV, ETR, and RPV. **F227S** is a highly unusual mutation at this position.

Mutation scoring: RT

HIVDB 9.5.1 (2023-11-05)

Drug resistance mutation scores of NRTI:

Download CSV



Rule	ABC ⚡	AZT ⚡	D4T ⚡	DDI ⚡	FTC ⚡	3TC ⚡	TDF ⚡
K70E	15	0	15	15	10	10	15
M184V	15	-10	-10	10	60	60	-10
K219Q	5	10	10	5	0	0	5
K70E + M184V	0	0	10	0	0	0	10
Total	35	0	25	30	70	70	20

Drug resistance mutation scores of NNRTI:

Download CSV



Rule	DOR ⚡	EFV ⚡	ETR ⚡	NVP ⚡	RPV ⚡
K103N	0	60	0	60	0
K238T	0	30	0	30	0
Total	0	90	0	90	0