PatientID: GU121691

Okitobba 06, 2023

## Color Code

HR: High-Level Resistance

LR: Low-Level Resistance

IR: Intermediate Resistance

S: Susceptible

DRUG.CLASS	DRUG	RESISTANCE.PROFILE	DRMS.above.20.percent.prevalence	
	BIC	$\mathbf{S}$		
	CAB	$_{ m LR}$		
INSTI	DTG	$\mathbf{S}$	Y143R;T97A	
	EVG	$_{ m LR}$		
	RAL	HR		

## Appendix

## Drug abbreviations in full

DRUG.CLASS	ABBREVIATION	DRUG.NAME
	ATV	Atazanavir
	DRV	Darunavir
	FPV	Fosamprenavir
PI	IDV	Indinavir
11	LPV	Lopinavir
	NFV	Nelfinavir
	SQV	Saquinavir
	TPV	Tipranavir
	ABC	Abacavir
	AZT	Azidothymidine
	DFT	Stavudine
NRTI	DDI	Didanosine
	FTC	Emtricitabine
	LMV	Lamivudine
	TDF	Tenofovir
	DOR	Doravirine
	EFV	Efavirenz
NNRTI	ETR	Etravirine
	NVP	Nevirapine
	RPV	Rilpivirine
	BIC	Bictegravir
	CAB	Cabotegravir
INSTI	DTG	Dolutegravir
	EVG	Elvitegravir
	RAL	Raltegravir

## Comments

DRUG.CLASS	COMMENTS
PI	
NRTI	
NNRTI	
INSTI	T97A is a polymorphic INSTI-selected mutation that, depending on subtype, occurs in 1% to 5% of viruses from untreated persons. Alone, it has minimal effects on INSTI susceptibility but in combination with other major resistance mutations, it synergistically reduces susceptibility to each of the INSTIs.  Y143C/R/H are non-polymorphic mutations associated with high-level RAL resistance. Alone, they have minimal effects on EVG susceptibility. However, they are associated with intermediate reductions in EVG susceptibility when they occur in combination with one or more accessory INSTI-resistance mutations. Y143 mutations do not reduce susceptibility to DTG, BIC, or CAB.