

PatientID: NC598-1997

Okitobba 06, 2023

### Color Code

■ HR: High-Level Resistance    ■ PLR: Potential Low-Level Resistance  
■ LR: Low-Level Resistance    ■ IR: Intermediate Resistance  
■ S: Susceptible

DRUG.CLASS	DRUG	RESISTANCE.PROFILE	DRMS.above.20.percent.prevalence
PI	ATV	HR	G48V;I54T;V82A;M46MI;K43T
	DRV	S	
	FPV	IR	
	IDV	HR	
	LPV	HR	
	NFV	HR	
	SQV	HR	
	TPV	IR	
NRTI	ABC	HR	K219KN;M41L;D67N;M184V;L210W;T215Y;T69D
	AZT	HR	
	D4T	HR	
	DDI	HR	
	FTC	HR	
	LMV	HR	
	TDF	HR	
NNRTI	DOR	IR	A98G;Y181C;K103N
	EFV	HR	
	ETR	IR	
	NVP	HR	
	RPV	HR	

## Appendix

### Drug abbreviations in full

DRUG.CLASS	ABBREVIATION	DRUG.NAME
<b>PI</b>	ATV	Atazanavir
	DRV	Darunavir
	FPV	Fosamprenavir
	IDV	Indinavir
	LPV	Lopinavir
	NFV	Nelfinavir
	SQV	Saquinavir
	TPV	Tipranavir
<b>NRTI</b>	ABC	Abacavir
	AZT	Azidothymidine
	DFT	Stavudine
	DDI	Didanosine
	FTC	Emtricitabine
	LMV	Lamivudine
	TDF	Tenofovir
<b>NNRTI</b>	DOR	Doravirine
	EFV	Efavirenz
	ETR	Etravirine
	NVP	Nevirapine
	RPV	Rilpivirine
<b>INSTI</b>	BIC	Bictegravir
	CAB	Cabotegravir
	DTG	Dolutegravir
	EVG	Elvitegravir
	RAL	Raltegravir

### Comments

DRUG.CLASS	COMMENTS
<b>PI</b>	G48V is a nonpolymorphic mutation selected by SQV and less often by IDV and LPV. It confers intermediate resistance to ATV but has little if any effect on LPV susceptibility.
	I54A/T/S are non-polymorphic PI-selected mutations that occur almost exclusively in viruses with multiple PI-resistance mutations. I54A/T/S are associated with reduced susceptibility to each of the PIs except DRV.
	K43T is a nonpolymorphic accessory mutation selected by ATV and LPV. Its phenotypic effect on currently used PIs is uncertain.
	M46I/L are relatively non-polymorphic PI-selected mutations. In combination with other PI-resistance mutations, they are associated with reduced susceptibility to each of the PIs except DRV.
	V82A is a non-polymorphic mutation selected primarily by IDV and LPV. It is associated with reduced susceptibility to LPV and to a lesser extent ATV. It increases DRV susceptibility.
	D67N is a non-polymorphic TAM associated with low-level resistance to AZT.
	K219E/Q/N/R are accessory TAMs that usually occur in combination with multiple other TAMs.

<b>NRTI</b>	L210W is a TAM that usually occurs in combination with M41L and T215Y. The combination of M41, L210W and T215Y causes high-level resistance to AZT and intermediate resistance to ABC and TDF.
	M184V/I cause high-level in vitro resistance to 3TC and FTC and low/intermediate resistance to ABC (3-fold reduced susceptibility). M184V/I are not contraindications to continued treatment with 3TC or FTC because they increase susceptibility to AZT and TDF and are associated with clinically significant reductions in HIV-1 replication.
	M41L is a TAM that usually occurs with T215Y. In combination, M41L plus T215Y confer intermediate / high-level resistance to AZT and d4T and contribute to reduced ddI, ABC and TDF susceptibility.
	T215Y/F are TAMs that causes intermediate/high-level resistance to AZT and potentially low-level resistance to ABC and TDF.
	T69D is a nonpolymorphic mutation selected by early NRTIs that does not appear to reduce AZT, ABC, or TDF susceptibility.
<b>NNRTI</b>	A98G is a non-polymorphic accessory mutation associated with low-level reduced susceptibility to each of the NNRTIs.
	K103N is a non-polymorphic mutation that confers high-level reductions in NVP and EFV susceptibility. It is the most commonly transmitted DRM.
	Y181C is a non-polymorphic mutation selected in persons receiving NVP, ETR and RPV. It confers high-level resistance to NVP, intermediate resistance to ETR and RPV, and low-level resistance to EFV. It does not significantly reduce DOR susceptibility.
<b>INSTI</b>	