

PatientID: HDR77

Okitobba 06, 2023

Color Code

■ HR: High-Level Resistance ■ PLR: Potential Low-Level Resistance
■ LR: Low-Level Resistance ■ IR: Intermediate Resistance
■ S: Susceptible

DRUG.CLASS	DRUG	RESISTANCE.PROFILE	DRMS.above.20.percent.prevalence
PI	ATV	S	
	DRV	S	
	FPV	S	
	IDV	S	
	LPV	S	
	NFV	S	
	SQV	S	
	TPV	S	
NRTI	ABC	IR	D67G;K70E;Y115F;K219R
	AZT	LR	
	D4T	IR	
	DDI	LR	
	FTC	PLR	
	LMV	PLR	
	TDF	IR	
NNRTI	DOR	HR	V106M;F227L;Y181S;G190A;K238T
	EFV	HR	
	ETR	LR	
	NVP	HR	
	RPV	IR	
INSTI	BIC	S	
	CAB	S	
	DTG	S	
	EVG	S	
	RAL	S	

Appendix

Drug abbreviations in full

DRUG.CLASS	ABBREVIATION	DRUG.NAME
PI	ATV	Atazanavir
	DRV	Darunavir
	FPV	Fosamprenavir
	IDV	Indinavir
	LPV	Lopinavir
	NFV	Nelfinavir
	SQV	Saquinavir
	TPV	Tipranavir
NRTI	ABC	Abacavir
	AZT	Azidothymidine
	DFT	Stavudine
	DDI	Didanosine
	FTC	Emtricitabine
	LMV	Lamivudine
	TDF	Tenofovir
NNRTI	DOR	Doravirine
	EFV	Efavirenz
	ETR	Etravirine
	NVP	Nevirapine
	RPV	Rilpivirine
INSTI	BIC	Bictegravir
	CAB	Cabotegravir
	DTG	Dolutegravir
	EVG	Elvitegravir
	RAL	Raltegravir

Comments

DRUG.CLASS	COMMENTS
PI	
NRTI	D67N is a non-polymorphic TAM associated with low-level resistance to AZT. D67G/E/S/T/H are non-polymorphic NRTI-selected mutations that generally occur in viruses with multiple TAMs.
	K219E/Q/N/R are accessory TAMs that usually occur in combination with multiple other TAMs.
	K70/E/Q/N/T/S/G cause low-level resistance to ABC and TDF.
	Y115F causes intermediate resistance to ABC and low-level resistance to TDF.
	F227L is a non-polymorphic mutation that usually occurs in combination with V106A. It is selected in vivo and in vitro with both NVP and DOR. In this context it is associated with high-level reductions in NVP and DOR susceptibility and intermediate reductions in EFV susceptibility. F227I/V are extremely rare mutations that have been selected in vitro by DOR.
	G190A is a non-polymorphic mutation that causes high-level resistance to NVP and intermediate resistance to EFV. It does not significantly reduce susceptibility to RPV, ETR, or DOR.

NNRTI	K238T/N are uncommon non-polymorphic mutations selected in persons receiving NVP and EFV usually in combination with K103N. Alone, K238T/N appear to have minimal effects on NNRTI susceptibility.
	V106M is a non-polymorphic mutation that confers high-level resistance to NVP and EFV. It is selected in vitro and in vivo by DOR and preliminary data suggests it reduces DOR susceptibility about 3-fold.
	Y181F/S/G are rare non-polymorphic NNRTI-associated mutations that are usually present as part of an electrophoretic mixture. They are likely to represent transitional mutations between Y and I or V.
INSTI	