PatientID: HIVDR-1676-23

Sebuttemba 27, 2023

Color Code

HR: High-Level Resistance
LR: Low-Level Resistance
IR: Intermediate Resistance

S: Susceptible

DRUG.CLASS	DRUG	RESISTANCE.PROFILE	DRMS.above.20.percent.prevalence
PI	ATV	S	
	DRV	\mathbf{S}	
	FPV	\mathbf{S}	
	IDV	\mathbf{S}	
	LPV	${f S}$	
	NFV	${f S}$	
	SQV	${f S}$	
	TPV	${f S}$	
NRTI	ABC	IR	
	AZT	IR	
	D4T	IR	
	DDI	IR	M41L;K70R;M184V;K219Q
	FTC	$_{ m HR}$	
	LMV	$_{ m HR}$	
	TDF	${f S}$	
NNRTI	DOR	$_{ m HR}$	
	EFV	$_{ m HR}$	
	ETR	\mathbf{S}	V106A;P225H;K103T
	NVP	HR	
	RPV	\mathbf{S}	

Appendix

Drug abbreviations in full

DRUG.CLASS	ABBREVIATION	DRUG.NAME
	ATV	Atazanavir
	DRV	Darunavir
	FPV	Fosamprenavir
PI	IDV	Indinavir
11	LPV	Lopinavir
	NFV	Nelfinavir
	SQV	Saquinavir
	TPV	Tipranavir
	ABC	Abacavir
	AZT	Azidothymidine
	DFT	Stavudine
NRTI	DDI	Didanosine
	FTC	Emtricitabine
	LMV	Lamivudine
	TDF	Tenofovir
	DOR	Doravirine
	EFV	Efavirenz
NNRTI	ETR	Etravirine
	NVP	Nevirapine
	RPV	Rilpivirine
	BIC	Bictegravir
	CAB	Cabotegravir
INSTI	DTG	Dolutegravir
	EVG	Elvitegravir
	RAL	Raltegravir

Comments

DRUG.CLASS	COMMENTS
PI	
NRTI	K219E/Q/N/R are accessory TAMS that usually occur in combination with multiple other TAMs. K70R is a TAM that confers intermediate resistance to AZT and contributes to reduced ABC and TDF susceptibility in combination with other TAMs. M184V/I cause high-level in vitro resistance to 3TC and FTC and low/intermediate resistance to ABC (3-fold reduced susceptibility). M184V/I are not contraindications to continued treatment with 3TC or FTC because they increase susceptibility to AZT and TDF and are associated with clinically significant reductions in HIV-1 replication. M41L is a TAM that usually occurs with T215Y. In combination, M41L plus T215Y confer intermediate / high-level resistance to AZT and d4T and contribute to reduced ddI, ABC and TDF susceptibility. K103T is an extremely rare non-polymorphic mutation that appears to confer intermediate/high-level resistance to NVP but it has little if any effect on EFV susceptibility. P225H is a non-polymorphic EFV-selected mutation that usually occurs in combination with K103N. The combination of P225H and K103N synergistically reduces NVP, EFV and
	DOR susceptibility.

\mathbf{NNRTI}	
	V106A is a non-polymorphic mutation that confers high-level resistance to NVP and DOR,
	and intermediate resistance to EFV. It is commonly selected in vitro and in vivo by DOR.
INSTI	