






PatientID: HIVDR-1769-23

Sebuttemba 27, 2023

### Color Code

 HR: High-Level Resistance     PLR: Potential Low-Level Resistance  
 LR: Low-Level Resistance     IR: Intermediate Resistance  
 S: Susceptible

DRUG.CLASS	DRUG	RESISTANCE.PROFILE	DRMS.above.20.percent.prevalence
NRTI	ABC	S	
	AZT	S	
	D4T	S	
	DDI	S	
	FTC	S	
	LMV	S	
	TDF	S	
NNRTI	DOR	S	
	EFV	S	
	ETR	S	
	NVP	S	
	RPV	S	
INSTI	BIC	HR	E138K;G140A;Q148K
	CAB	HR	
	DTG	HR	
	EVG	HR	
	RAL	HR	

## Appendix

### Drug abbreviations in full

DRUG.CLASS	ABBREVIATION	DRUG.NAME
<b>PI</b>	ATV	Atazanavir
	DRV	Darunavir
	FPV	Fosamprenavir
	IDV	Indinavir
	LPV	Lopinavir
	NFV	Nelfinavir
	SQV	Saquinavir
	TPV	Tipranavir
<b>NRTI</b>	ABC	Abacavir
	AZT	Azidothymidine
	DFT	Stavudine
	DDI	Didanosine
	FTC	Emtricitabine
	LMV	Lamivudine
	TDF	Tenofovir
<b>NNRTI</b>	DOR	Doravirine
	EFV	Efavirenz
	ETR	Etravirine
	NVP	Nevirapine
	RPV	Rilpivirine
<b>INSTI</b>	BIC	Bictegravir
	CAB	Cabotegravir
	DTG	Dolutegravir
	EVG	Elvitegravir
	RAL	Raltegravir

### Comments

DRUG.CLASS	COMMENTS
<b>PI</b>	
<b>NRTI</b>	
<b>NNRTI</b>	
	E138K/A/T are common nonpolymorphic accessory resistance mutations selected in patients receiving RAL, EVG, CAB, and DTG. Alone they do not reduce INSTI susceptibility. However, they contribute to reduced susceptibility in combination with other mutations particularly those at position 148.
	G140S/A/C are non-polymorphic mutations that usually occur with Q148 mutations. Alone, they have minimal effects on INSTI susceptibility. However, in combination with Q148 mutations they are associated with high-level resistance to RAL and EVG and intermediate reductions in DTG and BIC susceptibility.
	Q148H/K/R are nonpolymorphic mutations reported in persons receiving RAL, EVG, CAB, and DTG. They nearly always occur in combination with G140A/S or E138K. In this setting they are associated with near complete resistance to RAL and EVG, high-levels of reduction in CAB susceptibility, and low-to-intermediate reductions in DTG and BIC susceptibility.

## **INSTI**

S230R is a nonpolymorphic INSTI-selected mutation that primarily occurs in combination with other INSTI-resistance mutations. By itself, it appears to have minimal effect on susceptibility to available INSTIs.
---