

PatientID: HDR51

Okitobba 06, 2023

Color Code

■ HR: High-Level Resistance ■ PLR: Potential Low-Level Resistance
■ LR: Low-Level Resistance ■ IR: Intermediate Resistance
■ S: Susceptible

DRUG.CLASS	DRUG	RESISTANCE.PROFILE	DRMS.above.20.percent.prevalence
PI	ATV	S	
	DRV	S	
	FPV	S	
	IDV	S	
	LPV	S	
	NFV	S	
	SQV	S	
	TPV	S	
NRTI	ABC	IR	D67N;K70E;M184V;T215I
	AZT	LR	
	D4T	IR	
	DDI	IR	
	FTC	HR	
	LMV	HR	
	TDF	LR	
NNRTI	DOR	HR	Y188L;V179D
	EFV	HR	
	ETR	LR	
	NVP	HR	
	RPV	HR	

Appendix

Drug abbreviations in full

DRUG.CLASS	ABBREVIATION	DRUG.NAME
PI	ATV	Atazanavir
	DRV	Darunavir
	FPV	Fosamprenavir
	IDV	Indinavir
	LPV	Lopinavir
	NFV	Nelfinavir
	SQV	Saquinavir
	TPV	Tipranavir
NRTI	ABC	Abacavir
	AZT	Azidothymidine
	DFT	Stavudine
	DDI	Didanosine
	FTC	Emtricitabine
	LMV	Lamivudine
	TDF	Tenofovir
NNRTI	DOR	Doravirine
	EFV	Efavirenz
	ETR	Etravirine
	NVP	Nevirapine
	RPV	Rilpivirine
INSTI	BIC	Bictegravir
	CAB	Cabotegravir
	DTG	Dolutegravir
	EVG	Elvitegravir
	RAL	Raltegravir

Comments

DRUG.CLASS	COMMENTS
PI	
NRTI	D67N is a non-polymorphic TAM associated with low-level resistance to AZT.
	K70/E/Q/N/T/S/G cause low-level resistance to ABC and TDF.
	M184V/I cause high-level in vitro resistance to 3TC and FTC and low/intermediate resistance to ABC (3-fold reduced susceptibility). M184V/I are not contraindications to continued treatment with 3TC or FTC because they increase susceptibility to AZT and TDF and are associated with clinically significant reductions in HIV-1 replication.
	T215Y/F are TAMs that causes intermediate/high-level resistance to AZT and potentially low-level resistance to ABC and TDF. T215S/C/D/E/I/V/N/A/L do not reduce NRTI susceptibility but arise from viruses that once contained T215Y/F. The presence of one of these revertant mutations suggests that the patient may have once been infected with a virus containing T215Y/F.
NNRTI	V179D/E are somewhat polymorphic accessory NNRTI-selected mutation. In combination with other NNRTI DRMs, they appear to contribute low-levels of reduced susceptibility to each of the NNRTIs. In particular, the combinations of K103R/V179D and V106I/V179D act synergistically to reduce NVP and EFV susceptibility.
	Y188L is a non-polymorphic mutation that confers high-level resistance to NVP, EFV, RPV, and DOR, and potentially low-level resistance to ETR.

INSTI	
--------------	--