

PI Major Mutations:None

PI Accessory Mutations:None

PR Other Mutations:

L10X • V111L • T12Q • I13R • K14* • I15D • G16R • Q18K • E35D • M36I • R41K • R57K • I62V • L63P • I64L • H69K • L89M

Protease Inhibitors	
atazanavir/r (ATV/r)	Susceptible
darunavir/r (DRV/r)	Susceptible
fosamprenavir/r (FPV/r)	Susceptible
indinavir/r (IDV/r)	Susceptible
lopinavir/r (LPV/r)	Susceptible
nelfinavir (NFV)	Susceptible
saquinavir/r (SQV/r)	Susceptible
tipranavir/r (TPV/r)	Susceptible

PR comments

Other

- V111I/L are relatively non-polymorphic accessory mutation selected in persons receiving DRV. V111L is a nonpolymorphic PI-selected mutation associated with reduced in vitro DRV susceptibility when it occurs in combination with other PI-resistance mutations.

No drug resistance mutations were found for PI.

NRTI Mutations:

L74V • M184V

NNRTI Mutations:

K103N • P225H

RT Other Mutations:V35I • V60I • D121Y • K122E • I135T • I142V • K173L • Q174K • V179I • Q207A • R211S • V245E • E248D • S268G • A272S

Nucleoside Reverse Transcriptase Inhibitors		Non-nucleoside Reverse Transcriptase Inhibitors	
abacavir (ABC)	High-Level Resistance	doravirine (DOR)	Intermediate Resistance
zidovudine (AZT)	Susceptible	efavirenz (EFV)	High-Level Resistance
stavudine (D4T)	Susceptible	etravirine (ETR)	Susceptible
didanosine (DDI)	High-Level Resistance	nevirapine (NVP)	High-Level Resistance
emtricitabine (FTC)	High-Level Resistance	rilpivirine (RPV)	Susceptible
lamivudine (3TC)	High-Level Resistance		
tenofovir (TDF)	Susceptible		

RT comments

NRTI

- L74V causes intermediate ABC resistance. L74I causes low-level ABC resistance.
- M184V/I cause high-level in vitro resistance to 3TC and FTC and low/intermediate resistance to ABC (3-fold reduced susceptibility). M184V/I are not contraindications to continued treatment with 3TC or FTC because they increase susceptibility to AZT and TDF and are associated with clinically significant reductions in HIV-1 replication.

NNRTI

- K103N is a non-polymorphic mutation that confers high-level reductions in NVP and EFV susceptibility. It is the most commonly transmitted DRM.
- P225H is a non-polymorphic EFV-selected mutation that usually occurs in combination with K103N. The combination of P225H and K103N synergistically reduces NVP, EFV and DOR susceptibility.

Other

- V179I is a polymorphic mutation that is frequently selected in persons receiving ETR and RPV. However, it has little, if any, direct effect on NNRTI susceptibility.

Drug resistance mutation scores of NRTI:

Download CSV



Rule	ABC ⚡	AZT ⚡	D4T ⚡	DDI ⚡	FTC ⚡	3TC ⚡	TDF ⚡
<u>L74V</u>	30	0	0	60	0	0	0
<u>L74V + M184V</u>	15	0	0	0	0	0	0
<u>M184V</u>	15	-10	-10	10	60	60	-10
Total	60	-10	-10	70	60	60	-10

Drug resistance mutation scores of NNRTI:

Download CSV



Rule	DOR ⚡	EFV ⚡	ETR ⚡	NVP ⚡	RPV ⚡
<u>K103N + P225H</u>	10	0	0	0	0
<u>P225H</u>	20	45	0	45	0
<u>K103N</u>	0	60	0	60	0
Total	30	105	0	105	0