

PatientID: HDR120

Okitobba 06, 2023

Color Code

■ HR: High-Level Resistance ■ PLR: Potential Low-Level Resistance
■ LR: Low-Level Resistance ■ IR: Intermediate Resistance
■ S: Susceptible

DRUG.CLASS	DRUG	RESISTANCE.PROFILE	DRMS.above.20.percent.prevalence
PI	ATV	IR	I54V;V82A;L24I;L10F;L89V
	DRV	PLR	
	FPV	HR	
	IDV	HR	
	LPV	HR	
	NFV	HR	
	SQV	IR	
	TPV	LR	
NRTI	ABC	HR	M41L;L210W;T215Y;V75M
	AZT	HR	
	D4T	HR	
	DDI	HR	
	FTC	LR	
	LMV	LR	
	TDF	HR	
NNRTI	DOR	PLR	Y181C
	EFV	IR	
	ETR	IR	
	NVP	HR	
	RPV	IR	

Appendix

Drug abbreviations in full

DRUG.CLASS	ABBREVIATION	DRUG.NAME
PI	ATV	Atazanavir
	DRV	Darunavir
	FPV	Fosamprenavir
	IDV	Indinavir
	LPV	Lopinavir
	NFV	Nelfinavir
	SQV	Saquinavir
	TPV	Tipranavir
NRTI	ABC	Abacavir
	AZT	Azidothymidine
	DFT	Stavudine
	DDI	Didanosine
	FTC	Emtricitabine
	LMV	Lamivudine
	TDF	Tenofovir
NNRTI	DOR	Doravirine
	EFV	Efavirenz
	ETR	Etravirine
	NVP	Nevirapine
	RPV	Rilpivirine
INSTI	BIC	Bictegravir
	CAB	Cabotegravir
	DTG	Dolutegravir
	EVG	Elvitegravir
	RAL	Raltegravir

Comments

DRUG.CLASS	COMMENTS
PI	I54V is a non-polymorphic PI-selected mutation that contributes reduced susceptibility to each of the PIs except DRV.
	L10F is a common non-polymorphic, PI-selected accessory mutation associated with reduced in vitro susceptibility to LPV and DRV.
	L24I is a non-polymorphic mutation selected by IDV and LPV. It contributes reduced susceptibility to ATV and LPV.
	L89V is a nonpolymorphic accessory mutation weakly selected by each of the PIs. It appears to be minimally associated with reduced PI susceptibility. L89T is an uncommon non-polymorphic PI-selected mutation selected primarily by ATV.
	V82A is a non-polymorphic mutation selected primarily by IDV and LPV. It is associated with reduced susceptibility to LPV and to a lesser extent ATV. It increases DRV susceptibility.
	L210W is a TAM that usually occurs in combination with M41L and T215Y. The combination of M41, L210W and T215Y causes high-level resistance to AZT and intermediate resistance to ABC and TDF.

NRTI	M41L is a TAM that usually occurs with T215Y. In combination, M41L plus T215Y confer intermediate / high-level resistance to AZT and d4T and contribute to reduced ddI, ABC and TDF susceptibility.
	T215Y/F are TAMs that causes intermediate/high-level resistance to AZT and potentially low-level resistance to ABC and TDF.
	V75T/M/A/S are nonpolymorphic accessory NRTI-selected mutations. They appear to have minimal phenotypic effects on AZT, ABC, and TDF.
NNRTI	Y181C is a non-polymorphic mutation selected in persons receiving NVP, ETR and RPV. It confers high-level resistance to NVP, intermediate resistance to ETR and RPV, and low-level resistance to EFV. It does not significantly reduce DOR susceptibility.
INSTI	