

# Supplementary Information

## **Dynamics of Epiphytic Microbial Communities on Marine Macrophyte Surfaces**


1†

† To whom correspondence should be addressed: [marino.korlevic@irb.hr](mailto:marino.korlevic@irb.hr)


1. Ruđer Bošković Institute, Center for Marine Research, G. Paliaga 5, Rovinj, Croatia

2. University of Vienna, Department of Limnology and Bio-Oceanography, Althanstraße 14,  
Vienna, Austria


# 1 Supplementary Figures



**Figure S1.** Rarefaction curves of bacterial and archaeal communities from the surfaces of macrophytes (*Cymodocea nodosa* [Invaded] and *Caulerpa cylindracea* [Invaded and Noninvaded]) and in the surrounding seawater.



**Figure S2.** Seasonal dynamics of observed number of OTUs, Chao1, ACE, exponential of the Shannon Diversity Index and Inverse Simpson Index of bacterial and archaeal communities from the surfaces of macrophytes (*Cymodocea nodosa* [Invaded] and *Caulerpa cylindracea* [Invaded and Noninvaded]) and in the surrounding seawater.



**Figure S3.** Relative contribution of chloroplast sequences on the surfaces of macrophytes (*Cymodocea nodosa* [Invaded] and *Caulerpa cylindracea* [Invaded and Noninvaded]) and in the surrounding seawater.