

Selective DNA and Protein Isolation from Marine Macrophyte Surfaces

1†

† To whom correspondence should be addressed: marino.korlevic@irb.hr

1. Ruđer Bošković Institute, Center for Marine Research, G. Paliaga 5, Rovinj, Croatia

2. University of Vienna, Department of Limnology and Bio-Oceanography, Althanstraße 14,
Vienna, Austria

1 Abstract

2 Introduction

3 Surfaces of marine macrophytes are inhabited by a diverse microbial community whose
4 structure and function are poorly understood (Egan *et al.*, 2013). As less than 1 % of all
5 prokaryotic species are culturable, to study these organisms, molecular methods such as 16S
6 rRNA sequencing, metagenomics and metaproteomics are indispensable (Amann *et al.*, 1995;
7 Su *et al.*, 2012). Applying these techniques requires an initial isolation step, with the purpose of
8 obtaining high quality DNA and proteins.

9 Biological material (i.e. proteins and DNA) from pelagic microbial communities is usually
10 isolated by collecting cells onto filters and subsequently isolating the aimed compound (Gilbert
11 *et al.*, 2009). If a microbial size fraction is aimed sequential filtration is applied (Massana *et al.*,
12 1997; Andersson *et al.*, 2010). In contrast, obtaining biological materials from microorganism
13 inhabiting surfaces requires either a cell detachment procedure prior to isolation or the host material
14 is coextracted together with the targeted material. Methods for separating microbial cells from the
15 host include shaking of host tissue (Gross *et al.*, 2003; Nöges *et al.*, 2010), scraping of macrophyte
16 surfaces (Uku *et al.*, 2007) or the application of ultrasonication (Weidner *et al.*, 1996; Cai *et al.*,
17 2014). It was shown that shaking alone is not sufficient to remove microbial cells, at least from
18 plant root surfaces (Richter-Heitmann *et al.*, 2016). Manual separation methods, such as scraping
19 and brushing, are time consuming and subjective, as the detachment efficiency depends on host
20 tissue and the person performing the procedure (Cai *et al.*, 2014). Ultrasonication was proposed as
21 an alternative method as it is providing better results in terms of detachment efficiency (Cai *et al.*,
22 2014; Richter-Heitmann *et al.*, 2016). The downside of this procedure is that complete cell removal
23 was still not obtained and tissue disruption was observed especially after the application of probe
24 ultrasonication (Richter-Heitmann *et al.*, 2016). An alternative to these cell detachment procedures
25 is the isolation of targeted epiphytic compounds together with host materials (Staufenberger *et al.*,
26 2008; Jiang *et al.*, 2015). This procedure can lead to problems in the following processing
27 steps such as mitochondrial and chloroplast 16S rRNA sequence contaminations from the host

(Longford *et al.*, 2007; Staufenberg *et al.*, 2008). In addition, when performing metagenomics and metaproteomics host material can cause biased results towards more abundant host DNA and proteins.

An alternative to these procedures is a direct isolation of the targeted material by incubating macrophyte tissues in an extraction buffer. After the incubation is done the undisturbed host tissue is removed and the isolation procedure continues omitting host material contaminations. To our knowledge the only procedure describing a direct and selective epiphytic DNA isolation from the surfaces of marine macrophytes was provided by Burke *et al.* (2009). In contrast to previously described methods this protocol enables an almost complete removal of the surface community and was used for 16S rRNA gene clone libraries construction (Burke *et al.*, 2011b) and metagenomes sequencing (Burke *et al.*, 2011a). This method, though providing a selective isolation procedure, is using in the extraction buffer a rapid multienzyme cleaner (3M) that is not worldwide available and whose composition is not known (Burke *et al.*, 2009). Also to our knowledge, no selective isolation protocol for proteins from epiphytic communities inhabiting marine macrophytes was established.

In the present study, we adapted a protocol widely used for DNA isolation from filters (Massana *et al.*, 1997) and a protocol used for protein isolation from sediments (Chourey *et al.*, 2010) for selective extractions of DNA and proteins from epiphytic communities inhabiting the surfaces of two marine macrophytes: the seagrass *Cymodocea nodosa* and the alga *Caulerpa cylindracea*. In addition, we tested the removal efficiency of the protocol and the suitability of obtained DNA and proteins for 16S rRNA sequencing and metaproteomics.

Materials and Methods

Sampling

Leaves of *Cymodocea nodosa* were sampled in a *Cymodocea nodosa* meadow in the Bay of Saline (45°7'5"N; 13°37'20"E) and in a *Cymodocea nodosa* meadow invaded by *Caulerpa cylindracea* in the proximity of the village of Funtana (45°10'39"N 13°35'42"E). Thalli of *Caulerpa cylindracea* were sampled in the same *Cymodocea nodosa* invaded meadow in Funtana and on a locality of only *Caulerpa cylindracea* located close to the invaded meadow. Seagrasses and algae were collected the same day in two contrasting seasons, on 4 December 2017 and 18 June 2018. During spring 2018 the *Cymodocea nodosa* meadow in the Bay of Saline decayed to the extent that no leaves could be retrieved (Najdek *et al.*, unpublished data). Leaves and thalli were collected by diving and transported to the laboratory in containers placed on ice and filled with site seawater. Upon arrival to the laboratory, *Cymodocea nodosa* leaves were cut into sections of 1 – 2 cm, while *Caulerpa cylindracea* thalli were cut into 5 – 8 cm long sections. Leaves and thalli were washed three times with sterile artificial seawater (ASW) to remove loosely attached microbial cells.

DNA isolation

The DNA was isolated according to the protocol for isolation from filters described in Massana *et al.* (1997). This protocol was modified and adapted for DNA isolation from microbial communities from macrophytes surfaces as described below. 1 g wet weight of leaves and 2 g wet-weight of thalli were placed into 5 ml of lysis buffer (40 mM EDTA, 50 mM Tris-HCl, 0.75 M sucrose; pH 8.3). Lysozyme was added (final concentration 1 mg ml⁻¹) and the mixture was incubated at 37 °C for 30 minutes. Subsequently, proteinase K (final concentration 0.5 mg ml⁻¹) and SDS (final concentration 1 %) were added and the samples were incubated at 55 °C for 2

71 hours. Following the incubation, tubes were vortexed for 10 minutes and the mixture containign
72 lyzed cells was separated from host leaves or thalli by transferrring the solution into a clean tube.
73 The lysate was extracted twice with a mixture of phenol:chloroform:isoamyl alcohol (25:24:1;
74 pH 8) and once with chloroform-isoamyl alcohol (24:1). After each organic solvent mixture
75 addition tubes were slightly vortexed and centrifugated at 4,500 x g for 10 minutes. Following
76 each centrifugation aqueous phases were retrieved. After the final extraction 1/10 of 3 M sodium
77 acetate (ph 5.2) was added. DNA was precipitated by adding 1 volumne of isopropanol and
78 incubating the mixtures at -20 °C overnight.

79 **Protein isolation**

80 **Results and Discussion**

81 **Conclusions**

82 **Materials and Methods**

References

- Amann, R.I., Ludwig, W., and Schleifer, K.H. (1995) Phylogenetic identification and in situ detection of individual microbial cells without cultivation. *Microbiological reviews* **59**: 143–169.
- Andersson, A.F., Riemann, L., and Bertilsson, S. (2010) Pyrosequencing reveals contrasting seasonal dynamics of taxa within Baltic Sea bacterioplankton communities. *The ISME journal* **4**: 171–181.
- Burke, C., Kjelleberg, S., and Thomas, T. (2009) Selective extraction of bacterial DNA from the surfaces of macroalgae. *Applied and environmental microbiology* **75**: 252–256.
- Burke, C., Steinberg, P., Rusch, D., Kjelleberg, S., and Thomas, T. (2011a) Bacterial community assembly based on functional genes rather than species. *Proceedings of the National Academy of Sciences of the United States of America* **108**: 14288–14293.
- Burke, C., Thomas, T., Lewis, M., Steinberg, P., and Kjelleberg, S. (2011b) Composition, uniqueness and variability of the epiphytic bacterial community of the green alga *Ulva australis*. *The ISME journal* **5**: 590–600.
- Cai, X., Gao, G., Yang, J., Tang, X., Dai, J., Chen, D., and Song, Y. (2014) An ultrasonic method for separation of epiphytic microbes from freshwater submerged macrophytes. *Journal of basic microbiology* **54**: 758–761.
- Chourey, K., Jansson, J., VerBerkmoes, N., Shah, M., Chavarria, K.L., Tom, L.M., et al. (2010) Direct Cellular Lysis/Protein Extraction Protocol for Soil Metaproteomics. *Journal of Proteome Research* **9**: 6615–6622.
- Egan, S., Harder, T., Burke, C., Steinberg, P., Kjelleberg, S., and Thomas, T. (2013) The seaweed holobiont: understanding seaweed-bacteria interactions. *FEMS microbiology reviews* **37**:

462–476.

Gilbert, J.A., Field, D., Swift, P., Newbold, L., Oliver, A., Smyth, T., et al. (2009) The seasonal structure of microbial communities in the Western English Channel. *Environmental Microbiology* **11**: 3132–3139.

Gross, E.M., Feldbaum, C., and Graf, A. (2003) Epiphyte biomass and elemental composition on submersed macrophytes in shallow eutrophic lakes. *Hydrobiologia* **506-509**: 559–565.

Jiang, Y.-F., Ling, J., Dong, J.-D., Chen, B., Zhang, Y.-Y., Zhang, Y.-Z., and Wang, Y.-S. (2015) Illumina-based analysis the microbial diversity associated with *Thalassia hemprichii* in Xincun Bay, South China Sea. *Ecotoxicology (London, England)* **24**: 1548–56.

Longford, S., Tujula, N., Crocetti, G., Holmes, A., Holmström, C., Kjelleberg, S., et al. (2007) Comparisons of diversity of bacterial communities associated with three sessile marine eukaryotes. *Aquatic Microbial Ecology* **48**: 217–229.

Massana, R., Murray, A.E., Preston, C.M., and DeLong, E.F. (1997) Vertical distribution and phylogenetic characterization of marine planktonic Archaea in the Santa Barbara Channel. *Applied and environmental microbiology* **63**: 50–56.

Nõges, T., Luup, H., and Feldmann, T. (2010) Primary production of aquatic macrophytes and their epiphytes in two shallow lakes (Peipsi and Võrtsjärv) in Estonia. *Aquatic Ecology* **44**: 83–92.

Richter-Heitmann, T., Eickhorst, T., Knauth, S., Friedrich, M.W., and Schmidt, H. (2016) Evaluation of Strategies to Separate Root-Associated Microbial Communities: A Crucial Choice in Rhizobiome Research. *Frontiers in Microbiology* **7**: 773.

Staufenberger, T., Thiel, V., Wiese, J., and Imhoff, J.F. (2008) Phylogenetic analysis of

127 bacteria associated with *Laminaria saccharina*. *FEMS Microbiology Ecology* **64**: 65–77.

128 Su, C., Lei, L., Duan, Y., Zhang, K.-Q., and Yang, J. (2012) Culture-independent methods
129 for studying environmental microorganisms: methods, application, and perspective. *Applied*
130 *microbiology and biotechnology* **93**: 993–1003.

131 Uku, J., Björk, M., Bergman, B., and Díez, B. (2007) Characterization and comparison of
132 prokaryotic epiphytes associated with three East African seagrasses. *Journal of Phycology* **43**:
133 768–779.

134 Weidner, S., Arnold, W., and Puhler, A. (1996) Diversity of uncultured microorganisms
135 associated with the seagrass *Halophila stipulacea* estimated by restriction fragment length
136 polymorphism analysis of PCR-amplified 16S rRNA genes. *Applied and environmental*
137 *microbiology* **62**: 766–771.