

Supplementary Information

Selective DNA and Protein Isolation from Marine Macrophyte Surfaces

Marino Korlević^{1*}, Marsej Markovski¹, Zihao Zhao², Gerhard J. Herndl^{2,3}, Mirjana Najdek¹

*To whom correspondence should be addressed: marino.korlevic@irb.hr

1. Center for Marine Research, Ruđer Bošković Institute, Croatia
2. Department of Functional and Evolutionary Ecology, University of Vienna, Austria
3. Department of Marine Microbiology and Biogeochemistry, Royal Netherlands Institute for Sea Research, Utrecht University, The Netherlands

Supplementary Figures




Fig. S1. Rarefaction curves of bacterial and archaeal communities from the surfaces of macrophytes sampled in the Bay of Saline (*C. nodosa* [Monospecific Settlement]) and the Bay of Funatana (*C. nodosa* [Mixed Settlement] and *C. cylindracea* [Mixed and Monospecific Settlement]) in two contrasting seasons.

Supplementary Table

Table S1. Sample ID, Station, Community Type, Sampling Date, No. of Sequences and No. of OTUs of each sample. No. of Sequences and OTUs was calculated after the exclusion of eukaryotic, chloroplast, mitochondrial and no relative sequences.

Sample ID	Station	Community Type	Date	No. of Sequences	No. of OTUs
40	Saline	<i>Cymodocea nodosa</i> (Monospecific)	4 December 2017	19,176	942
41	Funtana	<i>Cymodocea nodosa</i> (Mixed)	4 December 2017	24,250	1,123
42	Funtana	<i>Caulerpa cylindracea</i> (Mixed)	4 December 2017	28,387	3,131
43	Funtana	<i>Caulerpa cylindracea</i> (Monospecific)	4 December 2017	13,661	1,809
61	Funtana	<i>Cymodocea nodosa</i> (Mixed)	19 June 2018	31,734	2,039
62	Funtana	<i>Caulerpa cylindracea</i> (Mixed)	19 June 2018	36,488	3,211
63	Funtana	<i>Caulerpa cylindracea</i> (Monospecific)	19 June 2018	41,871	3,851