

Order molecule	Nucleotide sequence (5'→3')	Accession / translated product
Synthetic Insulin Glargine A-Chain Forward Sequence	TATGGGTTCTTCTACCAACCACCACTCTTCTGGTCT GGTGCCGCGTGGTTCTGGCATCTGGAACAGTGTGCACC AGCATCTGCAGCCTGTACCACTGGAAACTACTGCGGCT AAC	GenBank accession number PQ362993 MGSSHHHHHHSSGLVPRGSGIVEQCCTSIQSLYQLENYCG (4352.87 Da)
Synthetic Insulin Glargine A-Chain Reverse Sequence	TCGAGTTAGCCGAGTAGTTTTCCAGCTGGTACAGGCTGCA GATGCTGGTGCAGCACTGTTCCAGATGCCAGAACACGC GGCACCAGACCAGAAGAGTGGTGGTGGTGGTGGAGAA GAACCCA	GenBank accession number PQ362993 MGSSHHHHHHSSGLVPRGSGIVEQCCTSIQSLYQLENYCG (4352.87 Da)
Synthetic Insulin Glargine B-Chain Forward Sequence	TATGGGTTCTTCTACCAACCACCACTCTTCTGGTCT GGTGCCGCGTGGTTCTTTGTGAACAGCATCTGTGCGGC AGCCATCTGGTGAAGCGCTGTACCTGGTGTGCGCGAAC GCGGCTTTTTTACACCCCAAAACCCGCGCTAAC	GenBank accession number PQ362994 MGSSHHHHHHSSGLVPRGSFVNQHLGSHLVEALYLVCGER GFFYTPKTRR (5768.54 Da)
Synthetic Insulin Glargine B-Chain Reverse Sequence	TCGAGTTAGCCGCGGTTTTGGGGTGTAAGAAAGCCGC GTTCCGCCGACACCAAGTACAGCGCTTCCACAGATGGCT GCCGCACAGATGCTGGTTCACAAAAGAACACGCGGCACC AGACCAGAAGAGTGGTGGTGGTGGTGGTGGAGAAGAACCC A	GenBank accession number PQ362994 MGSSHHHHHHSSGLVPRGSFVNQHLGSHLVEALYLVCGER GFFYTPKTRR (5768.54 Da)

S2. Molecular invariants enforced by SSD and ESD

- Input sequences are normalized to unambiguous DNA and coding designs are translated before release.
- SSD releases one complementary forward/reverse synthesis pair; ESD tiles longer targets into multiple ordered complementary pairs with directional internal junctions.
- Recognition sequences and strand cut offsets determine terminal products; overhangs are not stored as pair-specific oligo literals.
- An enzyme supplies an initiating ATG only when the bases retained after cleavage place ATG at the coding boundary.
- The forward and reverse oligonucleotide cores must be exact reverse complements.
- Annealing and ordered ligation must reconstruct the designed molecule on both strands, base for base.
- Internal assembly overhangs must be unique against all used overhangs and their reverse complements; designs widen beyond four bases when required.
- Outer reconstructed ends must equal the selected enzyme products with correct sequence and polarity.
- A cloning vector must contain exactly one site for each selected enzyme; physical vector and insert ends must anneal.
- Coding-junction translation, stop-codon consequences, vector-tag outcomes and site regeneration are reported explicitly.
- Full sequence verification requires complete requested-region coverage and zero confident unexplained differences.

S3. Reproducibility

Execute from the repository root with Python 3.12.13 and the locked project dependencies:

Commands

```
python tools/publication/validate_peptide_and_enzyme_logic.py
python tools/publication/design_glargine_case_study.py --source-record
tools/publication/insulin_glargine_experimental_record.json --output
publication_evidence/glargine_ab_design_and_cloning.json
python tools/publication/validate_insulin_correct_traces.py --root <reference-data-root> --trace-dir
publication_evidence/validated_traces --output-dir publication_evidence/sequencing_validation
python -m pytest gsynth_engine/tests -q
cd django_app && python -m pytest -q
cd ../frontend && npm test -- --run
```

Evidence object	Location	Purpose
Protein-to-clone JSON	publication_evidence/ peptide_and_enzyme_validation.json	Current glargine protein inputs, codon outputs, SSD molecules, duplexes, junctions and hashes
Design/cloning JSON	publication_evidence/ glargine_ab_design_and_cloning.json	Exact molecules, segments, translations, junctions, warnings and source hash
Sequencing JSON	publication_evidence/ sequencing_validation/ glargine_approved_trace_validation.json	Approved-file policy, hashes, primary F/R consensus metrics, post hoc quality sensitivity and Biopython check
Design script	tools/publication/ design_glargine_case_study.py	Executable glargine case study with equality assertions
Trace script	tools/publication/ validate_insulin_correct_traces.py	Executable four-file sequencing analysis and figure generation
Protein workflow script	tools/publication/ validate_peptide_and_enzyme_logic.py	Executable peptide-role, reverse-translation, SSD, cloning and enzyme checks

S4. Unabridged experimental oligonucleotide synthesis and cloning record

The record below preserves the complete experimental procedures relevant to the G-Synth validation. Laboratory notebooks and instrument exports remain the authoritative operational record. Oligonucleotides were synthesized on a MerMade 4 instrument using 1- μ mol, 1000-Å universal supports and DMT-off processing.

Purification used an equal volume of cold butanol, 30 s vortexing, 30 min at -20°C and 15 min centrifugation at 14,000 rpm and 4°C . Pellets were washed twice with 1 mL 70% ethanol at -20°C , centrifuged for 10 min, air-dried and resuspended. Exact rotor radius is not recorded, so relative centrifugal force cannot be reconstructed from rpm; future protocols should report $\times g$.

Hybridization combined 10 μL of each complementary strand with 20 μL 2 $\times$ SSC containing 0.1% SDS. Samples were heated at 95°C for 5 min and cooled to room temperature over 30 min. pET-21a(+) digestion used 1 μg vector, 10 U NdeI, 10 U XhoI and 1 $\times$ CutSmart buffer in 50 μL for 2 h at 37°C . Gel-purified vector was ligated to duplex insert at a nominal 3:1 molar ratio with 50 ng vector and 400 U T4 DNA ligase in 20 μL at 16°C for 2 h followed by 4°C overnight. E. coli DH5 α cells were heat-shock transformed for 50 s at 42°C , recovered in SOC and selected on 100 $\mu\text{g}/\text{mL}$ ampicillin.

Measurement	A forward	A reverse	B forward	B reverse
ssDNA concentration (ng/ μ L)	1702.52	1709.40	1688.51	1703.60
dsDNA concentration by ScanDrop (ng/ μ L)	830	830	1936	1936
dsDNA A260/A280	1.92	1.92	1.88	1.88
Bioanalyzer apparent size (bp)	129	129	173	173
Bioanalyzer concentration (ng/ μ L)	1.15	1.15	3.39	3.39

Caution: UV absorbance, gel mobility and Bioanalyzer sizing assess concentration, purity and approximate fragment size. They do not establish base identity. The different concentration values from ScanDrop and Bioanalyzer reflect method and dilution context and should not be combined without the original dilution records.




Figure S2. Primary wet-laboratory quality-control evidence. Agarose electrophoresis of the synthesized strands and hybridized duplexes, and Bioanalyzer electropherograms, are reproduced from the source experimental record. These assays support presence and approximate size, not nucleotide identity.

Alt text: Primary wet-laboratory quality-control evidence. Agarose electrophoresis of the synthesized strands and hybridized duplexes, and Bioanalyzer electropherograms, are reproduced from the source experimental record. These assays support presence and approximate size, not nucleotide identity.

S5. Complete MerMade 4 synthesis program

Script detail	Reagent / manufacturer	No./V primes	Injection	Reaction	Equalize
INITIALIZATION	INITIALIZATION	INITIALIZATION	INITIALIZATION	INITIALIZATION	INITIALIZATION
3 × Washing	Anhydrous Acetonitrile, (emp BIOTECH, Berlin, Germany)	1 / 10	180 μ L	00:00:02	00:00:12
RUN STEPS	RUN STEPS	RUN STEPS	RUN STEPS	RUN STEPS	RUN STEPS
Deblocking 1	3% Trichloroacetic Acid (TCA) in dichloromethane, (emp BIOTECH, Berlin, Germany)	1 / 10	130 μ L	00:00:02	00:00:59

Script detail	Reagent / manufacturer	No./V primes	Injection	Reaction	Equalize
Deblocking 2	3% Trichloroacetic Acid (TCA) in dichloromethane, (emp BIOTECH, Berlin, Germany)	1 / 10	100 µL	00:00:02	00:00:45
2 × Washing	Anhydrous Acetonitrile, emp BIOTECH	1 / 10	180 µL	00:00:02	00:00:12
Coupling 1 (2 subinjections)	Activator reagent, (emp BIOTECH, Berlin, Germany)	1 / 10	70 µL	00:00:02	00:01:20
Coupling 1 (2 subinjections)	DNA Amidites (A, C, G, T), (Biosearch Technologies, Petaluma, CA, USA)	1 / 10	50 µL	00:00:02	00:01:20
Coupling 2 (2 subinjections)	Activator reagent, emp BIOTECH	1 / 10	50 µL	00:00:02	00:00:40
Coupling 2 (2 subinjections)	DNA Amidites (A, C, G, T), (Biosearch Technologies, Petaluma, CA, USA)	1 / 10	30 µL	00:00:02	00:00:40
Washing	Anhydrous Acetonitrile, (emp BIOTECH, Berlin, Germany)	1 / 10	180 µL	00:00:02	00:00:12
Capping (2 subinjections)	Cap A: Acetic anhydride in THF, emp BIOTECH	1 / 10	65 µL	00:00:02	00:00:50
Capping (2 subinjections)	Cap B: Acetic anhydride in THF, emp BIOTECH	1 / 10	65 µL	00:00:02	00:00:50
Washing	Anhydrous Acetonitrile, (emp BIOTECH, Berlin, Germany)	1 / 10	180 µL	00:00:02	00:00:12
Oxidation	Iodine in water/pyridine/THF (0.02 M), emp BIOTECH	1 / 10	80 µL	00:00:02	00:01:00
2 × Washing	Anhydrous Acetonitrile, (emp BIOTECH, Berlin, Germany)	1 / 10	180 µL	00:00:02	00:00:12
FINALIZATION	FINALIZATION	FINALIZATION	FINALIZATION	FINALIZATION	FINALIZATION
3 × Deblocking	3% Trichloroacetic Acid (TCA) in dichloromethane, (emp BIOTECH, Berlin, Germany)	1 / 10	150 µL	00:00:02	00:00:45
3 × Washing	Anhydrous Acetonitrile, (emp BIOTECH, Berlin, Germany)	1 / 10	180 µL	00:00:02	00:00:12
DEPROTECTION OUTSIDE THE OLIGOSYNTHESIZER	DEPROTECTION OUTSIDE THE OLIGOSYNTHESIZER	DEPROTECTION OUTSIDE THE OLIGOSYNTHESIZER	DEPROTECTION OUTSIDE THE OLIGOSYNTHESIZER	DEPROTECTION OUTSIDE THE OLIGOSYNTHESIZER	DEPROTECTION OUTSIDE THE OLIGOSYNTHESIZER
Cleavage & Deprotection	Concentrated ammonia solution, (Sigma-Aldrich, St. Louis, MO, USA)	/	1 mL	18 h at 70°C	/

S6. Primer record

Primer	Sequence (5'→3')	Reported T _m (°C)	Expected product (bp)
SpAB F	ATGGGTCTCTCACCACCACCA	58.6	123 A / 156 B
SpA R	TTAGCCGAGTAGTTTCCA	52.3	123 A
SpB R	TTAGCGCGGGTTTTGG	53.8	156 B
SeqAB F	TGCATCCATATGGGTCTCTCACCA CCACCA	67.2	144 A / 177 B
SeqA R	TGCATCCTCGAGTTAGCCGAGTAGT TTCCA	65.8	144 A
SeqB R	TGCATCCTCGAGTTAGCGCGGGTT TTTGG	66.4	177 B
T7 F	TAATACGACTCACTATAGGG	52.0	307 A / 340 B
T7 R	GCTAGTTATTGCTCAGCGG	54.3	307 A / 340 B

The source table reported some A/B product sizes together in cells that do not apply to both reverse primers. The corrected product labels above associate SpA R/SeqA R with A and SpB R/SeqB R with B. Before a prospective run, each primer should be re-evaluated in G-Synth against the final recombinant plasmid and the laboratory's polymerase/buffer conditions.

S7. Wet-laboratory PCR and colony-screening evidence




Figure S3. PCR products obtained with insert-specific primers before cloning. SM, GeneRuler Express DNA Ladder; lanes 1–2, A insert; lanes 3–4, B insert.

Alt text: PCR products obtained with insert-specific primers before cloning. SM, GeneRuler Express DNA Ladder; lanes 1–2, A insert; lanes 3–4, B insert.




Figure S4. Colony-screening PCR. Lanes 1 and 3 are A and B products with insert-specific primers; lanes 2 and 4 are the corresponding products with T7 primers.

Alt text: Colony-screening PCR. Lanes 1 and 3 are A and B products with insert-specific primers; lanes 2 and 4 are the corresponding products with T7 primers.

S8. Sequencing evidence manifest

The validation set is limited to A Forward Seq.ab1, A Reverse Seq.ab1, B Forward Seq.ab1 and B Reverse Seq.ab1. Their SHA-256 hashes are recorded in the evidence JSON; no other chromatogram contributes to any reported result. Each F/R pair was oriented and assembled before consensus coverage was evaluated. Consensus coverage, identity, bidirectional overlap and F/R agreement are the primary retrospective endpoints.

Chain	Ref. bp	Consensus cov.	Identity	F/R overlap	F/R agree	Consensus diff.	Biopython cov.
A	123	100.0%	100.0%	56.1%	100.0%	0	100.0%
B	156	100.0%	100.0%	71.2%	100.0%	0	100.0%




Figure S5. Primary G-Synth F/R consensus evidence map. The two oriented reads jointly cover each reference completely; green shading marks bidirectional overlap, with 100% agreement in both constructs.

Alt text: Primary G-Synth F/R consensus evidence map. The two oriented reads jointly cover each reference completely; green shading marks bidirectional overlap, with 100% agreement in both constructs.

S9. Supplementary interface evidence




Figure S6. Primer-template hybridization view. The annealing 3' segment is base-paired; the 5' cloning extension is explicitly unpaired in cycle 1 and appears in the predicted product after extension.

Alt text: Primer-template hybridization view. The annealing 3' segment is base-paired; the 5' cloning extension is explicitly unpaired in cycle 1 and appears in the predicted product after extension.




Figure S7. Detailed SSD duplex hybridization. The two order molecules form a 123-bp exact antiparallel core; the NdeI-derived 5'-TA and XhoI-derived 5'-TCGA terminal products remain exposed and are labelled as cohesive ends.

Alt text: Detailed SSD duplex hybridization. The two order molecules form a 123-bp exact antiparallel core; the NdeI-derived 5'-TA and XhoI-derived 5'-TCGA terminal products remain exposed and are labelled as cohesive ends.





Figure S8. Pre-ligation vector–insert compatibility gate for the current glargine A design. The detailed view preserves enzyme identity, overhang sequence and physical polarity, and labels the joined duplex as an expected product while withholding product analyses.

Alt text: Pre-ligation vector–insert compatibility gate for the current glargine A design. The detailed view preserves enzyme identity, overhang sequence and physical polarity, and labels the joined duplex as an expected product while withholding product analyses.



Alt text: Product state after explicit in-silico ligation. The green status region reports the recombinant length and presents close-up sequence evidence for both closed junctions only after all blocking checks have passed.



Alt text: HindIII-containing diagnostic digest and predicted gel. The restriction site and complete digest fragments are displayed with a named broad-range marker. The panel is labelled as an in-silico prediction.




Figure S11. Restriction map showing HindIII as a named site derived from cut geometry.

Alt text: Restriction map showing HindIII as a named site derived from cut geometry.

S10. Secondary Geneious comparison and independent confirmation

The source experimental study used Benchling for reverse complements and Geneious Prime 2024.0.7 for cloning views and trace alignments. G-Synth independently regenerated the same four synthesis molecules from the two optimized coding inputs, and its recombinant maps placed the inserts downstream of the pET-21a(+) T7/lac expression elements with preserved coding frames. The Geneious figures and G-Synth therefore agree qualitatively on construct identity and placement. A new automated Geneious run was not performed because the reproducibility environment did not include a licensed command-line execution path. Biopython 1.87 was used as the independent, executable alignment comparator.

A prospective head-to-head study should freeze one set of reference plasmids and raw trace files, define identical trimming thresholds and variant rules, blind sample labels, and compare: read placement; orientation calls; covered intervals; mismatches/indels; consensus; runtime; manual interventions; exportability; and final verdict. The present study is a functional cross-check, not a statistical superiority benchmark.




Figure S12. Geneious Prime 2024.0.7 duplex representation of NdeI/XhoI-compatible A- and B-chain inserts from the experimental study. This is secondary qualitative concordance; G-Synth SSD is the primary in-silico reconstruction.

Alt text: Geneious Prime 2024.0.7 duplex representation of NdeI/XhoI-compatible A- and B-chain inserts from the experimental study. This is secondary qualitative concordance; G-Synth SSD is the primary in-silico reconstruction.




Figure S13. Geneious Prime construct annotation for the glargine A- and B-chain pET-21a(+) designs from the experimental study. G-Synth generated the primary coordinate, ORF and junction validation reported in the main article.

Alt text: Geneious Prime construct annotation for the glargine A- and B-chain pET-21a(+) designs from the experimental study. G-Synth generated the primary coordinate, ORF and junction validation reported in the main article.




Figure S14. Geneious Prime alignment made with the same four author-approved construct-confirmation files used by G-Synth. It provides secondary visual concordance and was not treated as an independent blinded benchmark.

Alt text: Geneious Prime alignment made with the same four author-approved construct-confirmation files used by G-Synth. It provides secondary visual concordance and was not treated as an independent blinded benchmark.

S11. Release and submission checklist

- Obtain written approval from every listed author for author order, affiliations, CRediT roles and AI disclosure.
- Run prospective bidirectional sequencing until both junctions and 100% of each insert are covered at the predefined quality threshold.
- Deposit raw trace files, reference FASTA/GenBank files, evidence JSON and analysis scripts in a stable repository, subject to institutional approval.
- Publish the version 1.0.0 GitHub release, connect Zenodo and insert the versioned DOI in the manuscript and CITATION.cff.
- Perform a clean-clone execution of all three test suites and both publication scripts; retain logs and environment versions.
- Prepare the target journal's graphical summary, high-resolution figures and machine-readable constructs; confirm its current author guidelines and retain the limitation that functional insulin glargine was not demonstrated.