

OODF: Optimized Opacity Distribution Functions for a New Generation of Solar and Stellar Brightness Variability Models

Miha Cernetic, Veronika Witzke, Alexander I. Shapiro, Natalia A. Krivova, Sami K. Solanki,
Rinat V. Tagirov

Max Planck Institute for Solar System Research
SOLVe group
miha.cernetic@mps.mpg.de

Mar. 22, '18



MAX-PLANCK-GESellschaft




SOLVe




Exemplary case of Strömgren y filter

- Complex structure of spectrum due to lines




Complex structure of opacity

- Opacity varies by multiple orders of magnitude within 1\AA




Sorting by opacity

- Sort wavelength points by corresponding values of opacity




Approximating the sorted values

- Approximate opacity with a stepwise function




High resolution spectrum and ODF spectrum

- Use the stepwise opacity to calculate the flux




High resolution spectrum and ODF spectrum

- Use the stepwise opacity to calculate the flux
- 4 sub bins -> This two integrals differ by just 2%




Generating ODFs

- Start with high resolution opacity




Generating ODFs

- Sort wavelength points by corresponding values of opacity; monotonically increasing opacity
- Integral is preserved by sorting




Generating ODFs

- All wavelength information within the bin is lost




Generating ODFs - Example with 10 uniform sub bins

- Approximate the sorted opacity with a step-wise function




ODF generation process

- Mean is skewed by extreme values




ODF performance analysis

- Synthesize spectrum using ODFs from 1000-9000Å with 10Å bins
- Compare the fluxes from the ODF spectrum with the high resolution spectrum in the bins




Analysis of different ODFs

- Uniform ODFs




Analysis of different ODFs

- Nonuniform ODFs
- The last sub bin is crucial after 5000Å




Comparison of nonuniform sub bins


- Legend specifies sub bin sizes, starting with the first one
- Last sub bin is the same for all




Best sub bin combinations using 3 sub bins



Best combinations of 3 sub bins for Strömgren y



Best combination of 3 sub bins for Strömgren y



Speedups in the case of Strömgren y

- Interval length: $\sim 1000\text{\AA}$

High resolution: 80 points per $\text{\AA} \sim 80\,000$ points

ODF: 12 points per $10\text{\AA} \sim 1200$ points
speedup 67 times

OODF: 3 points per $1000\text{\AA} \sim 3$ points
speedup 25 000 times



Conclusions

- An efficient procedure for radiative transfer is timely for new generation of solar and stellar variability models.
- We developed a novel method for fast spectral synthesis.
- Can be tailored for different filters: Strömgren $b + y$, Kepler, PLATO and others.
- Significant speed up relative to standard methods by a factor of at least two orders of magnitude.



Conclusions

- An efficient procedure for radiative transfer is timely for new generation of solar and stellar variability models.
- We developed a novel method for fast spectral synthesis.
- Can be tailored for different filters: Strömgren $b + y$, Kepler, PLATO and others.
- Significant speed up relative to standard methods by a factor of at least two orders of magnitude.

Thank you for your attention!

