

# Developing faster and more accurate radiative transfer calculations for solar and stellar applications

Miha Cernetic, Alexander I. Shapiro, Veronika Witzke, Natalia A. Krivova, Sami K. Solanki, Rinat V. Tagirov

Max Planck Institute for Solar System Research  
SOLVe group  
✉ cernetic@mps.mpg.de

Feb. 26, '19



## First project - Introduction


# **OODF: Optimized Opacity Distribution Functions for a New Generation of Solar and Stellar Brightness Variability Models**

Miha Cernetic, Alexander I. Shapiro, Veronika Witzke, Natalia A. Krivova, Sami K. Solanki,  
Rinat V. Tagirov




# Total Solar Irradiance

- TSI – spectrally integrated solar radiative flux at 1 AU from the sun



## 1.5D simulations


## 3D MHD simulations with MURaM



Structure of the magnetic features

+


## 1.5D radiative transfer



Spectra of the magnetic features


## Spectra of the individual components

### Observed solar spectra




## Importance of lines for variability

- TSI – Total Solar Irradiance, i.e. integrated over wavelengths
- SSI – Spectral Solar Irradiance, depends on wavelength
- $\Delta$  – difference between the solar minima and maxima




## Importance of lines for variability



## Importance of lines for variability


- 25% of the variability comes from molecular lines → accurate treatment of lines is crucial






## Generating ODFs

- Start with high resolution opacity




## Generating ODFs

- Sort wavelength points by corresponding values of opacity; monotonically increasing opacity
- Integral is preserved by sorting




## Generating ODFs

- All wavelength information within the bin is lost




## Generating ODFs - Example with 10 uniform sub bins

- Approximate the sorted opacity with a step-wise function




## ODF generation process

- Mean is skewed by extreme values




## ODF performance analysis

- Synthesize spectra with **NESSY** radiative transfer code using ODFs from 1000-9000Å in 10Å bins
- Compare the fluxes from the ODF spectrum with the high resolution spectrum in the bins




## Possible solutions

Number of calculations  $n_{\text{cal}} = n_{\text{bins}} \times n_{\text{subbins}}$



# Analysis of different ODFs


- Uniform ODFs







## Analysis of different ODFs

- Nonuniform ODFs
- The last sub bin is crucial after 5000Å




## Best sub bin combinations using 4 sub bins



Strömgren  $b$ 

## OODF treatment of filters

- ODF approach can be generalized to take into account any shape of transmission function
- Achieved by transforming the wavelength grid according to the transmission function



## Speedups in the case of Strömgren $b$

- Interval length (transmission curve  $> 1\%$ ):  $\sim 400\text{\AA}$

High resolution: 80 points per  $\text{\AA} \sim 32\,000$  points

ODF: 12 points per  $10\text{\AA} \sim 480$  points  
speedup 67 times


ODF: 4 points for the whole bin  
speedup  $\sim 8\,000$  times



## MSc project - Introduction

Title: **Fast Approximate non-LTE Treatment of the Chromosphere**

- The aim of the magnetohydrodynamic group at my institute is simulating the Sun from the upper part of the convection zone and up



[http://www.phy.cuhk.edu.hk/elearning/Solar\\_eclipse/Solar\\_structure.html](http://www.phy.cuhk.edu.hk/elearning/Solar_eclipse/Solar_structure.html)



## MSc project - Fast approximate non-LTE treatment of the chromosphere

### **Current solutions:**

- Currently used solution: pre-tabulated cooling rates derived from 1D RADYN radiation-hydrodynamic simulations



## MSc project - Fast approximate non-LTE treatment of the chromosphere

### Current solutions:

- Currently used solution: pre-tabulated cooling rates derived from 1D RADYN radiation-hydrodynamic simulations → difficult to expand to more than a handful of elements





## MSc project - Fast approximate non-LTE treatment of the chromosphere

### Current solutions:

- Currently used solution: pre-tabulated cooling rates derived from 1D RADYN radiation-hydrodynamic simulations → difficult to expand to more than a handful of elements

### My work:

- Escape probability methods



## MSc project - Fast approximate non-LTE treatment of the chromosphere

### Current solutions:

- Currently used solution: pre-tabulated cooling rates derived from 1D RADYN radiation-hydrodynamic simulations → difficult to expand to more than a handful of elements

### My work:

- Escape probability methods → problem with going from 1D to 3D



## MSc project - Fast approximate non-LTE treatment of the chromosphere

### Current solutions:

- Currently used solution: pre-tabulated cooling rates derived from 1D RADYN radiation-hydrodynamic simulations → difficult to expand to more than a handful of elements

### My work:

- Escape probability methods → problem with going from 1D to 3D
- Scharmer method: *A new approach to multi-level non-LTE radiative transfer problems, Scharmer's et al (1985)*
  - Eddington Barbier approximation + core saturation approximation



## Conclusions

### First project

- We developed a novel method for fast spectral synthesis.
- Found optimal sub bins for different wavelength regimes.
- Can be tailored for different filters: Strömgren  $b + y$ , Kepler, PLATO and others.
- Significant speed up relative to standard methods by a factor of at least two orders of magnitude.

### Second project


- Escape probabilities: try the 1.5D approach.
- Scharmer: implement the method into RH
- Compare both methods above to the pre-tabulated cooling rates.




Backup slides beyond

**Here be dragons!**



Best combinations of 4 sub bins for Strömgren  $b$ 

Formula: value weighted by the derivative



## Ascending vs descending sort

