

# OODF: Optimized Opacity Distribution Functions for a New Generation of Solar and Stellar Brightness Variability Models

Miha Cernetic, Alexander I. Shapiro, Veronika Witzke, Natalia A. Krivova, Sami K. Solanki, Rinat V. Tagirov


Max Planck Institute for Solar System Research  
SOLVe group  
✉ cernetic@mps.mpg.de

Aug. 23, '18




# Total Solar Irradiance

- TSI – spectrally integrated solar radiative flux at 1 AU from the sun




# Spectra of the individual components

## Solar spectra




## Importance of lines for variability

- TSI – Total Solar Irradiance, i.e. integrated over wavelengths
- SSI – Spectral Solar Irradiance, depends on wavelength
- $\Delta$  – difference between the solar minima and maxima




## Importance of lines for variability




## Importance of lines for variability

- 25% of the variability comes from molecular lines → accurate linelists are required



## 1.5D simulations


## 3D MHD simulations with MURaM



Structure of the magnetic features

+


## 1.5D radiative transfer



Spectra of the magnetic features

## Generating ODFs


- Start with high resolution opacity






## Generating ODFs

- Sort wavelength points by corresponding values of opacity; monotonically increasing opacity
- Integral is preserved by sorting




## Generating ODFs

- All wavelength information within the bin is lost




## Generating ODFs - Example with 10 uniform sub bins

- Approximate the sorted opacity with a step-wise function




## ODF generation process

- Mean is skewed by extreme values




## ODF performance analysis

- Synthesize spectra using ODFs from 1000-9000Å in 10Å bins
- Compare the fluxes from the ODF spectrum with the high resolution spectrum in the bins




## Possible solutions

Number of calculations goes as:  $n_{bins} \times n_{subbins}$




# Analysis of different ODFs

- Uniform ODFs




## Analysis of different ODFs

- Nonuniform ODFs
- The last sub bin is crucial after 5000Å






## Best sub bin combinations using 4 sub bins




## Best sub bin combinations using 10 sub bins



## Best combination of 4 sub bins for Strömgren b

- Total line contribution  $\sim 15\%$



## Speedups in the case of Strömgren $b$

- Interval length:  $\sim 400\text{\AA}$

High resolution: 80 points per  $\text{\AA} \sim 32\,000$  points

ODF: 12 points per  $10\text{\AA} \sim 480$  points  
speedup 67 times

OOF: 3 points for the whole bin  $\text{\AA} \sim 3$  points  
speedup  $\sim 11\,000$  times



## Conclusions

- We developed a novel method for fast spectral synthesis.
- Found optimal sub bins for different wavelength regimes.
- Can be tailored for different filters: Strömgren  $b + y$ , Kepler, PLATO and others.
- Significant speed up relative to standard methods by a factor of at least two orders of magnitude.




## Conclusions


- We developed a novel method for fast spectral synthesis.
- Found optimal sub bins for different wavelength regimes.
- Can be tailored for different filters: Strömgren  $b + y$ , Kepler, PLATO and others.
- Significant speed up relative to standard methods by a factor of at least two orders of magnitude.

Thank you for your attention!



Best combinations of 4 sub bins for Strömgren  $b$ 

Formula: value weighted by the derivative





## Ascending vs descending sort

