EAB 770 - Chapter 13

Miguel Fudolig

1 Categorized Time-to-Event Data

1.1 Outline

- Categorical Time-to-Event Data
- Life Table Estimation
- Mantel-Cox methods
- Piecewise Exponential Methods

1.2 Time-to-Event Data

Time-to-event data are generated from time to treatment/exposure until some event as their outcome.

> **Note**
>
> Sometimes, you can only determine the interval of time during which an event occurs. Examples include evaluating animals every four hours after exposure to bacteria and examining patients every half a year for the recurrence of their medical condition.
> Data from these studies are referred to as *grouped survival data* or *categorized survival data*.

1.3 Withdrawals

Withdrawals are instances when subjects leave the study before the study ends.

> **Note**
>
> Withdrawals are assumed to be independent of the condition being studied and that multiple withdrawals occur uniformly throughout the interval. After each interval between successive evaluations, the number of withdrawals are also measured.

> **Important**
>
> The survival status of subjects who withdrew is unknown, hence the data ends up being *censored*.

2 Life Table Estimation: Survival Rates

2.1 Survival Rates

We are often interested in estimating the survival rates for each method.

> **Note**
>
> The survival rate is the proportion of individuals in a study who have not experienced the event of interest.

The survivor function $S(y)$, defined to be the probability that the survival time is greater than or equal to y, can be expressed as

$$S(y) = 1 - F(y) = P(Y \geq y)$$

where Y is the random variable that denotes the continuous lifetime of a subject and $F(y)$ is the cumulative incidence function.

> **Note**
>
> The cumulative incidence function is defined as the cumulative probability of an event occurring before time y. This function is commonly chosen to be the Weibull distribution, but there are other functions that people can use.

2.2 Hazard Function

> **Note**
>
> The hazard function expresses the risk or hazard of the event in question occurring at some time y. This function is often referred to as $h(y)$.
> The cumulative hazard function is the cumulative risk of an event occurring by time y. This function is often referred to as $H(y)$.

The hazard function is related to the survival rate through the following equation:

$$S(y) = \exp\left(-\int_0^y h(t)dt\right) = \exp[-H(y)]$$

where $H(y)$ is the cumulative hazard function.

> **Note**
>
> $H(y)$ is the negative logarithm of the survivor function.

2.3 Life Table

The life table or actuarial method calculates the survival rates by determining the number of subjects at risk for each time interval. The number of subjects with no recurrence for each interval can be counted when the following is known:

- Number of withdrawals
- Number of recurrences
- Number of subjects who survived with no recurrence

2.4 Life Table

Each treatment group will have a life table with the following format:

Interval	No Events	Events	Withdrawals	At Risk

> **Note**
>
> While the Kaplan-Meier method is preferable in most cases, the life table method can be used for large data sets and grouped survival data.

2.5 Life Table Estimation

> **Note**
>
> Define n_{ijk} to be the number of patients in the ith group with the jth status for the kth time interval. $j = 0$ corresponds to no recurrence, $j = 1, 2$ corresponds to those with recurrence and those withdrawn during the interval, respectively. The interval can take the following values: $k = 1, 2, ..., t$

The number of patients in the ith group who have no recurrence at the kth interval can be expressed as:

$$n_{i0k} = n_{i0t} + \sum_{j=1}^{2} \sum_{g=k+1}^{t} n_{ijg}$$

The first term includes those who have no recurrence at time t, while the second term accounts for those with recurrence and withdrawn from $k + 1$ to t interval.

2.6 Survival Probability

The life table estimates for the probability of surviving at least k intervals are computed as:

$$G_{ik} = \prod_{g=1}^{k} \frac{n_{i0g} + 0.5n_{i2g}}{n_{i0g} + 0.5n_{i2g} + n_{i1g}} = \prod_{g=1}^{k} p_{ig}$$

2.6.1 Notes

> **Important**
>
> p_{ig} corresponds to the estimated conditional probability for surviving the gth interval given that survival of all preceding intervals has occurred.
> The term $0.5n_{i2g}$ accounts for the assumptions that the withdrawals are uniform throughout the interval and that withdrawals are not to have the event at the time of withdrawal.

2.6.2 Standard Error

The corresponding standard error of G_{ik} is:

$$se(G_{ik}) = G_{ik}\sqrt{\sum_{g=1}^{k}\left[\frac{(1-p_{ig})}{n_{i0g}+0.5n_{i2g}}\right]}$$

2.7 SAS Implementation

The LIFETEST procedure provides life table analysis for categorized time-to-event data.

> **Note**
>
> LIFETEST calculates survival estimates according to the Kaplan-Meier method by default, so the life table needs to be specified using the `method=lt` option in the `proc lifetest` statement.

Shown below is the sample code:

```
proc lifetest data=data method=lt plots=(s,ls) intervals=0 to k by 1;
freq count;
strata treatment;
time interval*censor(1);
run;
```

2.8 SAS: Comparison across strata

Survival curves are commonly compared across strata (treatment). SAS provides three different tests to test for a difference between the curves across the strata. This is found in the "Test of Equality over Strata" table

2.8.1 Log-Rank

The **Log-Rank** test is a nonparametric tests that require no assumption for the event time distribution. This test is more powerful when the two groups have similar rates of early events, with one having more long-term survivors than the other.

2.8.2 Wilcoxon

The **Wilcoxon** test is also a non-parametric test that is more powerful when one group has a tendency to have fewer early events and longer survival times.

2.8.3 LR

The **Likelihood Ratio** (-2LogLR) test is a parametric test that works best under the assumption that the time distribution is exponential. The LR test is best if there is a linear trend through the origin, indicating consistency with exponential event times.

2.8.4 Modified Peto-Peto Test

The Modified Peto-Peto Test can be used when survival curves cross. This test is a variation of the log-rank test that assumes weights that detect earlier event differences. This can be specified in the `STRATA` statement using `test=modpeto`.

2.9 `R` Implementation

Survival analysis in `R` is geared towards the Kaplan-Meier method compared to life tables. The `survival` and the `KMsurv` packages can be used to implement the life table analysis and create Kaplan-Meier curves like `SAS` with the following caveats:

2.9.1 Kaplan-Meier Curves

- The data needs to be disaggregated, i.e. we need to use `uncount()` to get "individual-level" data, to create survival curves from the `survfit()` function in the `survival` package.
- The `event` and `time` variables need to be defined as the following: the `time` variable should correspond to the mid point of each interval, while the `event` variable is defined as `1-censor`.

```r
library(survival)

data$time <- data$interval+0.5
data$event <- 1-data$censor

fit <- survfit(Surv(time, event) ~ group, data = data)
plot(fit, lty = 1:2, xlab = "Interval", ylab = "Survival")
```

2.9.2 Life Table

The aggregated counts have to be introduced using the `lifetab()` function in the `KMsurv` package in `R`.

```
library(KMsurv)
lifetab(tis, # Time points that will be used as intervals
        ninit, # Initial value for the group
        nlost, # Vector of censored for different intervals
        nevent # Vector of all events for different intervals
        )
```

2.9.3 Equality Tests for Strata

The function `survdiff()` can do a log-rank test and the modified Peto-Peto test.

> **Warning**
>
> `R` cannot perform the Wilcoxon and LR tests using a specific function in the `survival` and `KMsurv` packages.

```
library(survival)
survdiff(Surv(time,event)~group,data=data,rho=0) # Log-rank
survdiff(Surv(time,event)~group,data=data,rho=1) # Log-rank
```

2.10 Example

Investigators were interested in comparing an active and control treatment to prevent the recurrence of a medical condition that had been healed. They applied a diagnostic procedure at the end of the first, second, and third years to determine whether there was a recurrence. Use the life table method to test if there is a difference in survival curves between the active and control treatments.

2.10.1 SAS Data Set

The variable `INTERVAL` takes the value 0, 1, 2, or 3, corresponding to the following intervals: <1 Year, 1-2 Year, 2-3 Years, >3 Years (no recurrence at the end of the study). and the variable `CENSOR` takes the value 1 for censored subjects and 0 otherwise. The variable `TREATMENT` reflects whether the subjects received the active treatment or acted as controls, and the variable `COUNT` contains the frequency counts.

```
data medical;
input interval treatment $ censor count @@;
datalines;
0 control 1 9 0 control 0 15
1 control 1 7 1 control 0 13
2 control 1 6 2 control 0 7
3 control 1 17
0 active 1 9 0 active 0 12
1 active 1 3 1 active 0 7
2 active 1 4 2 active 0 10
3 active 1 45
;
```

2.10.2 R Data Set

```
library(tidyverse)
```

```
-- Attaching core tidyverse packages ------------------------ tidyverse 2.0.0 --
v dplyr     1.2.1     v readr     2.2.0
v forcats   1.0.1     v stringr   1.6.0
v ggplot2   4.0.3     v tibble    3.3.1
v lubridate 1.9.5     v tidyr     1.3.2
v purrr     1.2.2
-- Conflicts ------------------------------------------ tidyverse_conflicts() --
x dplyr::filter() masks stats::filter()
x dplyr::lag()    masks stats::lag()
i Use the conflicted package (<http://conflicted.r-lib.org/>) to force all conflicts to becor
```

```
medical_counts <- data.frame(
  interval  = c(0, 0, 1, 1, 2, 2, 3,  0, 0, 1, 1, 2, 2, 3),
  treatment = rep(c("control", "active"), each = 7),
  censor    = c(1, 0, 1, 0, 1, 0, 1,  1, 0, 1, 0, 1, 0, 1),
  count     = c(9, 15, 7, 13, 6, 7, 17,  9, 12, 3, 7, 4, 10, 45)
)

medical<- medical_counts|>
  uncount(weights = count)


medical$time  <- medical$interval + 0.5
medical$event <- 1 - medical$censor
```

2.10.3 SAS Code

```
proc lifetest data=medical method=lt plots=(s,ls) intervals=0 to 3 by 1;
freq count;
strata treatment/test=(logrank lr modpeto wilcoxon);
time interval*censor(1);
run;
```

2.10.4 R Code

```
library(survival)
library(KMsurv)
```

```
Warning: package 'KMsurv' was built under R version 4.6.1
```

```
medical_counts |>
  group_by(treatment) |>
  summarize(sum=sum(count))
```

```
# A tibble: 2 x 2
  treatment   sum
  <chr>     <dbl>
1 active       90
2 control      74
```

```
# Kaplan-Meier curves (SAS plots=(s,ls))
fit <- survfit(Surv(time, event) ~ treatment, data = medical)
summary(fit)
```


```
Call: survfit(formula = Surv(time, event) ~ treatment, data = medical)

                treatment=active
 time n.risk n.event survival std.err lower 95% CI upper 95% CI
  0.5     90      12    0.867  0.0358        0.799        0.940
  1.5     69       7    0.779  0.0450        0.695        0.872
  2.5     59      10    0.647  0.0534        0.550        0.760

                treatment=control
```

```
 time n.risk n.event survival std.err lower 95% CI upper 95% CI
  0.5     74      15    0.797  0.0467        0.711        0.894
  1.5     50      13    0.590  0.0603        0.483        0.721
  2.5     30       7    0.452  0.0649        0.341        0.599
```

```
plot(fit, lty = 1:2, xlab = "Interval", ylab = "Survival")
legend("topright", legend = c("Active", "Control"), lty = 2:1)
```




```
dev.off()
```

```
null device
          1
```

```
# Actuarial life-table estimates (SAS method=lt, intervals=0 to 3 by 1)
lt_control <- lifetab(tis = 0:4, ninit = 74,
                      nlost = c(9, 7, 6, 17), nevent = c(15, 13, 7, 0))
lt_active  <- lifetab(tis = 0:4, ninit = 90,
                      nlost = c(9, 3, 4, 45), nevent = c(12, 7, 10, 0))
lt_control
```

```
    nsubs nlost nrisk nevent      surv       pdf    hazard    se.surv
```

```
0-1    74     9  69.5     15 1.0000000 0.2158273 0.2419355 0.00000000
1-2    50     7  46.5     13 0.7841727 0.2192311 0.3250000 0.04934767
2-3    30     6  27.0      7 0.5649416 0.1464663 0.2978723 0.06266907
3-4    17    17   8.5      0 0.4184753        NA        NA 0.06652107
        se.pdf  se.hazard
0-1 0.04934767 0.06200874
1-2 0.05342128 0.08894071
2-3 0.05033961 0.11132948
3-4         NA         NA
```

```
lt_active
```

```
    nsubs nlost nrisk nevent      surv       pdf    hazard    se.surv
0-1    90     9  85.5     12 1.0000000 0.1403509 0.1509434 0.00000000
1-2    69     3  67.5      7 0.8596491 0.0891488 0.1093750 0.03756515
2-3    59     4  57.0     10 0.7705003 0.1351755 0.1923077 0.04638160
3-4    45    45  22.5      0 0.6353248        NA        NA 0.05449140
        se.pdf  se.hazard
0-1 0.03756515 0.04344933
1-2 0.03213710 0.04127800
2-3 0.03965962 0.06053125
3-4         NA         NA
```

```
# Log-rank Test

survdiff(Surv(time, event) ~ treatment, data = medical, rho = 0)
```

```
Call:
survdiff(formula = Surv(time, event) ~ treatment, data = medical,
    rho = 0)

                  N Observed Expected (O-E)^2/E (O-E)^2/V
treatment=active 90       29     37.7      2.00      5.88
treatment=control 74      35     26.3      2.87      5.88

 Chisq= 5.9  on 1 degrees of freedom, p= 0.02
```

```
# Modified Peto Test

survdiff(Surv(time, event) ~ treatment, data = medical, rho = 1)
```

```
Call:
survdiff(formula = Surv(time, event) ~ treatment, data = medical,
    rho = 1)

                   N Observed Expected (O-E)^2/E (O-E)^2/V
treatment=active  90     24.8     32.3      1.76       5.7
treatment=control 74     30.7     23.2      2.45       5.7

 Chisq= 5.7  on 1 degrees of freedom, p= 0.02
```

2.10.5 Answer

Based on the survival curves, the log-rank test might be preferred ($\chi^2 = 5.88$, p=0.015). At a significance level of 0.05, there is sufficient evidence of a difference between the survival curves between the two treatments.

2.11 Exercise

A study about mild traumatic brain injuries (mTBI) investigated the persistence of headaches for those who presented with mental health related sequelae and those who did not. Use the Modified Peto-Peto test and the life table method to test if there is a difference between the survival curves of the two profiles.

The variable `INTERVAL` takes the value 0, 1, 2, corresponding to the following intervals: <6 months, 6-12 months, >12 months (no occurrence at the end of the study). and the variable `CENSOR` takes the value 1 for censored subjects and 0 otherwise. The variable `PROFILE` reflects whether the subjects are part of the group that presented with mental health sequelae (MH) and those that did not (NoMH), and the variable `COUNT` contains the frequency counts.

2.11.1 SAS Data Set

```
data mTBI;
input interval profile $ censor count @@;
datalines;
0 NoMH 1 1 0 NoMH 0 17
1 NoMH 1 6 1 NoMH 0 9
2 NoMH 1 20
0 MH 1 8 0 MH 0 15
1 MH 1 2 1 MH 0 7
2 MH 1 10
;
```

2.11.2 R Data Set

```
library(tidyverse)
mTBI_counts <- data.frame(
  interval = c(0, 0, 1, 1, 2, 0, 0, 1, 1, 2),
  profile  = c("NoMH", "NoMH", "NoMH", "NoMH", "NoMH",
                       "MH", "MH", "MH", "MH", "MH"),
  censor   = c(1, 0, 1, 0, 1, 1, 0, 1, 0, 1),
  count    = c(1, 17, 6, 9, 20, 8, 15, 2, 7, 10)
)

mTBI_counts
```

```
   interval profile censor count
1         0    NoMH      1     1
2         0    NoMH      0    17
3         1    NoMH      1     6
4         1    NoMH      0     9
5         2    NoMH      1    20
6         0      MH      1     8
7         0      MH      0    15
8         1      MH      1     2
9         1      MH      0     7
10        2      MH      1    10
```

```
mTBI <- mTBI_counts |>
  uncount(weights=count)

mTBI$time  <- mTBI$interval + 0.5
mTBI$event <- 1 - mTBI$censor
```

2.11.3 SAS Code

```
proc lifetest data=mTBI method=lt plots=(s,ls)
intervals=0 to 2 by 1;
freq count;
strata profile/test=(logrank lr modpeto wilcoxon);
time interval*censor(1);
run;
```

2.11.4 R Code

```
library(survival)
library(KMsurv)

# Initial Counts

mTBI_counts |>
  group_by(profile) |>
  summarize(sum=sum(count))
```

```
# A tibble: 2 x 2
  profile   sum
  <chr>   <dbl>
1 MH         42
2 NoMH       53
```


```
# Kaplan-Meier curves (SAS plots=(s,ls))
fit <- survfit(Surv(time, event) ~ profile, data = mTBI)
summary(fit)
```

```
Call: survfit(formula = Surv(time, event) ~ profile, data = mTBI)

                profile=MH
 time n.risk n.event survival std.err lower 95% CI upper 95% CI
  0.5     42      15    0.643  0.0739        0.513        0.805
  1.5     19       7    0.406  0.0851        0.269        0.612

                profile=NoMH
 time n.risk n.event survival std.err lower 95% CI upper 95% CI
  0.5     53      17    0.679  0.0641        0.565        0.817
  1.5     35       9    0.505  0.0692        0.386        0.660
```

```
plot(fit, lty = 1:2, xlab = "Interval", ylab = "Survival")
legend("topright",legend = c("NoMH", "MH"), lty = 2:1)
```




```
lt_NoMH <- lifetab(tis = 0:3, ninit = 53,
                   nlost = c(1,6,20), nevent = c(17,9,0))
lt_MH  <- lifetab(tis = 0:3, ninit = 42,
                   nlost = c(8,2,10), nevent = c(15,7,0))
lt_NoMH
```

```
    nsubs nlost nrisk nevent      surv       pdf    hazard    se.surv
0-1    53     1  52.5     17 1.0000000 0.3238095 0.3863636 0.00000000
1-2    35     6  32.0      9 0.6761905 0.1901786 0.3272727 0.06458024
2-3    20    20  10.0      0 0.4860119        NA        NA 0.07101373
        se.pdf  se.hazard
0-1 0.06458024 0.09194179
1-2 0.05673015 0.10762044
2-3         NA         NA
```

```
lt_MH
```

```
    nsubs nlost nrisk nevent      surv       pdf    hazard    se.surv
0-1    42     8    38     15 1.0000000 0.3947368 0.4918033 0.00000000
1-2    19     2    18      7 0.6052632 0.2353801 0.4827586 0.07929287
2-3    10    10     5      0 0.3698830        NA        NA 0.08476377
        se.pdf se.hazard
```

```
0-1 0.07929287 0.1230840
1-2 0.07607697 0.1770703
2-3         NA        NA
```

```r
# Log-rank Test

survdiff(Surv(time, event) ~ profile, data = mTBI, rho = 0)
```

```
Call:
survdiff(formula = Surv(time, event) ~ profile, data = mTBI,
    rho = 0)

              N Observed Expected (O-E)^2/E (O-E)^2/V
profile=MH   42       22     19.8     0.250     0.625
profile=NoMH 53       26     28.2     0.175     0.625

 Chisq= 0.6  on 1 degrees of freedom, p= 0.4
```

```r
# Modified Peto Test

survdiff(Surv(time, event) ~ profile, data = mTBI, rho = 1)
```

```
Call:
survdiff(formula = Surv(time, event) ~ profile, data = mTBI,
    rho = 1)

              N Observed Expected (O-E)^2/E (O-E)^2/V
profile=MH   42     19.6     17.9     0.174     0.482
profile=NoMH 53     23.0     24.7     0.125     0.482

 Chisq= 0.5  on 1 degrees of freedom, p= 0.5
```

2.11.5 Answer

There is insufficient evidence of a difference between the survival curves of the two groups based on the modified Peto-Peto test ($\chi^2 = 0.5012$, p=0.48) and the Wilcoxon test ($\chi^2 = 0.4339$, p=0.51) .

2.12 Mantel-Cox Test

The Mantel-Cox test tests the null hypothesis that the survival rates are the same across the groups. The Mantel-Haenszel methodology was extended to survival data by formatting the frequencies as a set of 2x2 life tables and using the Mantel-Haenszel computations for these sets of tables.

> **Important**
>
> The treatment/groups define the rows, recurrence status define the columns, and intervals are the strata, under the assumption that the time interval results are uncorrelated. Withdrawals are grouped with the no recurrence group or eliminated entirely.

> **Note**
>
> The Mantel-Cox test for grouped data is equivalent to the log-rank test for comparing survival curves for ungrouped data.

2.13 Example

Investigators were interested in comparing an active and control treatment to prevent the recurrence of a medical condition that had been healed. They applied a diagnostic procedure at the end of the first, second, and third years to determine whether there was a recurrence. Use the Mantel-Cox method to test if there is a difference in survival rates between the active and control treatments.

2.13.1 SAS Data Set

The data set excludes censored data.

```
data medical_mc;
input time $ treatment $ status $ count @@;
datalines;
0-1 control recur 15 0-1 control not 50
0-1 active recur 12 0-1 active not 69
1-2 control recur 13 1-2 control not 30
1-2 active recur 7 1-2 active not 59
2-3 control recur 7 2-3 control not 17
2-3 active recur 10 2-3 active not 45
;
```

2.13.2 R Data Set

```
medical_mc <- data.frame(
  time      = c("0-1", "0-1", "0-1", "0-1", "1-2", "1-2", "1-2", "1-2", "2-3", "2-3", "2-3",
  treatment = c("control", "control", "active", "active", "control", "control", "active", "a
  status    = c("recur", "not", "recur", "not", "recur", "not", "recur", "not", "recur", "no
  count     = c(15, 50, 12, 69, 13, 30, 7, 59, 7, 17, 10, 45)
)
```

2.13.3 SAS Code

```
proc freq data=medical_mc order=data;
weight count;
tables time*treatment*status / cmh;
run;
```

2.13.4 R Code

```
library(vcdExtra)
```

```
Warning: package 'vcdExtra' was built under R version 4.6.1
```

```
Loading required package: vcd
```

```
Warning: package 'vcd' was built under R version 4.6.1
```

```
Loading required package: grid
```

```
Loading required package: gnm
```

```
Warning: package 'gnm' was built under R version 4.6.1
```

```
Registered S3 methods overwritten by 'vcdExtra':
  method      from
  pairs.table vcd
  print.Kappa vcd
```

```
Attaching package: 'vcdExtra'

The following objects are masked from 'package:vcd':

    pairs_diagonal_mosaic, woolf_test
```

```
med_cc <- xtabs(count~treatment+status+time,data=medical_mc)
med_cc
```

```
, , time = 0-1

         status
treatment not recur
  active   69    12
  control  50    15

, , time = 1-2

         status
treatment not recur
  active   59     7
  control  30    13

, , time = 2-3

         status
treatment not recur
  active   45    10
  control  17     7
```

```
CMHtest(med_cc,
        strata="time",
        overall=T)$ALL
```

```
Cochran-Mantel-Haenszel Statistics for treatment by status
    Overall tests, controlling for all strata

                AltHypothesis  Chisq Df      Prob
cor        Nonzero correlation 8.0294  1 0.0046024
rmeans  Row mean scores differ 8.0294  1 0.0046024
```

```
cmeans  Col mean scores differ 8.0294  1 0.0046024
general    General association 8.0294  1 0.0046024
```

2.13.5 Answer

The Cochran-Mantel-Haenszel test shows sufficient evidence of a treatment effect on survival rates (Q=8.03,p=0.0046).

> **Note**
>
> If the withdrawals were grouped with no recurrences, the results will be identical to the log-rank test in the previous example

2.14 Exercise

A study about mild traumatic brain injuries (mTBI) investigated the persistence of headaches for those who presented with mental health related sequelae and those who did not. Use the Mantel-Cox method to test if there is a difference in survival rates between the the two profiles.

2.14.1 SAS Data Set

```
data mTBI_mc;
input time $ profile $ status $ count @@;
datalines;
0-6 NoMH recur 17 0-6 NoMH not 35
0-6 MH recur 15 0-6 MH not 19
6-12 NoMH recur 9 6-12 NoMH not 20
6-12 MH recur 7 6-12 MH not 10
;
```

2.14.2 R Data Set

```
mTBI_mc <- data.frame(
  time    = c("0-6", "0-6", "0-6", "0-6", "6-12", "6-12", "6-12", "6-12"),
  profile = c("NoMH", "NoMH", "MH", "MH", "NoMH", "NoMH", "MH", "MH"),
  status  = c("recur", "not", "recur", "not", "recur", "not", "recur", "not"),
  count   = c(17, 35, 15, 19, 9, 20, 7, 10)
)
```

2.14.3 SAS Code

```
proc freq data=mTBI_mc order=data;
weight count;
tables time*profile*status / cmh;
run;
```

2.14.4 R Code

```
library(vcdExtra)
mtbi_cc <- xtabs(count~profile+status+time,data=mTBI_mc)
mtbi_cc
```

```
, , time = 0-6

       status
profile not recur
   MH    19    15
   NoMH  35    17

, , time = 6-12

       status
profile not recur
   MH    10     7
   NoMH  20     9
```

```
CMHtest(mtbi_cc,
        strata="time",
        overall=T)$ALL
```

```
Cochran-Mantel-Haenszel Statistics for profile by status
    Overall tests, controlling for all strata

                AltHypothesis  Chisq Df    Prob
cor        Nonzero correlation 1.6072  1 0.20489
rmeans  Row mean scores differ 1.6072  1 0.20489
cmeans  Col mean scores differ 1.6072  1 0.20489
general    General association 1.6072  1 0.20489
```

2.14.5 Answer

There is insufficient evidence of a difference between the survival rates of the two groups based on the Mantel-Cox test ($Q = 1.6072$, p=0.20).

2.15 Piecewise Exponential Models

Grouped survival data can be analyzed by providing a description of the pattern of event rates.

> **Note**
>
> The piecewise exponential model assumes that withdrawals are uniformly distributed during the time intervals in which they occur and unrelated to treatment failures. The within-interval probabilities of treatment failures are also assumed to be small and have independent exponential distributions.

2.16 Piecewise Exponential Likelihood

The piecewise exponential likelihood can be written as:

$$\Phi_{PE} = \prod_i \prod_k \lambda_{ik}^{n_i 1k} [\exp(-\lambda_{ik} N_{ik})]$$

where n_{i1k} is the number of failures of group i during the interval k, N_{ik} is the person-months of exposure, and λ_{ik} is the hazard parameter.

> **Important**
>
> The piecewise exponential model assumes independent exponential distributions with hazard parameters λ_{ik} for the respective time periods.

2.17 Piecewise Exponential Likelihood

N_{ik} is computed as

$$N_{ik} = a_k(n_{i0k} + 0.5n_{i1k} + 0.5n_{i2k})$$

where a_k is the length of the interval k. The conditional Poisson distribution of the number of failures/deaths conditioned on the exposure level also follows a Poisson likelihood.

> **Note**
>
> Since the likelihoods are proportional, whatever maximizes the conditional likelihood also maximizes the piecewise exponential likelihood.

2.18 Hazard Parameter

We can specify a model for the hazard parameter λ_{ik} such that

$$\log \lambda_{ik} = \alpha + \eta_k + \beta_1 X_{1i} + \beta_2 X_{2i} + ...$$

> **Note**
>
> This is similar to the proportional hazards model.

2.19 `R` Implementation

Both `LOGISTIC` and `GENMOD` can implement the piecewise exponential model analyses.

> **Important**
>
> The `GENMOD` procedure uses the Poisson likelihood to estimate failure rates, while `LOGISTIC` uses the response `failure/trials`. When using `LOGISTIC`, you need to rescale the exposure factor by a large number (at least 10,000 depending on the exposure variable) to scale the exposure from aggregate to individual.

2.20 `R` Implementation

The `glm()` function can implement the piecewise exponential model analyses. The Poisson (family="Poisson") is used to approximate the incidence rates, while the offset is commonly taken as the logarithm of the observation time.

2.21 Example

The data includes information pertaining to the experience of patients undergoing treatment for duodenal ulcers. types of surgeries was randomly assigned: vagotomy and drainage or antrectomy (VDA), or vagotomy and hemigastrectomy (VH). The patients were evaluated

at 6 months, 24 months, and 60 months. Use the piecewise exponential model to test for a treatment effect and if there is a difference in failure rates in time.

2.21.1 SAS Data Set

```
data vda;
input treatment $ time $ failure months;
nmonths=log(months);
datalines;
vda _0-6 23 3894
vda 7-24 32 10872
vda 25-60 45 18720
vh _0-6 9 2016
vh 7-24 5 5724
vh 25-60 10 10440
;
```

2.21.2 R Data Set

```
vda <- data.frame(
  treatment = c("vda", "vda", "vda", "vh", "vh", "vh"),
  time      = factor(c("0-6", "7-24", "25-60", "0-6", "7-24", "25-60"), levels=c("0-6", "7-24
  failure   = c(23, 32, 45, 9, 5, 10),
  months    = c(3894, 10872, 18720, 2016, 5724, 10440)
)
```

2.21.3 GENMOD

```
proc genmod data=vda;
class treatment time;
model failure = time treatment
/ dist=poisson link=log offset=nmonths;
estimate "failure rate 25-60 VH" int 1 time 1 0 0 treatment 0 1/exp;
estimate "failure rate 25-60 VDA" int 1 time 1 0 0 treatment 1 0/exp;
run;
```

2.21.4 R Code

```
mod1 <- glm(failure~time + treatment,
            data=vda,
            family="poisson",
            offset = log(months))
summary(mod1)
```

```
Call:
glm(formula = failure ~ time + treatment, family = "poisson",
    data = vda, offset = log(months))

Coefficients:
            Estimate Std. Error z value Pr(>|z|)
(Intercept)  -5.0093     0.1818 -27.546  < 2e-16 ***
time7-24     -0.8847     0.2414  -3.665 0.000247 ***
time25-60    -1.0429     0.2223  -4.691 2.72e-06 ***
treatmentvh  -0.8071     0.2273  -3.551 0.000384 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for poisson family taken to be 1)

    Null deviance: 37.3229  on 5  degrees of freedom
Residual deviance:  2.5529  on 2  degrees of freedom
AIC: 38.182

Number of Fisher Scoring iterations: 4
```

```
library(emmeans)
```

```
Welcome to emmeans.
Caution: You lose important information if you filter this package's results.
See '? untidy'
```

```
emmeans(mod1,~treatment*time,offset = 0,type="response")
```

```
 treatment time     rate       SE  df asymp.LCL asymp.UCL
 vda       0-6   0.00668 0.001210 Inf  0.004674   0.00953
```

```
 vh        0-6   0.00298 0.000761 Inf  0.001805   0.00492
 vda       7-24  0.00276 0.000468 Inf  0.001975   0.00385
 vh        7-24  0.00123 0.000303 Inf  0.000758   0.00199
 vda       25-60 0.00235 0.000335 Inf  0.001780   0.00311
 vh        25-60 0.00105 0.000238 Inf  0.000673   0.00164

Confidence level used: 0.95
Intervals are back-transformed from the log scale
```

2.21.5 Results

From the `GENMOD` results, the failure rates for the VDA treatment decreases across time. (β = -0.88,-1.04; p = <0.0001, 0.0002) The failure rate is higher for VDA compared to VH (β=0.8071, p=0.004)

The failure rates can be estimated using the `estimate` statement.The failure rate of VDA at 25-60 months is 0.0024 (0.0018,0.0031), while VH has a failure rate of 0.0010 (0.0007,0.0016).