

EAB 770 - Chapter 9

Miguel Fudolig

Table of contents

1	Logistic Regression - Polytomous Response	3
1.1	Outline	3
2	Ordinal Response Variables	3
2.1	Polytomous Variables?	3
2.2	Polytomous variables	3
2.3	Cumulative probabilities	4
2.4	Cumulative logits	4
2.5	Cumulative logits	4
2.6	Proportional Odds Assumption	5
2.7	Proportional Odds	5
2.8	Proportional Odds	5
2.9	Solving for Individual Probabilities	6
2.10	Proportional Odds Model in SAS	6
2.11	Proportional Odds Model in R	6
2.12	Example	7
2.12.1	Questions	7
2.12.2	Data	8
2.13	Example	9
2.13.1	SAS Data Set	9
2.13.2	SAS Code	10
2.13.3	R Code	10
2.13.4	Answer	15
2.14	Exercise	15
2.14.1	Questions	15
2.14.2	Data Set	16
2.15	Exercise	17
2.15.1	SAS Code	17
2.15.2	R Code	17

2.15.3	Answer	22
2.16	Partial Proportional Odds Model	23
2.17	Partial Proportional Odds Model in SAS	23
2.18	Partial Proportional Odds Model in R	24
2.19	Example	24
2.19.1	Questions	24
2.19.2	Data	24
2.19.3	R Data	25
2.20	Example	25
2.20.1	SAS Code	25
2.20.2	R Code	26
2.20.3	Answer	29
2.21	Exercise	29
2.22	Exercise	29
2.22.1	SAS Code	29
2.22.2	R Code	30
2.22.3	Answer	30
3	Nominal Response Variables	30
3.1	Generalized Logits Model	30
3.2	Generalized Logits Model	31
3.3	Generalized Logits Model: SAS	31
3.4	Generalized Logits Model: R	32
3.5	Example	32
3.5.1	Question	32
3.5.2	SAS Data Set	32
3.5.3	R Data Set	33
3.6	Example	33
3.6.1	SAS Code	33
3.6.2	R Code	33
3.6.3	Answer	35
3.7	Exercise	36
3.7.1	Question	36
3.7.2	Data	36
3.7.3	R Data	37
3.8	Exercise	37
3.8.1	SAS Code	37
3.8.2	R Code	37
3.8.3	Answer	39
3.9	Exact Generalized Logits Test	39
3.10	Example	40
3.10.1	Data Set	40

3.11 Example	40
3.11.1 SAS Code	40
3.11.2 Answer	41

1 Logistic Regression - Polytomous Response

1.1 Outline

- Ordinal Response
 - Proportional Odds Model
 - Multiple Qualitative Explanatory Variables
 - Partial Proportional Odds Model
- Nominal Response
 - Generalized Logits Model
 - Exact Methods: Generalized Logits Model

2 Ordinal Response Variables

2.1 Polytomous Variables?

Recall Chapter 8: dichotomous response variables can be analyzed using logistic regression.

Note

What if the response variable has more than two levels?

2.2 Polytomous variables

- Polytomous variables are usually described by the [multinomial distribution](#)

Tip

Unlike the binomial distribution where we only need one probability of success, more than one probability of success is needed to be defined in multinomial distributions.

2.3 Cumulative probabilities

Consider π_{hij} as the probability of success of the h^{th} strata and i^{th} explanatory variables and the j^{th} level of the response variable. We can define θ_{hij} such that:

$$\theta_{hiJ} = \sum_{j=1}^J \pi_{hij}$$

i Note

If $J = 1$, then $\theta_{hi1} = \pi_{hi1}$. If $J = 2$, then $\theta_{hi2} = \pi_{hi1} + \pi_{hi2}$. θ_{hiJ} is the *cumulative probability* of the response levels from $j = 1$ to $j = J$.

i Note

If there are $J + 1$ levels, then $\pi_{J+1} = 1 - \theta_{hiJ}$.

2.4 Cumulative logits

Cumulative logits can be defined based on the cumulative probabilities θ_{hij} . For a specific level k out of K levels,

$$\text{logit}(\theta_{hik}) = \log \left(\frac{\theta_{hik}}{1 - \theta_{hik}} \right) = \log \left(\frac{\sum_{j=1}^k \pi_{hij}}{1 - \sum_{j=1}^k \pi_{hij}} \right) = \log \left(\frac{\sum_{j=1}^k \pi_{hij}}{\sum_{j=k+1}^K \pi_{hij}} \right)$$

2.5 Cumulative logits

Suppose we define a linear predictor η_{hij} such that

$$\eta_{hij} = \text{logit}(\theta_{hij}) = \alpha_j + \mathbf{x}'_{hi} \boldsymbol{\beta}_k$$

where $\mathbf{x}'_{hi} \boldsymbol{\beta}_k$ represents the matrix product of the explanatory/strata variables x_{hi} and the coefficients for each response level β_k .

2.6 Proportional Odds Assumption

i Proportional Odds

The proportional odds assumption applies the following condition for a response variable with K total variable levels:

$$\mathbf{1} = \mathbf{2} = \mathbf{3} = \dots = \mathbf{K} =$$

Key assumption: the effect of the explanatory variables should not be disproportionate across all levels.

2.7 Proportional Odds

A consequence of the proportional odds assumption is that the difference between the levels of the explanatory variables are the same across different response levels.

i Example

Consider comparing the m^{th} and n^{th} level of the explanatory variable in the h^{th} strata for the first and second level of the response variable.

$$\eta_{hm1} - \eta_{hn1} = (\mathbf{x}'_{hm} - \mathbf{x}'_{hn})_1 = (\mathbf{x}'_{hm} - \mathbf{x}'_{hn})$$

$$\eta_{hm2} - \eta_{hn2} = (\mathbf{x}'_{hm} - \mathbf{x}'_{hn})_2 = (\mathbf{x}'_{hm} - \mathbf{x}'_{hn})$$

These are the same across all levels of the response variable. The difference between η correspond to the log odds, hence the odds ratios between explanatory variables are the same across all response levels.

2.8 Proportional Odds

Another consequence of the proportional odds assumption is that the difference between the response levels does not depend on the strata or explanatory variables.

i Example

Consider comparing the first and second level of the response variable.

$$\eta_{hm1} - \eta_{hm2} = \alpha_1 - \alpha_2 + (\mathbf{x}'_{hm} - \mathbf{x}'_{hm})\beta = \alpha_1 - \alpha_2$$

i Note

The proportional odds assumption can be tested using the Brant test. The null hypothesis of the test is that the proportional odds assumption holds. For the proportional odds assumption to hold, we must fail to reject the null hypothesis.

2.9 Solving for Individual Probabilities

Individual probabilities can be calculated by taking the differences between cumulative logits.

$$\pi_1 = \theta_1; \pi_2 = \theta_2 - \theta_1; \pi_k = \theta_k - \theta_{k-1}$$

2.10 Proportional Odds Model in SAS

- We can use the LOGISTIC, GENMOD, and GLIMMIX procedures to fit the proportional odds model.
- Dealing with ordinal variables means we need to specify order of the data.
- We can use the ESTIMATE and CONTRAST statements to determine and test the odds ratios between response levels, explanatory variable levels, and strata.
 - ESTIMATE: EXP option to get the estimated odds ratio, ILINK option to get the estimated cumulative probability θ .
 - CL option is used to get the confidence limits for odds ratio or estimated probabilities.

2.11 Proportional Odds Model in R

The `polr()` function in the MASS package in R performs the proportional odds model in R. Marginal/conditional means can still be calculated using `emmeans()` from the `emmeans` package.

i Note

The `poTest()` function in `car` tests whether the proportional odds assumption holds.

i Note

Because the `polr()` function uses a slightly different link between the cumulative logits, the interpretation for these numbers will be different from that of SAS. Calculating odds ratios will provide the odds ratio of recording higher level of outcomes between the two predictor levels.

2.12 Example

Male and female subjects received an active or placebo treatment for their arthritis pain, and the subsequent extent of improvement was recorded as marked, some, or none.

2.12.1 Questions

- Write the correct form of the linear predictor.
- Use the proportional odds model to determine whether there is an effect of sex and treatment.
- List the odds ratio estimates, 95% confidence intervals, and corresponding interpretations.
- Estimate the probability of marked improvement for females in the active group.

2.12.2 Data

Table 9.1 Arthritis

Sex	Treatment	Impr
		Marked
Female	Active	16
Female	Placebo	6
Male	Active	5
Male	Placebo	1

R Data Set

```
library(tidyverse)
```

```
-- Attaching core tidyverse packages ----- tidyverse 2.0.0 --
v dplyr      1.2.1      v readr      2.2.0
v forcats    1.0.1      v stringr    1.6.0
v ggplot2    4.0.3      v tibble     3.3.1
v lubridate  1.9.5      v tidyr      1.3.2
v purrr      1.2.2
-- Conflicts ----- tidyverse_conflicts() --
x dplyr::filter() masks stats::filter()
```



```
x dplyr::lag()      masks stats::lag()
i Use the conflicted package (<http://conflicted.r-lib.org/>) to force all conflicts to become
```

```
arthritis <- read.csv("arthritis_9.csv", stringsAsFactors = T)
glimpse(arthritis)
```

Rows: 84

Columns: 3

\$ Treatment <fct> Active, Active, Active, Active, Active, Active, Active, Ac~

\$ Sex <fct> Female, Female, Female, Female, Female, Female, Female, Fe~

\$ Improvement <fct> Marked, Marked, Marked, Marked, Marked, Marked, Marked, Ma~

It is important to check if the factor levels follow the order we want.

```
levels(arthritis$Improvement)
```

```
[1] "Marked" "None"   "Some"
```

```
levels(arthritis$Treatment)
```

```
[1] "Active" "Placebo"
```

```
levels(arthritis$Sex)
```

```
[1] "Female" "Male"
```

The code below fixes this.

```
arthritis <- arthritis |>
  mutate(Improvement = factor(Improvement, levels = c("None", "Some", "Marked"), ordered=T),
         Treatment = factor(Treatment, levels=c("Placebo", "Active")))
```

2.13 Example

Male and female subjects received an active or placebo treatment for their arthritis pain, and the subsequent extent of improvement was recorded as marked, some, or none.

2.13.1 SAS Data Set

```
data arthritis;
input sex $ treatment $ improve $ count @@;
datalines;
female active marked 16 female active some 5 female active none 6
female placebo marked 6 female placebo some 7 female placebo none 19
male active marked 5 male active some 2 male active none 7
male placebo marked 1 male placebo some 0 male placebo none 10
;
```

2.13.2 SAS Code

```
proc logistic data=arthritis order=data plots=effect(polybar x=treatment*sex); ①
freq count;
class treatment sex / param=ref;
model improve = sex treatment sex*treatment / selection=forward include=2 scale=none aggregate=none;
estimate "odds between marked vs. some+no improvement for female in placebo group" intercept 1 sex 1;
estimate "odds between marked+some vs. no improvement for female in placebo group" intercept 1 sex 1;
estimate "probability of marked improvement for female in the placebo group" intercept 1 sex 1;
estimate "probability of marked improvement for female in the active trt group" intercept 1 sex 1;
run;
```

- ① The polybar plot creates stacked bar plots showing probability and cumulative probability plots for each level dictated by x.
- ② `selection=forward include=2` tests whether the interaction is needed to be included in the model.
- ③ The `ilink` option calculates the cumulative probability, while `exp` calculates odds ratios. The reason it returns the probability for marked is that the marked level was inputted as the lowest level.

2.13.3 R Code

```
library(MASS)
```

Attaching package: 'MASS'

The following object is masked from 'package:dplyr':

```
select
```

```
library(car)
```

Loading required package: carData

Attaching package: 'car'

The following object is masked from 'package:dplyr':

recode

The following object is masked from 'package:purrr':

some

```
options(contrasts = c("contr.treatment", "contr.poly"))
mod_po <- polr(Improvement~Treatment+Sex,data=arthritis,Hess = TRUE)
summary(mod_po)
```

Call:

```
polr(formula = Improvement ~ Treatment + Sex, data = arthritis,
      Hess = TRUE)
```

Coefficients:

	Value	Std. Error	t value
TreatmentActive	1.797	0.4718	3.809
SexMale	-1.319	0.5381	-2.451

Intercepts:

	Value	Std. Error	t value
None Some	0.4941	0.3369	1.4665
Some Marked	1.3485	0.3667	3.6768

Residual Deviance: 150.0294

AIC: 158.0294

```
exp(coef(mod_po))
```

TreatmentActive	SexMale
6.0330950	0.2674786

```
poTest(mod_po)
```

Tests for Proportional Odds

```
polr(formula = Improvement ~ Treatment + Sex, data = arthritis,  
      Hess = TRUE)
```

	b[polr]	b[>None]	b[>Some]	Chisquare	df	Pr(>Chisq)
Overall				1.83	2	0.40
TreatmentActive	1.79726	1.78168	1.81602	0.01	1	0.94
SexMale	-1.31872	-1.46865	-0.93005	1.58	1	0.21

```
Anova(mod_po)
```

Analysis of Deviance Table (Type II tests)

Response: Improvement

	LR	Chisq	Df	Pr(>Chisq)
Treatment	16.0812	1	6.068e-05	***
Sex	6.5766	1	0.01033	*

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
library(emmeans)
```

Welcome to emmeans.

Caution: You lose important information if you filter this package's results.

See '? untidy'

```
emm_cum <- emmeans(mod_po, ~Treatment*Sex | cut, mode = "cum.prob")  
emm_cum
```

cut = None|Some:

Treatment	Sex	cumprob	SE	df	asyp.LCL	asyp.UCL
Placebo	Female	0.621	0.0793	Inf	0.4657	0.776
Active	Female	0.214	0.0658	Inf	0.0847	0.343
Placebo	Male	0.860	0.0682	Inf	0.7260	0.993
Active	Male	0.504	0.1180	Inf	0.2725	0.735

cut = Some|Marked:

Treatment	Sex	cumprob	SE	df	asympt.LCL	asympt.UCL
Placebo	Female	0.794	0.0600	Inf	0.6763	0.912
Active	Female	0.390	0.0870	Inf	0.2192	0.560
Placebo	Male	0.935	0.0368	Inf	0.8629	1.007
Active	Male	0.705	0.0999	Inf	0.5090	0.900

Confidence level used: 0.95

```
emm_prob <- emmeans(mod_po, ~Treatment*Sex | Improvement, mode="prob")
emm_prob
```

Improvement = None:

Treatment	Sex	prob	SE	df	asympt.LCL	asympt.UCL
Placebo	Female	0.6211	0.0793	Inf	0.46567	0.776
Active	Female	0.2136	0.0658	Inf	0.08466	0.343
Placebo	Male	0.8597	0.0682	Inf	0.72603	0.993
Active	Male	0.5039	0.1180	Inf	0.27255	0.735

Improvement = Some:


Treatment	Sex	prob	SE	df	asympt.LCL	asympt.UCL
Placebo	Female	0.1728	0.0457	Inf	0.08315	0.262
Active	Female	0.1760	0.0474	Inf	0.08320	0.269
Placebo	Male	0.0754	0.0361	Inf	0.00454	0.146
Active	Male	0.2008	0.0519	Inf	0.09904	0.303

Improvement = Marked:

Treatment	Sex	prob	SE	df	asympt.LCL	asympt.UCL
Placebo	Female	0.2061	0.0600	Inf	0.08850	0.324
Active	Female	0.6104	0.0870	Inf	0.43990	0.781
Placebo	Male	0.0649	0.0368	Inf	-0.00724	0.137
Active	Male	0.2953	0.0999	Inf	0.09954	0.491

Confidence level used: 0.95

```
emm_prob |> plot()
```



To calculate the odds ratio, we can use `contrast()` but we have to manually exponentiate the confidence limits.

```
emm_treatment <- emmeans(mod_po, ~Treatment)
contrast(emm_treatment, method="revpairwise", infer=T) |>
  as.data.frame() |>
  mutate(OR = exp(estimate),
         OR.LCL = exp(asymp.LCL),
         OR.UCL = exp(asymp.UCL))
```

	contrast	estimate	SE	df	asymp.LCL	asymp.UCL	z.ratio
1	Active - Placebo	1.79726	0.4717993	Inf	0.8725504	2.72197	3.809374
	p.value	OR	OR.LCL	OR.UCL			
1		0.0001393191	6.033095	2.393006	15.21025		

```
emm_sex <- emmeans(mod_po, ~Sex)
contrast(emm_sex, method="pairwise", infer=T) |>
  as.data.frame() |>
  mutate(OR = exp(estimate),
```

```
OR.LCL = exp(asymp.LCL),
OR.UCL = exp(asymp.UCL))
```

	contrast	estimate	SE	df	asymp.LCL	asymp.UCL	z.ratio	p.value
1	Female - Male	1.318716	0.5380997	Inf	0.2640599	2.373372	2.450691	0.01425824
	OR	OR.LCL	OR.UCL					
1		3.738617	1.302206		10.73352			

2.13.4 Answer

The linear predictor can be written as

$$\eta_k = \alpha_k + \beta_f x_f + \beta_a x_a$$

where $x_f = 1$ when the participant identifies as female, $x_a = 1$ when the participant is in the active treatment group. Using the residual chi-square test, the interaction term is not significant ($\chi^2 = 0.28, p = 0.60$).

Using the proportional odds model, we determine that there is a significant effect of sex ($\chi^2 = 6.2, p = 0.01$) and treatment ($\chi^2 = 14.4, p = 0.001$).

Females have 3.74 (95% CI: 1.33, 10.55) times the odds of marked improvement compared to males.

The odds of improvement after undergoing active treatment is 6.03 (95% CI: 2.39, 15.2) times that of the placebo group.

2.14 Exercise

Researchers collected information on subjects' exposure to general air pollution, exposure to pollution in their jobs, and whether they smoked. The response measured was chronic respiratory disease status. Subjects were assigned to one of four possible categories.

2.14.1 Questions

- Use the proportional odds main effects model to test the effect of air pollution, job exposure, and smoking status in the response.
- Estimate the odds ratio between Response I+II+III vs. IV for current vs. ex-smokers.
- Estimate the probability of developing a response level III for an ex-smoker in a high air pollution environment with job exposure.

2.14.2 Data Set

```
data respire;
input air $ exposure $ smoking $ level $ count @@;
datalines;
low no non 1 158 low no non 2 9
low no ex 1 167 low no ex 2 19
low no cur 1 307 low no cur 2 102
low yes non 1 26 low yes non 2 5
low yes ex 1 38 low yes ex 2 12
low yes cur 1 94 low yes cur 2 48
high no non 1 94 high no non 2 7
high no ex 1 67 high no ex 2 8
high no cur 1 184 high no cur 2 65
high yes non 1 32 high yes non 2 3
high yes ex 1 39 high yes ex 2 11
high yes cur 1 77 high yes cur 2 48
low no non 3 5 low no non 4 0
low no ex 3 5 low no ex 4 3
low no cur 3 83 low no cur 4 68
low yes non 3 5 low yes non 4 1
low yes ex 3 4 low yes ex 4 4
low yes cur 3 46 low yes cur 4 60
high no non 3 5 high no non 4 1
high no ex 3 4 high no ex 4 3
high no cur 3 33 high no cur 4 36
high yes non 3 6 high yes non 4 1
high yes ex 3 4 high yes ex 4 2
high yes cur 3 39 high yes cur 4 51
;
```

A csv file is uploaded on Canvas as `air_pollution.csv`.

```
library(tidyverse)
air_pollution <- read.csv("air_pollution.csv", stringsAsFactors = TRUE)
air_pollution <- air_pollution |>
  mutate(Level = factor(Level, ordered=T))

glimpse(air_pollution)
```

Rows: 2,089

Columns: 4

```
$ Air      <fct> high, high, high, high, high, high, high, high, high, high, h~
$ Exposure <fct> yes, yes, yes, yes, yes, yes, yes, yes, yes, yes, yes, yes, y~
$ Smoking  <fct> cur, cur, cur, cur, cur, cur, cur, cur, cur, cur, cur, cur, c~
$ Level    <ord> 4, 4, 4, 4, 4, 4, 4, 4, 4, 4, 4, 4, 4, 4, 4, 4, 4, 4, 4, 4, 4~
```

2.15 Exercise

Researchers collected information on subjects' exposure to general air pollution, exposure to pollution in their jobs, and whether they smoked. The response measured was chronic respiratory disease status. Subjects were assigned to one of four possible categories.

2.15.1 SAS Code

```
proc logistic data=respire order=data plots=effect(polybar x=air*exposure*smoking);;
freq count;
class air exposure smoking / param=reference;
model level = air exposure smoking air*exposure exposure*smoking air*smoking air*exposure*sm
estimate "odds between I+II+III vs. IV for current vs. ex smokers" smoking 1 -1 /exp cl cate
estimate "probability of at most III level for ex smoker, high air pollution, job exposure" :
estimate "probability of at most II level for ex smoker, high air pollution, job exposure" i
run;
```

2.15.2 R Code

```
library(MASS)
library(car)
options(contrasts = c("contr.treatment", "contr.poly"))
mod_po <- polr(Level~Air + Exposure + Smoking, data=air_pollution,Hess=TRUE)

summary(mod_po)
```

Call:

```
polr(formula = Level ~ Air + Exposure + Smoking, data = air_pollution,
      Hess = TRUE)
```

Coefficients:

	Value	Std. Error	t value
Airlow	0.03929	0.09384	0.4187

Exposureyes	0.86476	0.09576	9.0302
Smokingex	-1.45268	0.13725	-10.5842
Smokingnon	-1.85271	0.16481	-11.2416

Intercepts:

	Value	Std. Error	t value
1 2	0.2750	0.0880	3.1256
2 3	1.1562	0.0924	12.5133
3 4	2.0804	0.1044	19.9318

Residual Deviance: 4178.511
AIC: 4192.511

Anova(mod_po)

Analysis of Deviance Table (Type II tests)

Response: Level

	LR	Chisq	Df	Pr(>Chisq)
Air	0.175	1		0.6753
Exposure	81.308	1		<2e-16 ***
Smoking	255.541	2		<2e-16 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

poTest(mod_po)

Tests for Proportional Odds

```
polr(formula = Level ~ Air + Exposure + Smoking, data = air_pollution,
      Hess = TRUE)
```

	b[polr]	b[>1]	b[>2]	b[>3]	Chisquare	df	Pr(>Chisq)
Overall					10.66	8	0.222
Airlow	0.03929	0.00592	0.10350	0.02952	1.39	2	0.499
Exposureyes	0.86476	0.84954	0.85999	0.87715	0.04	2	0.981
Smokingex	-1.45268	-1.39976	-1.71927	-1.76946	3.61	2	0.165
Smokingnon	-1.85271	-1.83705	-1.75516	-3.02210	5.53	2	0.063 .

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
library(emmeans)
emm_cum <- emmeans(mod_po, ~Air+Exposure+Smoking|cut, mode="cum.prob")
emm_cum
```

cut = 1|2:

Air	Exposure	Smoking	cumprob	SE	df	asympt.LCL	asympt.UCL
high	no	cur	0.568	0.02160	Inf	0.526	0.611
low	no	cur	0.559	0.01790	Inf	0.524	0.594
high	yes	cur	0.357	0.02200	Inf	0.313	0.400
low	yes	cur	0.348	0.02140	Inf	0.306	0.390
high	no	ex	0.849	0.01880	Inf	0.812	0.886
low	no	ex	0.844	0.01780	Inf	0.809	0.879
high	yes	ex	0.703	0.03090	Inf	0.643	0.764
low	yes	ex	0.695	0.03060	Inf	0.635	0.755
high	no	non	0.894	0.01610	Inf	0.862	0.925
low	no	non	0.890	0.01600	Inf	0.858	0.921
high	yes	non	0.780	0.03000	Inf	0.721	0.838
low	yes	non	0.773	0.03070	Inf	0.713	0.833

cut = 2|3:

Air	Exposure	Smoking	cumprob	SE	df	asympt.LCL	asympt.UCL
high	no	cur	0.761	0.01680	Inf	0.728	0.794
low	no	cur	0.753	0.01440	Inf	0.725	0.782
high	yes	cur	0.572	0.02350	Inf	0.526	0.618
low	yes	cur	0.563	0.02310	Inf	0.517	0.608
high	no	ex	0.931	0.00978	Inf	0.912	0.951
low	no	ex	0.929	0.00939	Inf	0.910	0.947
high	yes	ex	0.851	0.01920	Inf	0.814	0.889
low	yes	ex	0.846	0.01930	Inf	0.808	0.884
high	no	non	0.953	0.00784	Inf	0.938	0.968
low	no	non	0.951	0.00783	Inf	0.936	0.967
high	yes	non	0.895	0.01670	Inf	0.862	0.928
low	yes	non	0.891	0.01720	Inf	0.858	0.925

cut = 3|4:

Air	Exposure	Smoking	cumprob	SE	df	asympt.LCL	asympt.UCL
high	no	cur	0.889	0.01030	Inf	0.869	0.909
low	no	cur	0.885	0.00923	Inf	0.867	0.903
high	yes	cur	0.771	0.01830	Inf	0.735	0.807
low	yes	cur	0.764	0.01830	Inf	0.728	0.800
high	no	ex	0.972	0.00450	Inf	0.963	0.980
low	no	ex	0.971	0.00436	Inf	0.962	0.979

high	yes	ex	0.935	0.00965	Inf	0.916	0.954
low	yes	ex	0.933	0.00976	Inf	0.914	0.952
high	no	non	0.981	0.00346	Inf	0.974	0.988
low	no	non	0.980	0.00347	Inf	0.973	0.987
high	yes	non	0.956	0.00782	Inf	0.940	0.971
low	yes	non	0.954	0.00809	Inf	0.938	0.970

Confidence level used: 0.95

```
emm_prob <- emmeans(mod_po, ~Air+Exposure+Smoking|Level, mode="prob")
emm_prob
```

Level = 1:

Air	Exposure	Smoking	prob	SE	df	asympt.LCL	asympt.UCL
high	no	cur	0.5683	0.02160	Inf	0.5260	0.6106
low	no	cur	0.5587	0.01790	Inf	0.5236	0.5937
high	yes	cur	0.3567	0.02200	Inf	0.3135	0.3999
low	yes	cur	0.3477	0.02140	Inf	0.3057	0.3898
high	no	ex	0.8491	0.01880	Inf	0.8123	0.8860
low	no	ex	0.8440	0.01780	Inf	0.8091	0.8790
high	yes	ex	0.7033	0.03090	Inf	0.6428	0.7638
low	yes	ex	0.6950	0.03060	Inf	0.6350	0.7551
high	no	non	0.8936	0.01610	Inf	0.8620	0.9251
low	no	non	0.8898	0.01600	Inf	0.8585	0.9211
high	yes	non	0.7795	0.03000	Inf	0.7207	0.8383
low	yes	non	0.7727	0.03070	Inf	0.7125	0.8329

Level = 2:

Air	Exposure	Smoking	prob	SE	df	asympt.LCL	asympt.UCL
high	no	cur	0.1923	0.01060	Inf	0.1716	0.2130
low	no	cur	0.1948	0.01020	Inf	0.1749	0.2147
high	yes	cur	0.2157	0.01070	Inf	0.1947	0.2366
low	yes	cur	0.2150	0.01070	Inf	0.1940	0.2360
high	no	ex	0.0823	0.00987	Inf	0.0630	0.1017
low	no	ex	0.0849	0.00940	Inf	0.0664	0.1033
high	yes	ex	0.1479	0.01370	Inf	0.1210	0.1749
low	yes	ex	0.1512	0.01360	Inf	0.1245	0.1778
high	no	non	0.0594	0.00875	Inf	0.0423	0.0766
low	no	non	0.0614	0.00868	Inf	0.0444	0.0784
high	yes	non	0.1156	0.01450	Inf	0.0872	0.1439
low	yes	non	0.1187	0.01470	Inf	0.0899	0.1475

Level = 3:

Air	Exposure	Smoking	prob	SE	df	asympt.LCL	asympt.UCL
high	no	cur	0.1283	0.00992	Inf	0.1089	0.1478
low	no	cur	0.1316	0.00919	Inf	0.1136	0.1496
high	yes	cur	0.1989	0.01280	Inf	0.1738	0.2240
low	yes	cur	0.2016	0.01280	Inf	0.1764	0.2267
high	no	ex	0.0402	0.00579	Inf	0.0288	0.0515
low	no	ex	0.0416	0.00559	Inf	0.0307	0.0526
high	yes	ex	0.0839	0.01070	Inf	0.0629	0.1050
low	yes	ex	0.0865	0.01080	Inf	0.0654	0.1077
high	no	non	0.0278	0.00468	Inf	0.0186	0.0370
low	no	non	0.0289	0.00468	Inf	0.0197	0.0380
high	yes	non	0.0604	0.00957	Inf	0.0417	0.0792
low	yes	non	0.0625	0.00985	Inf	0.0432	0.0818

Level = 4:

Air	Exposure	Smoking	prob	SE	df	asympt.LCL	asympt.UCL
high	no	cur	0.1110	0.01030	Inf	0.0908	0.1312
low	no	cur	0.1150	0.00923	Inf	0.0969	0.1330
high	yes	cur	0.2287	0.01830	Inf	0.1928	0.2646
low	yes	cur	0.2357	0.01830	Inf	0.1998	0.2716
high	no	ex	0.0284	0.00450	Inf	0.0196	0.0372
low	no	ex	0.0295	0.00436	Inf	0.0209	0.0380
high	yes	ex	0.0649	0.00965	Inf	0.0459	0.0838
low	yes	ex	0.0673	0.00976	Inf	0.0482	0.0864
high	no	non	0.0192	0.00346	Inf	0.0124	0.0260
low	no	non	0.0200	0.00347	Inf	0.0132	0.0268
high	yes	non	0.0444	0.00782	Inf	0.0291	0.0598
low	yes	non	0.0461	0.00809	Inf	0.0303	0.0620

Confidence level used: 0.95

To calculate the odds ratio, we can use `contrast()` but we have to manually exponentiate the confidence limits.

```
emm_Air <- emmeans(mod_po, ~Air)
contrast(emm_Air, method="revpairwise", infer=T) |>
  as.data.frame() |>
  mutate(OR = exp(estimate),
         OR.LCL = exp(asympt.LCL),
         OR.UCL = exp(asympt.UCL))
```

contrast	estimate	SE	df	asympt.LCL	asympt.UCL	z.ratio	p.value
----------	----------	----	----	------------	------------	---------	---------

1	low - high	0.0392934	0.09384313	Inf	-0.1446358	0.2232226	0.4187136	0.6754255
		OR	OR.LCL	OR.UCL				
1		1.040076	0.8653374	1.250099				

```
emm_Ex <- emmeans(mod_po,~Exposure)
contrast(emm_Ex,method="revpairwise",infer=T) |>
  as.data.frame() |>
  mutate(OR = exp(estimate),
         OR.LCL = exp(asymp.LCL),
         OR.UCL = exp(asymp.UCL))
```

	contrast	estimate	SE	df	asymp.LCL	asymp.UCL	z.ratio	p.value
1	yes - no	0.8647616	0.09576341	Inf	0.6770688	1.052454	9.030189	1.713763e-19
		OR	OR.LCL	OR.UCL				
1		2.37444	1.9681	2.864674				

```
emm_Smoking <- emmeans(mod_po,~Smoking)
contrast(emm_Smoking,method="pairwise",infer=T,adjust="tukey") |>
  as.data.frame() |>
  mutate(OR = exp(estimate),
         OR.LCL = exp(asymp.LCL),
         OR.UCL = exp(asymp.UCL))
```

	contrast	estimate	SE	df	asymp.LCL	asymp.UCL	z.ratio	p.value	OR	OR.LCL	OR.UCL
1	cur - ex	1.4526828	0.1372496	Inf	1.13101079	1.7743549	10.584238				
2	cur - non	1.8527107	0.1648078	Inf	1.46645064	2.2389708	11.241647				
3	ex - non	0.4000279	0.2007087	Inf	-0.07037313	0.8704288	1.993077				
1								2.786660e-14	4.274567	3.098787	5.896476
2								5.329071e-15	6.377082	4.333826	9.383668
3								1.138789e-01	1.491866	0.932046	2.387935

2.15.3 Answer

There is no evidence of an air pollution effect ($\chi^2 = 0.18, p = 0.68$). There is evidence of a strong association between exposure ($\chi^2 = 82.1, p = < 0.001$) and smoking history ($\chi^2 = 209.9, p < 0.001$).

The odds of being assigned to Response Level III and lower for current smokers is 0.23 times the odds for ex-smokers. The odds of being assigned to Response Level 4 for current smokers is 4.27 times the odds for ex-smokers.

The probability of receiving a response level III and below for an ex smoker, high air pollution, and with job exposure is 0.9351. The probability of receiving a response level II and below for an ex smoker, high air pollution, and with job exposure is 0.8512. Hence, the estimated probability of receiving a response level III is 0.0839 .

2.16 Partial Proportional Odds Model

Sometimes, the $\beta_k = \beta$ assumption does not hold for all explanatory variables, especially if some levels disproportionately affect the odds of success.



Tip

This assumption can be tested using the score test for the proportional odds assumption provided by the LOGISTIC procedure.

To address this, we specify unequal slopes for the explanatory variables that violate the proportional odds assumption.

$$\text{logit}(\theta_{hik}) = \alpha_k + x'_{hi}\beta_k + x'_{h'i'}\gamma$$

where x'_{hi} includes the explanatory variables with unequal slopes β_k and $x'_{h'i'}$ includes the explanatory variables with equal slopes γ . This model is called the **partial proportional odds model**.

2.17 Partial Proportional Odds Model in SAS

- If the score test for the proportional odds assumption is significant at the 0.10 level, we might need to use a partial proportional odds model.
- We can specify the model to have unequal slopes using the UNEQUALSLOPES option in the MODEL statement. UNEQUALSLOPES=X specifies that only the explanatory variable X has unequal slopes
- Testing for unequal slopes can be done using the TEST statement
 - LABEL: TEST statements effectively labels the test.
 - If the test yields a significant result, there is sufficient evidence for unequal slopes for that explanatory variable.
 - If the test is not significant, we can assume the proportional odds model holds.

2.18 Partial Proportional Odds Model in R

If the score test for the proportional odds assumption led to the rejection of the null hypothesis at the 0.10 level, we can use a partial proportional odds model.

i Note

The `vglm` package in the `VGAM` package in R can be used to perform partial proportional odds model. Suppose we have two predictors, `x1` and `x2`, with `x2` not satisfying the proportional odds model.

```
mod_vglm <- vglm(outcome~x1 + x2,  
                 data= sample_data,  
                 family=cumulative(parallel=FALSE~x2))
```

It can also be used to estimate proportional odds models. However, `vglm()` does not work with `emmeans` as of September 2026.

2.19 Example

The table contains data from a study by a bicycling clothing manufacturer who wanted to assess their test glove. It was designed to combat the hand problems experienced by cyclists dealing with carpal tunnel syndrome. Cyclists wore either the company's standard gel glove or the new test glove for a week's worth of their standard rides. Then they reported whether they experienced major, moderate, or no relief from their usual symptoms (numbness and nerve pain) on the bike.

2.19.1 Questions

- Assume an unequal slopes, main effect model and test for the equality of slopes for glove and gender.
- Perform logistic regression and interpret the results.

2.19.2 Data

```
data wrist;  
input glove $ gender $ relief $ count @@;  
datalines;  
test female major 12 test female moderate 8 test female none 5  
test male major 8 test male moderate 14 test male none 15
```



```
gel female major 15 gel female moderate 5 gel female none 9
gel male major 8 gel male moderate 4 gel male none 20
;
```

2.19.3 R Data

```
library(tidyverse)
cycling <- expand.grid(Relief = c("Major","Moderate","None"),
                      Gender = c("Female","Male"),
                      Type = c("Test","Gel"))

cycling$count <- c(12,8,5,8,14,15,15,5,9,8,4,20)
cycling <- cycling |>
  mutate(Relief = ordered(Relief, levels = c("Major","Moderate","None"))) |>
  uncount(weights=count)
```

2.20 Example

The table contains data from a study by a bicycling clothing manufacturer who wanted to assess their test glove. It was designed to combat the hand problems experienced by cyclists dealing with carpal tunnel syndrome. Cyclists wore either the company's standard gel glove or the new test glove for a week's worth of their standard rides. Then they reported whether they experienced major, moderate, or no relief from their usual symptoms (numbness and nerve pain) on the bike.

2.20.1 SAS Code

```
proc logistic order=data data=wrist;
freq count;
class glove gender / param=reference order=data;
model relief= glove gender / scale=none aggregate unequalslopes; ①
pogender: test genderfemale_major=genderfemale_moderate; ②
poglove: test glovetest_major=glovetest_moderate; ③
run;

proc logistic order=data data=wrist;
freq count;
class glove gender / param=reference order=data;
```

```
model relief= glove gender / scale=none aggregate unequalslopes=glove; ④
run;
```

- ① `unequal slopes` tells SAS to provide unequal slopes for each explanatory level at all response levels.
- ② `pogender` is the label for testing whether the slopes should be the same for different response levels for the gender explanatory variable
- ③ `poglove` is the label for testing whether the slopes should be the same for different response levels for the glove explanatory variable
- ④ Only assumes unequal slopes for glove, but proportional odds assumption holds for gender.

2.20.2 R Code

```
library(MASS)
library(ordinal)
```

Warning: package 'ordinal' was built under R version 4.6.1

Attaching package: 'ordinal'

The following object is masked from 'package:dplyr':

slice

```
library(car)
mod_po <- polr(Relief~Gender+Type,data=cycling)
poTest(mod_po)
```

Tests for Proportional Odds

```
polr(formula = Relief ~ Gender + Type, data = cycling)
```

	b[polr]	b[>Major]	b[>Moderate]	Chisquare	df	Pr(>Chisq)
Overall				6.85	2	0.0326 *
GenderMale	1.1766	1.1873	1.1747	0.00	1	0.9724
TypeGel	0.3504	-0.1683	0.7811	6.75	1	0.0094 **

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

The proportional odds test shows that there is evidence of unequal slopes for type, but not for gender.

```
library(VGAM)
```

Warning: package 'VGAM' was built under R version 4.6.1

Loading required package: stats4

Loading required package: splines

Attaching package: 'VGAM'

The following objects are masked from 'package:ordinal':

dgumbel, dloggamma, pgumbel, plgamma, qgumbel, rgumbel, wine

```
library(car)
modv<- vglm(Relief~Gender+Type,
  family = cumulative(parallel=FALSE~ Type),
  data=cycling)

summary(modv)
```

Call:

```
vglm(formula = Relief ~ Gender + Type, family = cumulative(parallel = FALSE ~
  Type), data = cycling)
```

Coefficients:

	Estimate	Std. Error	z value	Pr(> z)
(Intercept):1	-0.0826	0.3379	-0.244	0.806893
(Intercept):2	1.5170	0.3677	4.125	3.7e-05 ***
GenderMale	-1.1819	0.3548	-3.331	0.000866 ***
TypeGel:1	0.1485	0.3926	0.378	0.705225
TypeGel:2	-0.7952	0.3894	-2.042	0.041167 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Names of linear predictors: logitlink(P[Y<=1]), logitlink(P[Y<=2])

Residual deviance: 247.0353 on 241 degrees of freedom

Log-likelihood: -123.5177 on 241 degrees of freedom

Number of Fisher scoring iterations: 4

Exponentiated coefficients:

GenderMale	TypeGel:1	TypeGel:2
0.3067019	1.1600929	0.4515106

```
Anova(modv)
```

Analysis of Deviance Table (Type II tests)

Response: Relief

	Df	Chisq	Pr(>Chisq)
Gender	1	11.0946	0.0008658 ***
Type	2	7.7644	0.0206052 *

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Let's calculate the probabilities and odds ratios.

```
# Odds Ratio
```

```
exp(coef(modv))
```

(Intercept):1	(Intercept):2	GenderMale	TypeGel:1	TypeGel:2
0.9207215	4.5584608	0.3067019	1.1600929	0.4515106

```
exp(confint(modv))
```

	2.5 %	97.5 %
(Intercept):1	0.4747861	1.7854949
(Intercept):2	2.2172362	9.3718316
GenderMale	0.1529985	0.6148169
TypeGel:1	0.5374483	2.5040837
TypeGel:2	0.2104667	0.9686180

```
# For reverse pairwise order: (note that the limits are reversed.)
exp(-coef(modv))
```

```
(Intercept):1 (Intercept):2  GenderMale  TypeGel:1  TypeGel:2
      1.0861047      0.2193723      3.2604949      0.8619999      2.2147872
```

```
exp(-confint(modv))
```

```

                2.5 %    97.5 %
(Intercept):1  2.1062117 0.5600688
(Intercept):2  0.4510119 0.1067027
GenderMale     6.5360121 1.6265005
TypeGel:1      1.8606441 0.3993477
TypeGel:2      4.7513445 1.0323988
```

2.20.3 Answer

Using a partial proportional odds model, there is a significant effect of the type of glove and gender. the odds of experiencing major relief for females is 3.26 (95% CI: 1.63, 6.54) times that of males. The odds of major relief when using the test glove is 1.16 (0.54, 2.50) times that of gel. Additionally, the odds of moderate relief when using the test glove is 2.22 (1.03, 4.75) times that of gel.

2.21 Exercise

Consider the arthritis example discussed earlier. Test the hypothesis of equal slopes for both sex and treatment.

2.22 Exercise

Consider the arthritis example discussed earlier. Test the hypothesis of equal slopes for both sex and treatment.

2.22.1 SAS Code

```
proc logistic data=arthritis order=data plots=effect(polybar x=treatment*sex);;
freq count;
class treatment sex / param=reference;
model improve = sex treatment / unequalslopes scale=none aggregate;
equal_sex: test sexfemale_marked=sexfemale_some;
equal_treatment: test treatmentactive_marked=treatmentactive_some;
run;
```

2.22.2 R Code

```
mod_po <- polr(Improvement~Sex+Treatment,data=arthritis)
poTest(mod_po)
```

Tests for Proportional Odds

```
polr(formula = Improvement ~ Sex + Treatment, data = arthritis)
```

	b[polr]	b[>None]	b[>Some]	Chisquare	df	Pr(>Chisq)
Overall				1.83	2	0.40
SexMale	-1.31872	-1.46865	-0.93005	1.58	1	0.21
TreatmentActive	1.79726	1.78168	1.81602	0.01	1	0.94

2.22.3 Answer

We suppose the slopes are unequal, then test whether there is evidence against the null hypothesis of equal slopes using `test`.

SAS: There is insufficient evidence to claim that the β for sex ($\chi^2 = 1.35, p = 0.25$) and treatment ($\chi^2 = 0.03, p = 0.86$) are different across the different outcome levels. R: There is insufficient evidence to claim that the β for sex ($\chi^2 = 1.58, p = 0.21$) and treatment ($\chi^2 = 0.01, p = 0.94$) are different across the different outcome levels.

3 Nominal Response Variables

3.1 Generalized Logits Model

When the response variables are nominal, the concept of cumulative probabilities is less applicable. The **generalized logits model** is used for nominal polytomous response variables.

Generalized Logit Model

The logit function is the logarithm of the ratio of the probability of the level of interest, π_j and the *reference* category π_r .

$$\text{logit}\theta_j = \log\left(\frac{\pi_j}{\pi_r}\right) = \alpha_j + \sum_g \beta_{jg}x_g$$

where g comprises of ALL explanatory variables

Reference level

For the reference level, $\alpha_r = 0, \beta_{rg} = 0$.

3.2 Generalized Logits Model

The resulting expression for π_j can be written as:

$$\pi_j = \frac{\exp[\alpha_j + \sum_g \beta_{jg}x_g]}{\sum_{k=1}^r \exp[\alpha_k + \sum_g \beta_{kg}x_g]}$$

with the constraint $\alpha_r = 0, \beta_{rg} = 0$.

3.3 Generalized Logits Model: SAS

Implementing generalized logit models in SAS

The LOGISTIC procedure will model the logit models for all levels simultaneously. If there are r levels of the response variable, there will be $(r - 1)$ degrees of freedom for a single logit model.

Link function

SAS does not know whether your response variable is ordinal or nominal. The generalized logits model can be specified using `link=glogit` in the `model` statement.

Syntax

- Unless specified, the last variable level based on the specified order in the `class` statement is set as the reference level.
- Syntax is the same as the PO model, but the assumptions are different.

3.4 Generalized Logits Model: R

The `vglm()` function in the `VGAM` package performs the generalized logits model using the `multinomial()` function to define the family of models.

```
modv <- vglm(Outcome~x1 + x2,  
             data=sample_data,  
             family=multinomial(parallel=FALSE, refLevel=1)) #reflevel=1 just specifies the 1
```

3.5 Example

Schoolchildren in experimental learning settings were surveyed to determine which learning style they preferred. Investigators were interested in whether response was associated with school and their program, which was either a standard school day or also included after-hours care.

3.5.1 Question

Use logistic regression to test for program and school effects and calculate the relevant odds ratios.

3.5.2 SAS Data Set

```
data school;  
input school program $ style $ count @@;  
datalines;  
1 regular self 10 1 regular team 17 1 regular class 26  
1 after self 5 1 after team 12 1 after class 50  
2 regular self 21 2 regular team 17 2 regular class 26  
2 after self 16 2 after team 12 2 after class 36  
3 regular self 15 3 regular team 15 3 regular class 16  
3 after self 12 3 after team 12 3 after class 20
```



```
;
proc freq data=school order=data;
weight count;
tables school*program*style/ nocol norow nopct;
run;
```

3.5.3 R Data Set

```
library(tidyverse)
school <- expand.grid(Program = c("Regular","After"),
                     Style = c("Self","Team","Class"),
                     School = c("1","2","3"))

school$Count <- c(10,5,17,12,26,50,21,16,17,12,26,36,15,12,15,12,16,20)

school <- school |>
  uncount(weights = Count)
```

3.6 Example

Schoolchildren in experimental learning settings were surveyed to determine which learning style they preferred. Investigators were interested in whether response was associated with school and their program, which was either a standard school day or also included after-hours care. Use logistic regression to test for program and school effects and calculate the relevant odds ratios.

3.6.1 SAS Code

```
proc logistic data=school order=data;
freq count;
class school (ref=first) program (ref=last) / param=ref;
model style= program school school*program / link=glogit selection=forward include=2 scale=n;
run;
```

3.6.2 R Code

```

modv <- vglm(Style~School + Program,
             data=school,
             family=multinomial(parallel=FALSE, refLevel=3))

summary(modv)

```

Call:

```

vglm(formula = Style ~ School + Program, family = multinomial(parallel = FALSE,
  refLevel = 3), data = school)

```

Coefficients:

	Estimate	Std. Error	z value	Pr(> z)	
(Intercept):1	-1.2233	0.3154	-3.879	0.000105	***
(Intercept):2	-0.5662	0.2586	-2.189	0.028593	*
School2:1	1.0828	0.3539	3.059	0.002218	**
School2:2	0.1801	0.3172	0.568	0.570163	
School3:1	1.3147	0.3839	3.424	0.000616	***
School3:2	0.6556	0.3395	1.931	0.053455	.
ProgramAfter:1	-0.7474	0.2820	-2.651	0.008028	**
ProgramAfter:2	-0.7426	0.2706	-2.745	0.006057	**

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Names of linear predictors: log(mu[,1]/mu[,3]), log(mu[,2]/mu[,3])

Residual deviance: 666.9339 on 668 degrees of freedom

Log-likelihood: -333.4669 on 668 degrees of freedom

Number of Fisher scoring iterations: 4

Reference group is level 3 of the response

```

library(car)
Anova(modv)

```

Analysis of Deviance Table (Type II tests)

Response: Style

```

      Df  Chisq Pr(>Chisq)
School   4 14.843   0.005039 **
Program   2 10.916   0.004262 **

```

```
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
# Odds Ratio
exp(coef(modv))
```

```

(Intercept):1 (Intercept):2 School2:1 School2:2 School3:1
0.2942532      0.5676812      2.9530080      1.1973459      3.7236794
School3:2 ProgramAfter:1 ProgramAfter:2
1.9262906      0.4735774      0.4758599

```

```
exp(confint(modv))
```

```

      2.5 %    97.5 %
(Intercept):1 0.1585883 0.5459732
(Intercept):2 0.3419345 0.9424668
School2:1      1.4757049 5.9092141
School2:2      0.6430168 2.2295485
School3:1      1.7545863 7.9025968
School3:2      0.9902930 3.7469675
ProgramAfter:1 0.2725131 0.8229898
ProgramAfter:2 0.2800064 0.8087053

```

3.6.3 Answer

There is sufficient evidence of a program ($\chi^2 = 10.9, p = 0.004$) and school ($\chi^2 = 14.8, p = 0.005$).

The odds of preferring self-directed learning for those assigned to after-hours care is 0.47 (0.27-0.82) times that of those who assigned to regular class hours. The odds of students from school 2 preferring self-directed learning is 2.95 (1.48-5.91) times that of students from school 1. Similarly, the odds of students from school 3 preferring self-directed learning is 3.72 (1.76-7.90) times that of students from school 1.

The odds of preferring team-based learning for those assigned to after-hours care is 0.48 (0.27-0.82) times that of those who assigned to regular class hours. The odds of students from school 2 preferring self-directed learning is 1.20 (0.64-2.23) times that of students from school 1. Similarly, the odds of students from school 3 preferring self-directed learning is 1.93 (0.99-3.75) times that of students from school 1.

3.7 Exercise

The data set “firecracker” shows data of injuries due to firecrackers from three community health centers in rural Philippines. The age groups were coded as A: <25 years of age, B: 25 years of age. Use a generalized logit main effect model to determine whether there is a significant effect of age and clinic in the occurrence of face and finger injury compared to other parts of the body.

3.7.1 Question

Use the estimate statement to test and estimate the odds ratio between clinics 1 and 2 for fingers vs. other injury and face vs. other injury.

3.7.2 Data

```
data firecracker;
input clinic $ agegroup $ injury $ count;
datalines;
1 A Fingers 25
1 A Face 18
1 A Other 5
1 B Fingers 13
1 B Face 5
1 B Other 3
2 A Fingers 42
2 A Face 7
2 A Other 12
2 B Fingers 23
2 B Face 5
2 B Other 5
3 A Fingers 10
3 A Face 4
3 A Other 4
3 B Fingers 9
3 B Face 7
3 B Other 5
;
run;
```

3.7.3 R Data

```
firecracker <- expand.grid(Injury=c("Fingers","Face","Other"),
                          Agegroup=c("A","B"),
                          Clinic = c("1","2","3"))

firecracker$Count <- c(25,18,5,13,5,3,42,7,12,23,5,5,10,4,4,9,7,5)
firecracker <- firecracker |>
  uncount(weights=Count)
```

3.8 Exercise

The data set “firecracker” shows data of injuries due to firecrackers from three community health centers in rural Philippines. The age groups were coded as A: <25 years of age, B: 25 years of age. Use a generalized logit main effect model to determine whether there is a significant effect of age and clinic in the occurrence of face and finger injury compared to other parts of the body.

3.8.1 SAS Code

```
proc logistic data=firecracker order=data;
freq count;
class clinic agegroup;
model injury= clinic agegroup clinic*agegroup/link=glogit selection=forward include=2 scale=1;
estimate "clinic 1 vs clinic 2 - face" clinic 1 -1/ exp cl category="Face";
estimate "clinic 1 vs clinic 2 - fingers" clinic 1 -1/ exp cl category="Fingers";
run;
```

3.8.2 R Code

```
modv <- vglm(Injury~Clinic+Agegroup,
             data=firecracker,
             family=multinomial(parallel=FALSE, refLevel=3))

summary(modv)
```

Call:

```
vglm(formula = Injury ~ Clinic + Agegroup, family = multinomial(parallel = FALSE,
```

```
refLevel = 3), data = firecracker)
```

Coefficients:

	Estimate	Std. Error	z value	Pr(> z)
(Intercept):1	1.54406	0.40779	3.786	0.000153 ***
(Intercept):2	1.05057	0.43512	2.414	0.015759 *
Clinic2:1	-0.21911	0.47524	-0.461	0.644763
Clinic2:2	-1.40520	0.55777	-2.519	0.011758 *
Clinic3:1	-0.82186	0.56949	-1.443	0.148976
Clinic3:2	-0.85967	0.61912	-1.389	0.164975
AgegroupB:1	0.04654	0.40891	0.114	0.909377
AgegroupB:2	0.01831	0.48319	0.038	0.969767

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Names of linear predictors: log(mu[,1]/mu[,3]), log(mu[,2]/mu[,3])

Residual deviance: 367.3984 on 396 degrees of freedom

Log-likelihood: -183.6992 on 396 degrees of freedom

Number of Fisher scoring iterations: 4

Reference group is level 3 of the response

```
library(car)
Anova(modv)
```

Analysis of Deviance Table (Type II tests)

Response: Injury

	Df	Chisq	Pr(>Chisq)
Clinic	4	11.9012	0.0181 *
Agegroup	2	0.0154	0.9923

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
# Odds Ratio - note that the confidence interval limits are reversed.
exp(-coef(modv))
```

(Intercept):1 (Intercept):2 Clinic2:1 Clinic2:2 Clinic3:1

0.2135119	0.3497396	1.2449671	4.0763280	2.2747209
Clinic3:2	AgegroupB:1	AgegroupB:2		
2.3623847	0.9545231	0.9818537		

```
exp(-confint(modv))
```

	2.5 %	97.5 %
(Intercept):1	0.4748214	0.09600943
(Intercept):2	0.8205713	0.14906415
Clinic2:1	3.1599718	0.49049272
Clinic2:2	12.1629723	1.36615043
Clinic3:1	6.9450478	0.74504244
Clinic3:2	7.9496566	0.70202549
AgegroupB:1	2.1274023	0.42827548
AgegroupB:2	2.5312568	0.38085301

3.8.3 Answer

There is a significant effect of clinic ($\chi^2 = 11.9, p = 0.02$), but no evidence of an age effect ($\chi^2 = 0.02, p = 0.99$).

The odds of receiving patients with face injuries in clinic 1 is 4.08 (1.37-12.2) times that of clinic 2. Meanwhile, the odds of receiving patients with finger injuries in clinic 1 is 1.25 (0.49-3.16) times that of clinic 2.

3.9 Exact Generalized Logits Test

- Exact methods are more appropriate for tables with small or zero counts.

Exact Method Warning

Generalized logits tests have more parameters and exact methods are typically more computationally expensive. When using exact methods, try to keep the model small for relatively small data sets.

Exact Method in SAS

The **exact** statement can perform joint tests as well as calculate exact confidence intervals for odds ratios. Use the “Exact” p-value column in the “Probability” row in the exact conditional tests.

3.10 Example

The following data come from a study in which a physical therapy practice experimented with different treatment modalities in a strengthening program. The therapists rotated patients through three programs for maintenance after basic strength had been attained. The strengthening programs included free weights, resistance bands, and weight machines. The therapists are interested in whether gender and age affect choice of strengthening programs.

3.10.1 Data Set

```
data pt;
format therapy $11.;
input gender $ age $ therapy $ count @@;
datalines;
males adult machines 2 males senior machines 10
males adult freeweights 13 males senior freeweights 9
males adult bands 3 males senior bands 3
females adult machines 3 females senior machines 8
females adult freeweights 9 females senior freeweights 0
females adult bands 1 females senior bands 1
;
```

3.11 Example

The following data come from a study in which a physical therapy practice experimented with different treatment modalities in a strengthening program. The therapists rotated patients through three programs for maintenance after basic strength had been attained. The strengthening programs included free weights, resistance bands, and weight machines. The therapists are interested in whether gender and age affect choice of strengthening programs.



Warning

Small cell counts means we need to use exact methods.

3.11.1 SAS Code

```
proc logistic data=pt;
title "asymptotic";
freq count;
```



```

class gender(ref='females') age(ref='senior') / param=ref;
model therapy(ref='machines') = gender age / link=glogit scale=none aggregate;
run;

proc logistic data=pt;
title "exact";
freq count;
class gender(ref='females') age(ref='senior') / param=ref;
model therapy(ref='machines') = gender age / link=glogit scale=none aggregate;
exact gender age/joint estimate=both;
run;

```

3.11.2 Answer

There is a marginal effect of gender (\$ p=0.099), *but there is a significant effect of age* ($\chi^2 = 12.2$, p=0.003\$).

The odds of males preferring bands is 4.1 (0.49-59.55) times than that of females. Additionally, the odds of males preferring freeweights is 4.49 (0.93-30.2) times than that of females.

The odds of an adult preferring bands is 5.08 (0.60-53.2) times that of seniors. However, the odds of an adult preferring freeweights is 12.47 (2.7-84.0) times that of seniors.