

# Quantum chaos meets quantum channels

September 13, 2024

## 1 Model

The spin chain we are interested in studying first is that studied by Mirkin and Wisniacki in Ref. [1]:

$$H = \sum_{i=1}^L (h_x \sigma_i^x + h_z \sigma_i^z) - \sum_{i=1}^{L-1} J_z \sigma_i^z \sigma_{i+1}^z. \quad \text{eq:H:wisniacki:ising:chain} \quad (1)$$

## 2 Mean level spacing ratio

The level spacing ratio  $\tilde{r}_n$  is defined as:

$$\tilde{r}_n = \frac{\min(s_n, s_{n-1})}{\max(s_n, s_{n-1})}, \quad \text{eq:level:spacing:ratio} \quad (2)$$

where  $s_n = E_{n+1} - E_n$ . The mean level spacing ratio  $\langle \tilde{r}_n \rangle$  is known to attain the value  $\langle \tilde{r}_n \rangle \approx 0.5207$  when the level spacing distribution  $P(s)$  is Wigner-Dyson and  $\langle \tilde{r}_n \rangle \approx 0.386$  when it is Poisson.

## 3 Spectral form factor

The spectral form factor  $K(t)$  is defined as:

$$K(t) = \frac{1}{2^L} \left\langle |\text{Tr } U(t)|^2 \right\rangle = \frac{1}{2^L} \left\langle \sum_{i,j} e^{i(E_i - E_j)t} \right\rangle, \quad \text{eq:sff} \quad (3)$$

where  $\langle \cdot \rangle$  denotes the ensemble-average over statistically-similar systems.

## 4 Chaometer's quantum channel

The reduced dynamics of the chaometer is described by the quantum channel:

$$\mathcal{E}(\rho) = \text{Tr}_E \left( e^{-iHt} \rho \otimes \left| \psi_0^{(E)} \right\rangle \left\langle \psi_0^{(E)} \right| e^{iHt} \right), \quad \text{eq:chaometer:channel} \quad (4)$$

where  $H$  is that of eq. (1),  $\left| \psi_0^{(E)} \right\rangle$  the initial state of all spins except the chaometer, and  $\rho$  the initial state of the chaometer.

The chaometer's quantum channel  $\mathcal{E}$ , in general, is divisible into:

1. A unitary operation rotating the Bloch's sphere.
2. A quantum channel that deforms the Bloch's sphere and translates its origin.

Both operations do not commute.

JA: Comentar soluci3n de los estados aleatorios.




Figure 1: Mean level spacing ratio  $\langle \tilde{r}_n \rangle$  [c.f. eq. (2)] of the Ising chain with Hamiltonian (1) as a function of ratios  $h_z/h_x$  and  $J/h_x$ . We assume  $J_z = J \forall k$ . fig:mean:level:spacing:ratio




Figure 2: Spectral form factor (SFF) [c.f. (3)] in regular region:  $h_z/h_x = 2.5$  and  $J/h_x = 1$ . (a) Whole spectrum. (b) Even-parity subspace spectrum. fig:sff:regular




Figure 3: Spectral form factor (SFF) [c.f. (3)] in chaotic region:  $h_z/h_x = 0.5$  and  $J/h_x = 1$ . (a) Whole spectrum. (b) Even-parity subspace spectrum. fig:sff:chaotic

## 5 Purity of the chaometer

Averaged purity  $\mathcal{P}$  is defined in Ref. [1] as:

$$\bar{\mathcal{P}} = \frac{1}{N} \sum_{i=1}^N \left( \frac{1}{T} \int_0^T \text{Tr} [\rho_i^2(t)] \right) \quad \text{eq:avg:purity} \quad (5)$$

where:

- $\rho_i(t)$ : chaometer's density matrix.
- $N$ : number of different random initial states of the whole chain.  $N = 50$  in Mirkin and Wisniacki [1].
- $T$ : maximum time.  $T = 50$  in Mirkin and Wisniacki [1].

Also, the normalized averaged purity is defined as

$$\bar{\mathcal{P}}_{\text{Norm}} = \frac{\bar{\mathcal{P}} - \min(\bar{\mathcal{P}})}{\max(\bar{\mathcal{P}}) - \min(\bar{\mathcal{P}})}, \quad \text{eq:avg:norm:purity} \quad (6)$$

where  $\max(\bar{\mathcal{P}})$  and  $\min(\bar{\mathcal{P}})$  are the minimum and maximum value obtained when sweeping the parameter range ( $h_z$  in their case).

In Fig. 5 we plot one realization of the dynamics of purity of the chaometer. We compare in Fig. 6 our results and those of Mirkin and Wisniacki [1].

## 6 Purity of Choi-Jamiołkowski matrix

We investigate the purity of Choi-Jamiołkowski matrix of quantum channel  $\mathcal{E}(t)$  of the chaometer [c.f. eq. (4)] in Fig. 4.

To compute the Choi-Jamiołkowski matrix  $\mathcal{D}(t)$  of the chaometer's quantum channel in eq. (4) we use the definition  $(\mathcal{E} \otimes \mathbb{1})[|\phi^+\rangle\langle\phi^+|]$ , with  $|\phi^+\rangle = 1/\sqrt{2}(|0,0\rangle + |1,1\rangle)$  is the maximally entangled state between two spins, and obtain

$$\mathcal{D}(t) = \frac{1}{2} \sum_{i,j,p,q} \langle j, \psi_{0E} | U^\dagger(t) (|p\rangle\langle q| \otimes \mathbb{1}_E) U(t) | i, \psi_{0E} \rangle | q, i \rangle \langle p, j |. \quad \text{eq:choi:chaometer} \quad (7)$$




Figure 4: Purity of Choi-Jamiołkowski matrix in (a) regular ( $h_z = 2.5$ ) and (b) chaotic ( $h_z = 0.5$ ) for the three random initial states showed in the video. fig:choi:purity

Let us write  $\mathcal{E}(\rho)$  in computational basis. From eq. (4) we write

$$\mathcal{E}(\rho) = \sum_{k,l} e^{-i(E_k - E_l)t} \langle E_k | (\rho \otimes |\psi_E\rangle\langle\psi_E|) | E_l \rangle \text{Tr}_E (|E_k\rangle\langle E_l|) \quad (8)$$

$$= \sum_{k,l,p,p',\vec{q},\vec{q}'} e^{-i(E_k - E_l)t} \langle E_k | (\rho \otimes |\psi_E\rangle\langle\psi_E|) | E_l \rangle \text{Tr}_E (|p, \vec{q}\rangle \langle p, \vec{q}'| E_k \rangle \langle E_l | p', \vec{q}' \rangle \langle p', \vec{q}' |), \quad (9)$$

where  $\vec{q}$  and  $\vec{q}'$  are indices of elements of the basis of the environment,

$$= \sum_{p,p',\vec{q}} \langle p, \vec{q} | U(t) (\rho \otimes |\psi_E\rangle\langle\psi_E|) U^\dagger(t) | p', \vec{q} \rangle | p \rangle \langle p' | \quad (10)$$

$$= \sum_{p,p'} \langle p | \text{Tr}_E [U(t) (\rho \otimes |\psi_E\rangle\langle\psi_E|) U^\dagger(t)] | p' \rangle | p \rangle \langle p' |. \quad (11)$$

To compute the Choi-Jamiołkowski matrix  $\mathcal{D}(t)$  of the chaometer's quantum channel in eq. (4) we use the definition  $(\mathcal{E} \otimes \mathbb{1})[|\phi^+\rangle\langle\phi^+|]$ , with  $|\phi^+\rangle = 1/\sqrt{2}(|0,0\rangle + |1,1\rangle)$  is the maximally entangled state between two spins, and eq. (11) to obtain

$$\mathcal{D}(t) = \frac{1}{2} \sum_{i,j,p,p',\vec{q}} \langle p, \vec{q} | U(t) (|i, \psi_E\rangle\langle j, \psi_E|) U^\dagger(t) | p', \vec{q} \rangle | p, i \rangle \langle p', j | \quad (12)$$

$$= \frac{1}{2} \sum_{i,j,p,p'} \langle j, \psi_{0E} | U^\dagger(t) (|p\rangle\langle p| \otimes \mathbb{1}_E) U(t) | i, \psi_{0E} \rangle | p, i \rangle \langle p', j |, \quad (13)$$

where we have only reordered the factors and used the resolution of identity over  $\mathcal{H}_E$ .

We compute the purity of  $\mathcal{D}(t)$ . From eq. (7) it is straightforward to obtain

$$\text{Tr}[\mathcal{D}^2(t)] = \frac{1}{4} \sum_{i,j,p,q} \left| \langle i, \psi_{0E} | U^\dagger(t) (|p\rangle\langle q| \otimes \mathbb{1}_E) U(t) | j, \psi_{0E} \rangle \right|^2. \quad (14)$$

Moreover, reordering this expression an interesting interpretation of the purity of the Choi-Jamiołkowski matrix is revealed. Let us conveniently rewrite eq. (14) as

$$\text{Tr}[\mathcal{D}^2(t)] = \frac{1}{2} \sum_i \langle i, \psi_{0E} | U^\dagger(t) \sum_{p,q} \left[ \left( \frac{|p\rangle\langle q|}{\sqrt{2}} \otimes \mathbb{1}_E \right) U(t) \left( \frac{\mathbb{1}_S}{2} \otimes |\psi_{0E}\rangle\langle\psi_{0E}| \right) U^\dagger(t) \left( \frac{|q\rangle\langle p|}{\sqrt{2}} \otimes \mathbb{1}_E \right) \right] U(t) | i, \psi_{0E} \rangle. \quad (15)$$

In other words, the purity of  $\mathcal{D}(t)$  represents the probability that the environment remains in its initial state after the following process: First, the system and environment are initialized in a product state, with the system in a maximally mixed state. Next, the combined system evolves. Then, a completely depolarizing channel is applied to the system. Finally, the system and environment undergo reverse evolution.




Figure 5: Dynamics of chaometer's purity of a single realization with the same initial state of the environment of the first chaometer's quantum channel of the videos. fig:purity:one:realization

## 7 Non complete positiveness of $\Lambda(t, s)$

Any quantum channel  $\mathcal{E}(t)$  can be composed as

$$\mathcal{E}(t) = \Lambda(t, s) \circ \mathcal{E}(s, 0), \quad (16)$$

nonetheless,  $\Lambda(t, s)$  is not in general completely positive. A way to quantify how far is  $\Lambda(t, s)$  from being completely positive is through  $\tilde{\lambda}$ : eq:lambda:tilde

$$\tilde{\lambda} = |\min(0, \lambda_{\text{smallest}})|, \quad (17)$$

where  $\lambda_{\text{smallest}}$  is the smallest eigenvalue of  $\Lambda^R(t, s)/2$ . We have added the factor  $1/2$  just so  $\text{Tr}[\Lambda^R(t, s)/2] = 1$ .

Let us fix  $s = 0.1$  and investigate the complete positiveness of  $\Lambda(t, s)$ , see Fig. 7.

## References

- [1] Nicolás Mirkin and Diego Wisniacki. Quantum chaos, equilibration, and control in extremely short spin chains. *Phys. Rev. E*, 103:L020201, Feb 2021.




Figure 6: Averaged  $\bar{P}$  and averaged normalized purities  $\bar{P}_{Norm}$  [c.f. eqs. (5) and (6).] for  $N = 50$  random initial states of the chaometer for each quantum channel showed in the videos. *JA: Tendría más sentido sacar la pureza del canal. Lo pienso* Taken and modified from Ref. [1]. *fig:mirkin2021:fig2:modified*




Figure 7: Most negative eigenvalue  $\tilde{\lambda}$  of map  $\Lambda(t, s)/2$ , with  $s = 0.1$  [c.f. eqs. (16) and (17)]. *fig:lambda.index:Chir-“jami-”*




Figure 8: Burst of the Bloch sphere at  $t = 0.5$  s.

*fig:burst*