

# Fractal in Action

G. V. V. Sharma  
Dept. of Electrical Engg.,  
IIT Hyderabad.

June 15, 2024

- 1 Class 10
- 2 Class 12
- 3 CLASS 10
- 4 Market
- 5 Product
- 6 Financials
- 7 Team

## Question

The centre of a circle is at  $(2,0)$ . If one end of a diameter is at  $(6,0)$ , then the other end is at :

- ①  $(0,0)$
- ②  $(4,0)$
- ③  $(-2,0)$
- ④  $(-6,0)$

## Solution

Let

$$\mathbf{O} = \begin{pmatrix} 2 \\ 0 \end{pmatrix}, \mathbf{A} = \begin{pmatrix} 6 \\ 0 \end{pmatrix} \quad (1)$$

Then,

$$\mathbf{O} = \frac{(\mathbf{A} + \mathbf{B})}{2} \quad (2)$$

$$\Rightarrow \mathbf{B} = 2\mathbf{O} - \mathbf{A} \quad (3)$$

$$= 2 \begin{pmatrix} 2 \\ 0 \end{pmatrix} - \begin{pmatrix} 6 \\ 0 \end{pmatrix} = \begin{pmatrix} -2 \\ 0 \end{pmatrix} \quad (4)$$

## Question

2)  $AD$  is a median of  $\triangle ABC$  with vertices  $A(5, -6)$ ,  $B(6, 4)$  and  $C(0, 0)$ . Length  $AD$  is equal to:

- ①  $\sqrt{68}$
- ②  $2\sqrt{15}$
- ③  $\sqrt{101}$
- ④ 10

## Solution

The midpoint of **BC** is

$$\mathbf{D} = \frac{\mathbf{B} + \mathbf{C}}{2} \quad (5)$$

$$= \frac{1}{2} \begin{pmatrix} 6 \\ 4 \end{pmatrix} + \frac{1}{2} \begin{pmatrix} 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \end{pmatrix}, \quad (6)$$

Since

$$\mathbf{A} - \mathbf{D} = \begin{pmatrix} 5 \\ -6 \end{pmatrix} - \begin{pmatrix} 3 \\ 2 \end{pmatrix} = \begin{pmatrix} 2 \\ -8 \end{pmatrix} \quad (7)$$

$$\Rightarrow \|\mathbf{A} - \mathbf{D}\| \triangleq \sqrt{(\mathbf{A} - \mathbf{D})^\top (\mathbf{A} - \mathbf{D})} \quad (8)$$

$$= \sqrt{\begin{pmatrix} 2 & -8 \end{pmatrix} \begin{pmatrix} 2 \\ -8 \end{pmatrix}} = \sqrt{2^2 + 8^2} = \sqrt{68} \quad (9)$$

## Question

If the distance between the points  $(3, -5)$  and  $(x, -5)$  is 15 units, then the values of  $x$  are

- ①  $12, -18$
- ②  $-12, 18$
- ③  $18, 5$
- ④  $-9, -12$

## Solution

$$\mathbf{A} = \begin{pmatrix} 3 \\ -5 \end{pmatrix}, \mathbf{B} = \begin{pmatrix} x \\ -5 \end{pmatrix} \quad (10)$$

$$\Rightarrow \mathbf{A} - \mathbf{B} = \begin{pmatrix} 3 - x \\ -5 - (-5) \end{pmatrix} = \begin{pmatrix} 3 - x \\ 0 \end{pmatrix} \quad (11)$$

$$\Rightarrow \|\mathbf{A} - \mathbf{B}\| = \sqrt{(3 - x \quad 0) \begin{pmatrix} 3 - x \\ 0 \end{pmatrix}} = \sqrt{(3 - x)^2} \quad (12)$$

$$\Rightarrow 15 = \pm(3 - x) \quad (13)$$

$$\Rightarrow x = -12, 18 \quad (14)$$



## Question

Solve the following system of linear equations algebraically:

$$2x + 5y = -4; 4x - 3y = 5$$

## Solution

The above system of equations can be written as:

$$\begin{pmatrix} 2 & 5 \\ 4 & -3 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -4 \\ 5 \end{pmatrix} \quad (15)$$

Writing the augmented matrix for using Gauss elimination

$$\left( \begin{array}{cc|c} 2 & 5 & -4 \\ 4 & -3 & 5 \end{array} \right) \xleftarrow{R_2 \rightarrow R_2 - 2R_1} \left( \begin{array}{cc|c} 2 & 5 & -4 \\ 0 & -13 & 13 \end{array} \right) \quad (16)$$

$$\left( \begin{array}{cc|c} 2 & 5 & -4 \\ 0 & -13 & 13 \end{array} \right) \xleftarrow{R_1 \rightarrow \frac{13}{5}R_1 + R_2} \left( \begin{array}{cc|c} \frac{26}{5} & 0 & \frac{13}{5} \\ 0 & -13 & 13 \end{array} \right) \quad (17)$$

$$\Rightarrow \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} \frac{1}{2} \\ -1 \end{pmatrix} \quad (18)$$

## Question

- 5) The sum of the digits of a 2-digit number is 14. The number obtained by interchanging its digits exceeds the given number by 18. Find the number.

## Solution

Let the digits of the number be  $x_1$ (tens) and  $x_2$ (units). Given

$$x_1 + x_2 = 14 \quad (19)$$

$$10x_2 + x_1 = 18 + 10x_1 + x_2 \quad (20)$$

$$\implies x_1 - x_2 = -2 \quad (21)$$

Solving the equations (19),(21) in their matrix forms

$$\begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 14 \\ -2 \end{pmatrix} \quad (22)$$

$$(23)$$

## Solution

For the matrix

$$\mathbf{A} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix} \quad (24)$$

$$\mathbf{A}^\top \mathbf{A} = \mathbf{I} \quad (25)$$

$\mathbf{A}$  is an orthogonal matrix.

$$\begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 14 \\ -2 \end{pmatrix} \quad (26)$$

$$\implies 2\mathbf{I}\mathbf{x} = \begin{pmatrix} 12 \\ 16 \end{pmatrix} \quad (27)$$

$$\implies \mathbf{x} = \begin{pmatrix} 6 \\ 8 \end{pmatrix} \quad (28)$$

## Question

Find the ratio in which the point  $C \left( \frac{8}{5}, y \right)$  divides the line segment joining the points  $A(1, 2)$  and  $B(2, 3)$ . Also, find the value of  $y$ .

## Solution

For collinearity,

$$\text{rank} \begin{pmatrix} 1 & 1 & 1 \\ A & B & C \end{pmatrix} = 2 \quad (29)$$

$$\begin{pmatrix} 1 & 1 & 1 \\ 1 & 2 & 8/5 \\ 2 & 3 & y \end{pmatrix} \leftrightarrow \begin{pmatrix} 1 & 1 & 1 \\ 0 & (2-1) & (8/5-1) \\ 0 & (3-2) & (y-3) \end{pmatrix} \leftrightarrow \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 3/5 \\ 0 & 1 & y-3 \end{pmatrix} \quad (30)$$

$$\xleftrightarrow{R_3 \rightarrow R_3 - R_2} \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 3/5 \\ 0 & 0 & y - \frac{18}{5} \end{pmatrix} \implies y = \frac{18}{5} \quad (31)$$

## Question

$ABCD$  is a rectangle formed by the points  $A(-1, -1)$ ,  $B(-1, 6)$ ,  $C(3, 6)$  and  $D(3, -1)$ .  $P, Q, R$  and  $S$  are mid-points of sides  $AB, BC, CD$  and  $DA$  respectively. Show that the diagonal of the quadrilateral  $PQRS$  bisect each other.



## Solution

$$\mathbf{P} = \frac{\mathbf{A} + \mathbf{B}}{2}, \quad \mathbf{Q} = \frac{\mathbf{B} + \mathbf{C}}{2} \quad (32)$$

$$\mathbf{R} = \frac{\mathbf{C} + \mathbf{D}}{2}, \quad \mathbf{S} = \frac{\mathbf{D} + \mathbf{A}}{2} \quad (33)$$

Let  $\mathbf{O}_1$  and  $\mathbf{O}_2$  be the midpoints of  $PR$  and  $QS$  respectively

$$\mathbf{O}_1 = \frac{\mathbf{P} + \mathbf{R}}{2} = \frac{\mathbf{A} + \mathbf{B} + \mathbf{C} + \mathbf{D}}{4} \quad (34)$$

$$\mathbf{O}_2 = \frac{\mathbf{Q} + \mathbf{S}}{2} = \frac{\mathbf{A} + \mathbf{B} + \mathbf{C} + \mathbf{D}}{4} \quad (35)$$

Since the midpoints of the diagonals coincide, the diagonals bisect each other.

# Topics covered so far

- ① Vectors
- ② Section Formula
- ③ Norm
- ④ Gauss Elimination
- ⑤ Orthogonal matrix
- ⑥ Rank

## Question

If  $\vec{a} = 2\hat{i} - \hat{j} + \hat{k}$  and  $\vec{b} = \hat{i} + \hat{j} - \hat{k}$ , then  $\vec{a}$  and  $\vec{b}$  are:

- ① Collinear vectors which are not parallel
- ② Parallel vectors
- ③ Perpendicular vectors
- ④ Unit vectors

## Solution

Let

$$\mathbf{a} = \begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix}, \mathbf{b} = \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix} \quad (36)$$

Applying concept of rank from (29)

$$\text{rank} \begin{pmatrix} 1 & 1 & -1 \\ 2 & -1 & 1 \end{pmatrix} = 2 \neq 1, \text{ Not parallel} \quad (37)$$

Applying condition for perpendicularity:

$$\mathbf{a}^\top \mathbf{b} = (2 \quad -1 \quad 1) \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix} = 0 \implies \mathbf{a} \perp \mathbf{b} \quad (38)$$

## Question

2) If  $\alpha, \beta$  and  $\gamma$  are the angles which a line makes with positive directions of  $x, y$  and  $z$  axes respectively, then which of the following are not true?

- ①  $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$
- ②  $\sin^2 \alpha + \sin^2 \beta + \sin^2 \gamma = 2$
- ③  $\cos 2\alpha + \cos 2\beta + \cos 2\gamma = -1$
- ④  $\cos \alpha + \cos \beta + \cos \gamma = 1$

## Solution

Let  $\mathbf{m}$  represent the unit direction vector of the line. Then,

$$\mathbf{m} = \begin{pmatrix} \cos \alpha \\ \cos \beta \\ \cos \gamma \end{pmatrix} \quad (39)$$

with

$$\|\mathbf{m}\| = 1 \quad (40)$$

From (15)

$$2x + 5y = -4 \quad (41)$$

can be expressed as

$$\mathbf{n}^T \mathbf{x} = c \quad (42)$$

where

$$\mathbf{n} = \begin{pmatrix} 2 \\ 5 \end{pmatrix}, \quad c = -4 \quad (43)$$

## Solution

This is defined to be the normal equation of a line, where  $\mathbf{n}$  represents the normal vector. Also,

$$2x + 5y = -4 \quad (44)$$

$$\implies 2x = -4 - 5y \quad (45)$$

$$\implies \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -2 \\ 0 \end{pmatrix} + y \begin{pmatrix} -\frac{5}{2} \\ 1 \end{pmatrix} \quad (46)$$

$$\mathbf{x} = \begin{pmatrix} -2 \\ 0 \end{pmatrix} - \frac{5y}{2} \begin{pmatrix} 1 \\ -\frac{2}{5} \end{pmatrix} \quad (47)$$

$$= \mathbf{A} + k\mathbf{m} \quad (48)$$

The general equation written above is of the parametric form.

## Question

$\vec{a}, \vec{b}$  and  $\vec{c}$  are three mutually perpendicular unit vectors. If  $\theta$  is the angle between  $\vec{a}$  and  $(2\vec{a} + 3\vec{b} + 6\vec{c})$ , find the value of  $\cos \theta$ .



## Solution

Given:

$$\mathbf{a}^\top \mathbf{b} = \mathbf{b}^\top \mathbf{c} = \mathbf{c}^\top \mathbf{a} = 0 \quad (49)$$

$$||\mathbf{a}|| = ||\mathbf{b}|| = ||\mathbf{c}|| = 1 \quad (50)$$

$$\cos \theta = \frac{\mathbf{a}^\top (2\mathbf{a} + 3\mathbf{b} + 6\mathbf{c})}{||\mathbf{a}||||2\mathbf{a} + 3\mathbf{b} + 6\mathbf{c}||} \quad (51)$$

Now,

$$\mathbf{a}^\top (2\mathbf{a} + 3\mathbf{b} + 6\mathbf{c}) = 2\mathbf{a}^\top \mathbf{a} + 3\mathbf{a}^\top \mathbf{b} + 6\mathbf{a}^\top \mathbf{c} = 2 + 0 + 0 = 2 \quad (52)$$

$$||\mathbf{a}||||2\mathbf{a} + 3\mathbf{b} + 6\mathbf{c}|| = ||2\mathbf{a} + 3\mathbf{b} + 6\mathbf{c}|| \quad (53)$$

## Solution

From (8) norm definition:

$$(\|2\mathbf{a} + 3\mathbf{b} + 6\mathbf{c}\|)^2 = \|4\mathbf{a}^2\| + \|9\mathbf{b}^2\| + \|36\mathbf{c}^2\| = 49 \quad (54)$$

$$\implies \|2\mathbf{a} + 3\mathbf{b} + 6\mathbf{c}\| = +7 \quad (55)$$

$$\implies \cos \theta = \frac{2}{7} \quad (56)$$

## Question

Find the position vector of point **C** which divides the line segment joining points **A** and **B** having position vectors  $\hat{i} + 2\hat{j} - \hat{k}$  and  $-\hat{i} + \hat{j} + \hat{k}$  respectively in the ratio 4 : 1 externally. Further, find  $|\overrightarrow{AB}| : |\overrightarrow{BC}|$ .

## Solution

We know that

$$\mathbf{C} = \frac{4\mathbf{B} - \mathbf{A}}{4 - 1} \quad (57)$$

Simplify the above for  $\mathbf{C}$ .

## Question

- 5) Find the equation of the line passing through the point of intersection of the lines  $\frac{x}{1} = \frac{y-1}{2} = \frac{z-2}{3}$  and  $\frac{x-1}{0} = \frac{y}{-3} = \frac{z-7}{2}$  and perpendicular to these given lines.

## Solution

Let the given lines be denoted by  $\mathbf{x}_1$  and  $\mathbf{x}_2$  respectively. From (48):

$$\mathbf{x}_1 = \begin{pmatrix} 0 \\ 1 \\ 2 \end{pmatrix} + k_1 \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} = \mathbf{A} + k_1 \mathbf{m}_1 \quad (58)$$

$$\mathbf{x}_2 = \begin{pmatrix} 1 \\ 0 \\ 7 \end{pmatrix} + k_2 \begin{pmatrix} 0 \\ -3 \\ 2 \end{pmatrix} = \mathbf{B} + k_2 \mathbf{m}_2 \quad (59)$$

## Solution

The two equations required to solve for the direction of line are

$$\mathbf{m}^T \mathbf{m}_1 = 0 \quad (60)$$

$$\mathbf{m}^T \mathbf{m}_2 = 0 \quad (61)$$

$$\Rightarrow (\mathbf{m}_1 \quad \mathbf{m}_2)^T \mathbf{m} = 0 \quad (62)$$

$$\begin{pmatrix} 1 & 2 & 3 \\ 0 & -3 & 2 \end{pmatrix} \mathbf{m} \xleftarrow{R_1 \rightarrow 2R_2 + 3R_1} \begin{pmatrix} 3 & 0 & 13 \\ 0 & -3 & 2 \end{pmatrix} \mathbf{m} = 0 \quad (63)$$

$$\Rightarrow \begin{pmatrix} 3 & 0 & 13 \\ 0 & -3 & 2 \end{pmatrix} \begin{pmatrix} m_x \\ m_y \\ m_z \end{pmatrix} = 0 \quad (64)$$

$$\Rightarrow \mathbf{m} = \begin{pmatrix} \frac{-13}{3} \\ \frac{2}{3} \\ 1 \end{pmatrix} \quad (65)$$

## Solution

Let the unknown line in its parametric form be denoted as follows from (48).

$$\mathbf{x}_3 = \mathbf{C} + k_3 \mathbf{m} \quad (66)$$

For finding the point of intersection of the lines( $\mathbf{C}$ ), equating (58) and (59)

$$\mathbf{A} + k_1 \mathbf{m}_1 = \mathbf{B} + k_2 \mathbf{m}_2 \quad (67)$$

$$(\mathbf{m}_1 \quad \mathbf{m}_2) \begin{pmatrix} k_1 \\ -k_2 \end{pmatrix} = \mathbf{B} - \mathbf{A} \quad (68)$$

From the above  $k_1$  and  $k_2$  can be found by gauss elimination given in (17) and thus  $\mathbf{C}$ .



## Question

Two vertices of the parallelogram **ABCD** are given as **A**  $(-1, 2, 1)$  and **B**  $(1, -2, 5)$ . If the equation of the line passing through **C** and **D** is  $\frac{x-4}{1} = \frac{y+7}{-2} = \frac{z-8}{2}$ , then find the distance between the sides **AB** and **CD**. Hence, find the area of parallelogram **ABCD**

## Solution

Let the two parallel lines be

$$\mathbf{x} = \mathbf{A} + k_1 \mathbf{m} \quad (69)$$

$$\mathbf{x} = \mathbf{B} + k_2 \mathbf{m} \quad (70)$$

If  $\mathbf{P}$  be a point on the second line,

$$\mathbf{P} = \mathbf{B} + k_2 \mathbf{m} \quad (71)$$

$$(\mathbf{A} - \mathbf{P})^\top \mathbf{m} = 0 \quad (72)$$

From the above,

$$(\mathbf{A} - \mathbf{B})^\top \mathbf{m} - k_2 \|\mathbf{m}\|^2 = 0 \quad (73)$$

$$\implies k_2 = \frac{(\mathbf{A} - \mathbf{B})^\top \mathbf{m}}{\|\mathbf{m}\|^2} \quad (74)$$

## Question

- 8) The sum of first and eight terms of an A.P is 32 and their product is 60. Find the first term and common difference of the A.P. Hence, also find the sum of its first 20 terms.

## Solution

Let the first and eighth terms be  $x(0)$  and  $x(7)$  respectively, given:

$$x(0) + x(7) = 32 \quad (75)$$

$$x(0) = 32 - x(7) \quad (76)$$

$$x(0)x(7) = 60 \quad (77)$$

From (76) and (77)

$$x(7)(32 - x(7)) = 60 \quad (78)$$

The roots are  $(30, 2)$ , therefore, if  $x(7) = 30$  then  $x(0) = 2$  and if  $x(7) = 2$  then  $x_0 = 30$

Now

$$x(n) = (x(0) + nd)u(n) \quad (79)$$

Where  $d$  is the common difference of the A.P and  $u(n)$  is the unit step function.

$$(u(n) = 0 \forall n < 0, u(n) = 1 \forall n \geq 0)$$

## Solution

$$\implies x(7) = (x(0) + 7d) \quad (80)$$

$$\implies 7d = \pm 28 \implies d = \pm 4 \quad (81)$$

Therefore the A.P is 2, 6, 10... or 30, 26, 22....

Considering the former for calculations and taking Z-Transform of (79) for sum.

Since

$$X(z) = \sum_{n=-\infty}^{\infty} x(n)z^{-n} \quad (82)$$

Let  $y(n)$  denote the sum, let:

$$y(n) = x(n) * h(n) \quad (83)$$

$$= \sum_{k=-\infty}^{\infty} x(k)h(n-k) \quad (84)$$

## Solution

Replace  $h(n)$  with  $u(n)$ .

$$y(n) = \sum_{k=0}^n x(k)u_{(n-k)} \quad (85)$$

$$= x(0)u(n) + x(1)u(n-1) + \dots x(n)u_{(0)} \quad (86)$$

This denotes the sum of terms  $x(0), x(1) \dots x(n)$  i.e. first  $n+1$  terms.  
From (82)

$$u(n) \xrightarrow{Z} \frac{1}{(1-z^{-1})} \quad (87)$$

$$nu(n) \xrightarrow{Z} \frac{z^{-1}}{(1-z^{-1})^2} \quad (88)$$

$$\Rightarrow X(z) = \frac{2}{(1-z^{-1})} + \frac{4z^{-1}}{(1-z^{-1})^2}, \quad |z| > |1| \quad (89)$$

## Solution

Now as convolution in the time domain corresponds to multiplication in the frequency domain and (89) and (87).

$$Y(z) = X(z) * H(z) \quad (90)$$

$$= \left( \frac{2}{(1 - z^{-1})} + \frac{4z^{-1}}{(1 - z^{-1})^2} \right) \left( \frac{1}{(1 - z^{-1})} \right), \quad |z| > |1| \quad (91)$$

Using normal inversion for inverse Z-transform:

$$Y(z) = \frac{2}{(1 - z^{-1})^2} + \frac{4z^{-1}}{(1 - z^{-1})^3}, \quad |z| > |1| \quad (92)$$

$$= \frac{8z^{-1}}{1 - z^{-1}} + \frac{10z^{-2}}{(1 - z^{-1})^2} + \frac{4z^{-3}}{(1 - z^{-1})^3} + 2 \quad (93)$$

## Solution

For proceeding forwards here are some important generalizations.

Shifting property

$$x(n - k) \leftrightarrow z^{-k}X(z) \quad (94)$$

Differentiation property

$$nx(n) \leftrightarrow -zX'(z) \quad (95)$$

From (87) and (94)

$$u(n - 1) \xrightarrow{\mathcal{Z}} \frac{z^{-1}}{1 - z^{-1}} \quad (96)$$

From (87) and (95)

$$nu(n) \xrightarrow{\mathcal{Z}} -z \frac{d}{dz} \left( \frac{1}{1 - z^{-1}} \right) \quad (97)$$

$$nu(n) \xrightarrow{\mathcal{Z}} \frac{z^{-1}}{(1 - z^{-1})^2} \quad (98)$$



## Solution

From (94)

$$(n-1)u(n-1) \xrightarrow{Z} z^{-1} \frac{z^{-1}}{(1-z^{-1})^2} \quad (99)$$

$$(n-1)u(n-1) \xrightarrow{Z} \frac{z^{-2}}{(1-z^{-1})^2} \quad (100)$$

Now, using (95) and writing the corresponding L.H.S

$$(n)(n-1)u(n-1) \xrightarrow{Z} \frac{2z^{-2}}{(1-z^{-1})^3} \quad (101)$$

Using (94)

$$\frac{(n-1)(n-2)u(n-2)}{2} \xrightarrow{Z} \frac{z^{-3}}{(1-z^{-1})^3} \quad (102)$$

## Solution

The inverse-Z of a constant will be  $\delta(n)$ , so it is ruled out. Plugging these values in (93) we get

$$y(n) = 8u(n-1) + 10(n-1)u(n-1) + 4\frac{(n-1)(n-2)u(n-2)}{2} + 2\delta(n) \quad (103)$$

Putting  $n = 19$

$$y(19) = 2(19+1)^2 = 800 \quad (104)$$

## Solution

Using contour integration for inverse Z-transform

$$y(19) = \frac{1}{2\pi j} \oint_C Y(z) z^{18} dz \quad (105)$$

$$= \frac{1}{2\pi j} \oint_C \left( 2z^{20} (z-1)^{-2} + 4z^{20} (z-1)^{-3} \right) dz \quad (106)$$

$$R = \frac{1}{(m-1)!} \lim_{z \rightarrow a} \frac{d^{m-1}}{dz^{m-1}} ((z-a)^m f(z)) \quad (107)$$

For  $R_1$ ,  $m = 2$ , where  $m$  corresponds to number of repeated poles.

$$R_1 = \frac{1}{(1)!} \lim_{z \rightarrow 1} \frac{d}{dz} \left( (z-1)^2 2z^{20} (z-1)^{-2} \right) \quad (108)$$

$$= 2 \lim_{z \rightarrow 1} \frac{d}{dz} (z^{20}) \quad (109)$$

$$= 40 \quad (110)$$

## Solution

$$R_2 = \frac{1}{(2)!} \lim_{z \rightarrow 1} \frac{d^2}{dz^2} \left( (z-1)^3 4z^{20} (z-1)^{-3} \right) \quad (111)$$

$$= (2) \lim_{z \rightarrow 1} \frac{d^2}{dz^2} (z^{20}) \quad (112)$$

$$= 760 \quad (113)$$

$$R_1 + R_2 = 800 \quad (114)$$

$$\implies y(19) = 800 \quad (115)$$

Similarly, the sum for the A.P. 30, 26, 22... can be found by the same procedure.

## Question

- 9) In an A.P. of 40 terms, the sum of first 9 terms is 153 and the sum of last 6 terms is 687. Determine the first term and the common difference of the A.P. Also find the sum of all the terms of the A.P.

Given:

$$y(8) = 153 \quad (116)$$

$$y(39) - y(34) = 687 \quad (117)$$

Now, let the first term be  $x(0)$  and common difference be  $d$ . From (82) and (79)

$$X(z) = \frac{x(0)}{1 - z^{-1}} + \frac{dz^{-1}}{(1 - z^{-1})^2}, \quad |z| > |1| \quad (118)$$

For finding the sum (Assuming  $h(n) = u(n)$ )

$$y(n) = x(n) * h(n) \quad (119)$$

$$Y(z) = X(z) * H(z) \quad (120)$$

$$= \left( \frac{x(0)}{(1 - z^{-1})} + \frac{dz^{-1}}{(1 - z^{-1})^2} \right) \left( \frac{1}{(1 - z^{-1})} \right), \quad |z| > |1| \quad (121)$$

## Solution

$$Y(z) = \frac{(2x(0) + d)z^{-1}}{1 - z^{-1}} + \frac{(x(0) + 2d)z^{-2}}{(1 - z^{-1})^2} + \frac{dz^{-3}}{(1 - z^{-1})^3} + x(0) \quad (122)$$

Using normal inversion for inverse Z-transform:

Using the results (96),(100) and (102)

$$y(n) = (2x_0 + d)u(n-1) + (n-1)u(n-1)(x_0 + 2d) + \frac{d(n-1)(n-2)u(n-2)}{2} + x(0)\delta(n) \quad (123)$$

Now use (116) and (117) to solve for  $x(0)$  and  $d$  and put in (133) for the sum of 40 terms.

## Solution

Using contour integration for inverse Z-transform

$$y(n) = \frac{1}{2\pi j} \oint_C Y(z) z^{n-1} dz \quad (124)$$

$$= \frac{1}{2\pi j} \oint_C \left( x(0)z^{n+1} (z-1)^{-2} + dz^{20} (z-1)^{-3} \right) dz \quad (125)$$

$$R = \frac{1}{(m-1)!} \lim_{z \rightarrow a} \frac{d^{m-1}}{dz^{m-1}} ((z-a)^m f(z)) \quad (126)$$

For  $R_1$  ,  $m = 2$  , where  $m$  corresponds to number of repeated poles .

$$R_1 = \frac{1}{(1)!} \lim_{z \rightarrow 1} \frac{d}{dz} \left( (z-1)^2 x(0)z^{n+1} (z-1)^{-2} \right) \quad (127)$$

$$= x(0) \lim_{z \rightarrow 1} \frac{d}{dz} (z^{n+1}) \quad (128)$$

$$= (n+1) x(0) \quad (129)$$



## Solution

For  $R_2$ ,  $m = 3$

$$R_2 = \frac{1}{(2)!} \lim_{z \rightarrow 1} \frac{d^2}{dz^2} \left( (z-1)^3 dz^{n+1} (z-1)^{-3} \right) \quad (130)$$

$$= \left( \frac{d}{dz} \right) \lim_{z \rightarrow 1} \frac{d^2}{dz^2} (z^{n+1}) \quad (131)$$

$$= \left( \frac{d}{dz} \right) (n)(n+1) \quad (132)$$

$$y(n) = R_1 + R_2 = \left( \frac{n+1}{2} \right) (2x(0) + nd) \quad (133)$$

Now use (116) and (117) to solve for  $x(0)$  and  $d$  and put in (133) for the sum of 40 terms.

# Solution

- Hardware products to be available at low cost.
- Related skill sets to be provided to the public through simple projects.

# Market

- Students
- Working professionals
- Industry and academia

# Product




Figure: Decimal Game

# Financials

- Investment - negligible
- Expected profit -  $> 50\%$

# Team

- G V V Sharma
- Faculty in the EE Dept at IITH
- 22 years exp.
- Currently running the FWC certificate program at IITH.
- Leading the SatCom 6G team at IITH.