Supplementary Table S1 Neonatal outcomes

	Daily iron group ($N = 39$)			Alternate-day iron group ($N = 36$)			
	N	Median (IQR)	Range	*N*	Median (IQR)	Range	*p*-Value
GA at delivery (wk)	39	39 (38–39.6)	26.9–41	36	38.9 (37.7–39.4)	26–40.4	0.68
Birthweight of babies (g)	39	3,240 (2,910–3,390)	1,180–4,170	36	3,220 (2,937.5–3,505)	650–4,140	0.50
Apgar 1 min	39	8 (8–9)	7–9	36	9 (8–9)	2–9	0.91
Apgar 5 min	39	9 (9–9)	8–9	36	9 (9–9)	6–9	0.39
	No./no. observed (%)			No./no. observed (%)			
Neonatal complications							
Any[b]	4/39 (11)			5/36 (14)			0.73
Hypoglycemia	1/39 (3)			1/36 (3)			>0.99
Jaundice	1/39 (3)			1/36 (3)			>0.99
Prematurity	2/39 (5)			2/36 (6)			>0.99
Transient tachypnea	0/39 (0)			1/36 (3)			0.48
Congenital cystic adenomatoid malformation	1/39 (3)			0/36 (0)			>0.99
Very low birthweight	1/39 (3)			0/36 (0)			>0.99
Respiratory distress	1/39 (3)			2/36 (6)			0.60
Maternal postpartum complications							
Any	3/39 (3)			3/36 (3)			>0.99
Endometriosis treated with antibiotics	1/39 (3)			1/36 (3)			>0.99
Gestational hypertension	1/39 (3)			0/36 (0)			>0.99
Postpartum hemorrhage	0/39 (0)			1/36 (3)			0.48
Preeclampsia without severe features	0/39 (0)			1/36 (3)			0.48
Symptomatic anemia	1/39 (3)			0/36 (0)			>0.99
Adverse pregnancy outcomes							
NICU admission	3/38 (8)			5/36 (14)			0.48
Need for IV Iron	1/39 (3)			1/36 (3)			>0.99
Blood transfusion predelivery	0/37 (0)			0/35 (0)			–
Blood transfusion postdelivery	1/39 (3)			1/36 (3)			>0.99

Abbreviations: IQR, interquartile range; IV, intravenous; NICU, neonatal intensive care unit.
[a]$N = 75$ patients delivered by time of analysis.
[b]Some children had >1 neonatal complication, so the "any" neonatal complication row is less than the sum of all individual complications.

 © 2024. Thieme. All rights reserved.

Supplementary Table S2 Iron labs at baseline and 6 weeks after treatment among compliant patients

	Randomization group	Baseline			6 wk			Change (6 wk − baseline)				
		No.	Median (IQR)	Mean ± SD	No.	Median (IQR)	Mean ± SD	No.	Median (IQR)	Mean ± SD	Baseline adjusted difference in means (95% CI)[a]	*p*-Value
Hb	Daily	36	10.6 (10.3,10.8)	10.5 ± 0.5	34	11.3 (10.6,11.8)	11.4 ± 1.0	34	0.9 (0.3,1.5)	0.9 ± 0.9		
	Alternate	29	10.7 (10.3,10.8)	10.5 ± 0.5	29	11.3 (10.7,11.6)	11.1 ± 1.2	29	0.8 (0, 1)	0.7 ± 1.0	−0.2 (−0.7,0.2)	0.32
Hct	Daily	36	32 (30.6,33)	31.9 ± 1.6	34	34.5 (33,35.6)	34.6 ± 3.3	34	2.5 (1.3,4.5)	2.7 ± 2.9		
	Alternate	29	31.5 (29.7,32.7)	30.2 ± 4.5	29	34 (32.7,35.4)	33.9 ± 3.5	29	2.7 (0.6,4.1)	3.6 ± 5.5	−0.4 (−2.2,1.3)	0.63
Ferritin	Daily	36	11.5 (8.5,15)	12.2 ± 5.3	33	20 (12,28)	24.3 ± 24.3	33	7 (3,15)	12 ± 23.3		
	Alternate	29	13 (10,17)	13.6 ± 5.3	27	21 (14,26)	21.3 ± 10.4	27	8 (1,12)	8.1 ± 8.5	−4.1 (−13.7,5.4)	0.39
Hepcidin	Daily	26	8.7 (6.6,10.7)	9.9 ± 3.8	30	9.5 (8.5,11.1)	11.1 ± 6.2	26	0.1 (−2.1,3.5)	1.6 ± 6.5		
	Alternate	21	7.6 (6.3,13.4)	10.1 ± 5.0	25	8.9 (7.9,10.3)	9.5 ± 3	21	−0.9 (−5,2.6)	−0.5 ± 6.2	−2.0 (−5.3,1.3)	0.23

Abbreviations: CI, confidence interval; Hb, hemoglobin; Hct, hematocrit; IQR, interquartile range; SD, standard deviation.

 © 2024. Thieme. All rights reserved.

Supplementary Table S3 Side effects by randomization group among compliant patients

Iron side effects	Daily iron group ($N = 36$) No./no. observed (%)	Alternate-day iron group ($N = 29$) No./no. observed (%)	*p*-Value
Constipation	12/36 (33)	9/29 (31)	>0.99
Nausea	12/36 (33)	10/29 (34)	>0.99
Vomiting	6/36 (17)	5/29 (17)	>0.99
Abdominal pain	1/36 (3)	1/29 (3)	>0.99
Diarrhea	0/36 (0)	1/29 (3)	0.45
Headache	3/36 (8)	4/29 (14)	0.69
Other	2/36 (6)	2/29 (7)	>0.99

 © 2024. Thieme. All rights reserved.