

Questions

1. Show that a projective transformation maps a quadric surface to a quadric surface.

Here, by a “projective transformation”, I mean any invertible 4×4 matrix that operates on 3D points which are written in 4D homogeneous coordinates.

2. Give the 4×4 projective transformation matrix that is defined by

`gluPerspective(theta, aspect, near, far)` .

Rather than just plugging in the values given in the lecture, think about the special constraints on `gluPerspective`, as opposed to `glFrustum`.

3. How would one choose the first four arguments of `glFrustum` to achieve the same view volume as `gluPerspective(60, 2, 4, 20)` ?
4. Show that a 4×4 projective matrix \mathbf{M} maps lines to lines.
5. In the lecture I showed that when \mathbf{M} maps a plane π to a plane, the normal \mathbf{n} of the mapped plane $\mathbf{M}\pi$ is $(\mathbf{M}^{-1})^T \mathbf{n}$. Suppose you are given a polygon (and say its normal too) and you apply \mathbf{M} to the vertices of a polygon and to the normal. Suppose you want to know the normal of the polygon in the mapped space. Do you need to know \mathbf{M}^{-1} to compute this new normal?
6. **[ADDED Feb. 15]**

- (a) Write the ellipsoid

$$(x - 3)^2 + y^2 + 4z^2 = 20,$$

in the form $\mathbf{x}^T \mathbf{Q} \mathbf{x} = 0$ where $\mathbf{x}^T = (x, y, z, 1)$.

- (b) What is the equation of this ellipsoid after it has undergone a projective transformation to the view volume between planes $z = -1$ and $z = -8$.

Hint: It is sufficient for you to write your answer as a product of matrices and vectors. One of these matrices should be of the form:

$$\mathbf{M} = \begin{bmatrix} f_0 & 0 & 0 & 0 \\ 0 & f_0 & 0 & 0 \\ 0 & 0 & f_0 + f_1 & -f_0 f_1 \\ 0 & 0 & 1 & 0 \end{bmatrix}.$$

Answers

1. Consider an invertible 4×4 matrix \mathbf{M} . Let \mathbf{x} be the 4D column vector $(x, y, z, 1)^T$. Then,

$$\begin{aligned} 0 &= \mathbf{x}^T \mathbf{Q} \mathbf{x} \\ &= \mathbf{x}^T \mathbf{M}^T (\mathbf{M}^T)^{-1} \mathbf{Q} \mathbf{M}^{-1} \mathbf{M} \mathbf{x} \\ &= (\mathbf{M} \mathbf{x})^T (\mathbf{M}^T)^{-1} \mathbf{Q} \mathbf{M}^{-1} \mathbf{M} \mathbf{x} \end{aligned}$$

For a quadric surface, we have a symmetric matrix \mathbf{Q} , i.e. $(\mathbf{Q} = \mathbf{Q}^T)$, and so the matrix $(\mathbf{M}^T)^{-1} \mathbf{Q} \mathbf{M}^{-1}$ is also symmetric, as we show below. Hence, the transformed points $\mathbf{M} \mathbf{x}$ also lie on a quadric surface.

To show that

$$((\mathbf{M}^T)^{-1} \mathbf{Q} \mathbf{M}^{-1})^T = (\mathbf{M}^T)^{-1} \mathbf{Q} \mathbf{M}^{-1}$$

use the property that $(\mathbf{A} \mathbf{B})^T = \mathbf{B}^T \mathbf{A}^T$. We get

$$((\mathbf{M}^T)^{-1} \mathbf{Q} \mathbf{M}^{-1})^T = (\mathbf{M}^{-1})^T \mathbf{Q}^T ((\mathbf{M}^T)^{-1})^T$$

Now use the property that $(\mathbf{M}^T)^{-1} = (\mathbf{M}^{-1})^T$ for any invertible \mathbf{M} , and \mathbf{Q} symmetric, and we are done.

2. OpenGL first applies (lecture 5, bottom p. 3)

$$\mathbf{M}_{projective} = \begin{bmatrix} \text{near} & 0 & 0 & 0 \\ 0 & \text{near} & 0 & 0 \\ 0 & 0 & \text{near} + \text{far} & \text{near} * \text{far} \\ 0 & 0 & -1 & 0 \end{bmatrix}.$$

OpenGL then normalizes the view volume so it maps to $[-1, 1] \times [-1, 1] \times [-1, 1]$. It does so in a few steps. First, the x and y range of the viewing window is already centered on $(0, 0)$ so we don't need to translate it. We do need to translate z , however, to bring $z = -\text{near}$ to $z = 0$. After that, we scale each of the x, y, z dimensions, remembering to flip the z axis. The scale factors can be derived from the definitions of θ and **aspect** and use basic trigonometry. Finally, we translate so that the volume is centered at the origin. Assuming I have not made an error, the normalization is:

$$\mathbf{M}_{normalize} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \frac{2}{\text{near} * \text{aspect} * 2 * \tan(\frac{\theta}{2})} & 0 & 0 & 0 \\ 0 & \frac{2}{\text{near} * 2 * \tan(\frac{\theta}{2})} & 0 & 0 \\ 0 & 0 & \frac{2}{\text{near} - \text{far}} & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & \text{near} \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

3. Since the **aspect** parameter is 2, we have

$$\text{left} = -\text{right} = 2 * \text{bottom} = -2 * \text{top}.$$

Since the **fov** is 60 degrees, the angle from the center of the image to the edge of the image (along either the x or y axis) is 30 degrees. The **near** plane is at a distance 4. Using $\tan(30) = \frac{1}{\sqrt{3}}$ we get that

$$(\text{left}, \text{right}, \text{bottom}, \text{top}) = \left(-\frac{8}{\sqrt{3}}, \frac{8}{\sqrt{3}}, -\frac{4}{\sqrt{3}}, \frac{4}{\sqrt{3}}\right)$$

4. I showed in the lecture that a projective transformation maps **M** planes to planes. It follows that **M** maps lines to lines. The reason is that any line can be represented as the intersection of two planes. But since two planes π_1 and π_2 map to planes, **M** π_1 and **M** π_2 , the line which is the intersection of π_1 and π_2 must map to the intersection of planes **M** π_1 and **M** π_2 which itself must be a line (since the intersection of two planes is a line).

Note that it follows immediately that triangles map to triangles. I made that claim during the lecture but didn't prove it. So now its proved.

5. No, we can compute the normal instead by taking the cross product

$$\mathbf{n} \equiv \mathbf{M}\mathbf{e}_1 \times \mathbf{M}\mathbf{e}_2$$

for two consecutive edges \mathbf{e}_1 and \mathbf{e}_2 in the polygon. That is, $\mathbf{M}\mathbf{e}_1$ and $\mathbf{M}\mathbf{e}_2$ are direction vectors, representing the directions of transformed edges of the polygon.

6. (a) We can get the matrix **Q** in two ways. One is to use the formula from class (not so interesting and requires less understanding). Expanding the equation, we get

$$x^2 - 6x - 11 + y^2 + 4z^2 = 0,$$

which can be written in the form $\mathbf{x}^T \mathbf{Q} \mathbf{x} = 0$ with:

$$\mathbf{Q} = \begin{bmatrix} 1 & 0 & 0 & -3 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 4 & 0 \\ -3 & 0 & 0 & -11 \end{bmatrix}.$$

A second way to get **Q** is to notice that x is translated by 3 and that otherwise we can write the matrix down by inspection.

$$\mathbf{x}^T \mathbf{Q} \mathbf{x} = \mathbf{x}^T \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ -3 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 4 & 0 \\ 0 & 0 & 0 & -20 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & -3 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \mathbf{x}.$$

- (b) The projective transformation for near and far clipping planes at $z = -1$ and $z = -8$, is:

$$\mathbf{M} = \begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -9 & -8 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

There are two ways we can write out the equation for the transformed ellipsoid.

The first is easier, and closer to the description I gave in Exercises 3 Question 5 (see also slides of the review lecture 13). If \mathbf{x} is a point on the ellipsoid, then $\mathbf{x}' = \mathbf{M}\mathbf{x}$ is a point on the transformed ellipsoid. The points on the original ellipsoid satisfy $\mathbf{x}^T \mathbf{Q} \mathbf{x} = 0$. We write $\mathbf{M}^{-1}\mathbf{x}' = \mathbf{x}$ and substitute, which gives

$$(\mathbf{M}^{-1}\mathbf{x}')^T \mathbf{Q} \mathbf{M}^{-1}\mathbf{x}' = 0$$

or

$$\mathbf{x}'^T \mathbf{M}^{-T} \mathbf{Q} \mathbf{M}^{-1} \mathbf{x}' = 0.$$

The second (which is the one I originally posted) involves a trick. Write $\mathbf{x}^T \mathbf{Q} \mathbf{x} = 0$ by inserting two products of matrices with their inverses:

$$(\mathbf{x}^T \mathbf{M}^T)(\mathbf{M}^{-T} \mathbf{Q} \mathbf{M}^{-1})(\mathbf{M} \mathbf{x}) = 0.$$

Thus, the coefficient matrix of the quadric in the projected view volume is $\mathbf{M}^{-T} \mathbf{Q} \mathbf{M}^{-1}$, i.e. the equation for the transformed ellipsoid is

$$\mathbf{x}'^T \mathbf{M}^{-T} \mathbf{Q} \mathbf{M}^{-1} \mathbf{x}' = 0$$

where $\mathbf{x}' = \mathbf{M}\mathbf{x}$ is the transformed \mathbf{x} . The notation \mathbf{M}^{-T} stands for $(\mathbf{M}^{-1})^T$.