




This page shows the frequency of the top 6 "L2" lineages, across recent months.

The detailed Lineage classifications are provided by Nextclade. I roll those up into "L2" groups, which roughly follow the WHO Variant definitions. For example, my "BA.2.86.*" group includes BA.2.86 and all it's descendants, e.g. the JN.* lineages.

The detailed Lineage classifications are quite numerous and dynamic, so the "Lineage L2" groups give a simpler and more stable basis for analysis and comparison.

The frequency shown at each point is based on the 7-day rolling average across all lineages.


The grey column chart across the bottom shows the volume of sequences available by date. As there can be long sample and data processing times, it is quite routine for recent dates to show lower sample sizes.

The frequency results calculated for the most recent dates might not be representative, due to those lower sample sizes.

n=13,470 sequenced genomes, from 1 March 2026 up to 21 June 2026

Global - BA.3.* vs XFG.*

● $\log (1st \text{ Lineage} / 2nd \text{ Lineage})$ ● trend




This page compares the relative frequency of 2 selected "Lineage L2" groups, over recent months. A challenging Lineage L2 is selected first, and compared to the incumbent.

The trend is shown as a green line and expressed as a daily growth % advantage. If the green line crosses over the 0.0 line, the date when that occurred or is predicted to occur will be shown. At that point the challenging Lineage L2 is considered to have "crossed over" or taken over dominance from the incumbent Lineage L2.


The Lineage classifications are provided by Nextclade. I add the "Lineage L2" groups, typically following common variant groupings, but occasionally being "creative".

The grey column chart across the bottom shows the volume of sequences available by date. As there can be long sample and data processing times, it is quite routine for recent dates to show lower sample sizes.



This page shows the frequency of the top lineages and intensity of samples, across recent months, for Asia.


Location of samples (approx)



n=1,988 sequenced genomes, from 1 March 2026 up to 21 June 2026

Asia - JN.1.* +FLiRT vs NB.1.8.1.* Nimbus


● log (1st Lineage / 2nd Lineage) ● trend



This page shows the growth rate of the top challenger lineage vs the incumbent, for Asia.


Samples by Country


Country	#	Latest Collection date	by Collection date
Singapore	1,288	21/06/2026	
Japan	190	08/06/2026	
South Korea	181	15/06/2026	
Hong Kong	88	19/06/2026	
China	52	26/03/2026	
Malaysia	49	13/05/2026	
Taiwan	43	16/06/2026	
Israel	40	21/06/2026	
Thailand	28	17/06/2026	
Pakistan	14	19/06/2026	
Vietnam	7	13/04/2026	
Kazakhstan	4	15/04/2026	
Armenia	3	24/03/2026	
Cambodia	1	18/05/2026	
Total	1,988	21/06/2026	



This page shows the frequency of the top lineages and intensity of samples, across recent months, for Africa.

Location of samples (approx)






This page shows the growth rate of the top challenger lineage vs the incumbent, for Africa.


Samples by Country


Country	#	Latest Collection date	by Collection date
Ghana	37	20/05/2026	
South Africa	34	27/05/2026	
Reunion	11	15/04/2026	
Uganda	8	09/05/2026	
Cote d'Ivoire	4	30/04/2026	
Zambia	3	09/06/2026	
Total	97	09/06/2026	



This page shows the frequency of the top lineages and intensity of samples, across recent months, for Europe.

Location of samples (approx), excluding Russia







This page shows the growth rate of the top challenger lineage vs the incumbent, for Europe.


Samples by Country

Country	#	Latest Collection date	by Collection date
Spain	280	19/06/2026	
Russia	256	12/06/2026	
Sweden	212	27/05/2026	
France	176	17/06/2026	
Norway	117	28/04/2026	
Italy	99	04/06/2026	
Germany	77	15/06/2026	
Finland	73	21/04/2026	
Netherlands	51	28/04/2026	
Ireland	43	30/05/2026	
Luxembourg	43	03/04/2026	
Estonia	35	23/04/2026	
Poland	34	02/06/2026	
Denmark	30	13/04/2026	
United Kingdom	23	22/04/2026	
Portugal	22	23/04/2026	
Croatia	13	27/04/2026	
Slovenia	10	02/06/2026	
Slovakia	7	30/04/2026	
Czechia	6	13/04/2026	
Ukraine	5	23/03/2026	
Lithuania	4	16/03/2026	
Belgium	3	26/04/2026	
Romania	2	12/05/2026	
Total	1,621	19/06/2026	



This page shows the frequency of the top lineages and intensity of samples, across recent months, for North America






This page shows the growth rate of the top challenger lineage vs the incumbent, for North America.


Samples by Country


Country	#	Latest Collection date	by Collection date
United States	3,474	21/06/2026	
Canada	2,852	19/06/2026	
Costa Rica	46	10/06/2026	
Dominican R...	37	19/06/2026	
Guatemala	29	21/06/2026	
Puerto Rico	23	09/06/2026	
Mexico	2	14/03/2026	
Total	6,463	21/06/2026	



This page shows the frequency of the top lineages and intensity of samples, across recent months, for South America

Location of samples (approx)






This page shows the growth rate of the top challenger lineage vs the incumbent, for South America.


Samples by Country


Country	#	Latest Collection date	by Collection date
Brazil	604	15/06/2026	
Chile	322	20/04/2026	
Ecuador	25	29/05/2026	
French Guiana	18	20/06/2026	
Paraguay	9	20/03/2026	
Argentina	2	30/03/2026	
Colombia	1	15/03/2026	
Total	981	20/06/2026	



This page shows the frequency of the top lineages and intensity of samples, across recent months, for Oceania.

Location of samples (approx)






This page shows the growth rate of the top challenger lineage vs the incumbent, for Oceania.

Samples by Country

Country	#	Latest Collection date	by Collection date
Australia	1,852	21/06/2026	
New Zealand	439	12/06/2026	
New Caledonia	20	13/05/2026	
Guam	9	26/05/2026	
Total	2,320	21/06/2026	



This page shows the frequency of the top 7 lineages, across recent months. The lineages are filtered for a "Lineage L2" group of interest - currently NB.1.8.1.* "Nimbus".

The Lineage classifications are provided by Nextclade. The colour assignments are random.

The frequency shown at each point is based on the 7-day rolling average across all lineages.


The grey column chart across the bottom shows the volume of sequences available by date. As there can be long sample and data processing times, it is quite routine for recent dates to show lower sample sizes.

The frequency results calculated for the most recent dates might not be representative, due to those lower sample sizes.

Global

n=13,470 sequenced genomes, from 1 March 2026 up to 21 June 2026

● RF.5.1 ● RK.1 ● RK.1.2 ● RV.1.3 ● SV.2 ● SV.2.1 ● XFG.1.1




This page shows the frequency of the top 7 lineages, across recent months.

The Lineage classifications are provided by Nextclade. The colour assignments are random.

The frequency shown at each point is based on the 7-day rolling average across all lineages.

The grey column chart across the bottom shows the volume of sequences available by date. As there can be long sample and data processing times, it is quite routine for recent dates to show lower sample sizes.

The frequency results calculated for the most recent dates might not be representative, due to those lower sample sizes.



This page shows the frequency of a selected "Lineage L2" group of interest, for the 7 countries reporting the most samples over recent months.


The detailed Lineage classifications are provided by Nextclade. I roll those up into "L2" groups, which roughly follow the WHO Variant definitions. For example, my "JN.1.* +FLiRT" group includes the descendants of JN.1.* with the mutations: F456L & R346T.

The detailed Lineage classifications are quite numerous and dynamic, so the "Lineage L2" groups give a simpler and more stable basis for analysis and comparison.

The frequency shown at each point is based on the 7-day rolling average across all lineages, for that country.

The grey column chart across the bottom shows the volume of sequences available by date. As there can be long sample and data processing times, it is quite routine for recent dates to show lower sample sizes.

The frequency results calculated for the most recent dates might not be representative, due to those lower sample sizes.



This page shows the frequency of a selected "Lineage L2" group of interest, for the 7 countries reporting the most samples over recent months.


The detailed Lineage classifications are provided by Nextclade. I roll those up into "L2" groups, which roughly follow the WHO Variant definitions. For example, my "JN.1.* +FLiRT" group includes the descendants of JN.1.* with the mutations: F456L & R346T.

The detailed Lineage classifications are quite numerous and dynamic, so the "Lineage L2" groups give a simpler and more stable basis for analysis and comparison.

The frequency shown at each point is based on the 7-day rolling average across all lineages, for that country.

The grey column chart across the bottom shows the volume of sequences available by date. As there can be long sample and data processing times, it is quite routine for recent dates to show lower sample sizes.

The frequency results calculated for the most recent dates might not be representative, due to those lower sample sizes.



This page shows the frequency of a selected Lineage, for the 7 countries reporting the most samples over recent months.

The detailed Lineage classifications are provided by Nextclade.

The frequency shown at each point is based on the 7-day rolling average across all lineages, for that country.


The grey column chart across the bottom shows the volume of sequences available by date. As there can be long sample and data processing times, it is quite routine for recent dates to show lower sample sizes.

The frequency results calculated for the most recent dates might not be representative, due to those lower sample sizes.

n=13,470 sequenced genomes, from 1 March 2026 up to 21 June 2026

Global - XFG.1.1 vs XFG.*

● $\log (1st \text{ Lineage} / 2nd \text{ Lineage})$ ● trend




This page compares the relative frequency of a selected vs a "Lineage L2" group, over recent months. A challenging Lineage is selected first, and compared to the incumbent "Lineage L2" group.

The trend is shown as a green line and expressed as a daily growth % advantage. If the green line crosses over the 0.0 line, the date when that occurred or is predicted to occur will be shown. At that point the challenging Lineage L2 is considered to have "crossed over" or taken over dominance from the incumbent Lineage L2.

The Lineage classifications are provided by Nextclade. I add the "Lineage L2" groups, typically following common variant groupings, but occasionally being "creative".

The grey column chart across the bottom shows the volume of sequences available by date. As there can be long sample and data processing times, it is quite routine for recent dates to show lower sample sizes.


Global



The vertical height of each bar segment represents the relative volume of all the samples of that specific Lineage, as well as all it's descendants.

The full picture is typically quite busy, so insignificant Lineages (with few samples, or at the extreme top or bottom of the hierarchy) are not shown.

The Lineage classifications are provided by Nextclade.



This page shows the frequency of the top 6 "L2" lineages, across recent months, for "International Traveller" samples.

This is probably a more randomised sample than the "Global" aggregate of all samples submitted to GISAID, as those are dominated by the US and Canada

These samples are mainly collected from arrivals into the US and Japan.

Data Submitted in the last 8 weeks

Country	# Samples Sequenced	Latest Collection date	by Collection date	Latest Submission date	by Submission date
Total					

This page shows the volume and currency/timeliness of the genomic sequencing data shared via GISAID, over the last 8 weeks, for the countries sharing the most samples.

Each sample shared comes with a Collection date - when the PCR test for that sample was collected. The GISAID system also records a Submission date for each sample, which is typically the date that sample was uploaded.

The latest date of each type is shown, along with "sparkline"-style mini charts to give a flavour for the spread of recent data by Collection date and by Submission date.