

Probability Formula Review

Free Quick Notes Books has your copies of Quick Notes Accounting, Economics, Statistics, and Excel Statistics Lab Manual.

I. Types and characteristics of probability

A. Types of probability

1. Classical: $P(A) = \frac{A}{N}$
2. Empirical: $P(A) = \frac{A}{n}$
3. Subjective: Use empirical formula assuming past data of similar events is appropriate.

B. Probability characteristics

1. Range for probability: $0 \leq P(A) \leq 1$
2. Value of complements: $P(\bar{A}) = 1 - P(A)$

II. Probability rules

A. Addition is used to find the sum or union of 2 events.

1. General rule: $P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$
2. Special rule: $P(A \text{ or } B) = P(A) + P(B)$ is used when events are mutually exclusive.

B. Multiplication is used to determine joint probability or the intersection of 2 events.

1. General rule: $P(A \text{ and } B) = P(A) \times P(B | A)$
2. Special rule: $P(A \text{ and } B) = P(A) \times P(B)$ is used when the events are independent.

Note: For independent events, the joint probability is the product of the marginal probabilities.

C. Bayes' theorem is used to find conditional probability.

$$P(A|B) = \frac{P(A) \times P(B|A)}{P(A) \times P(B|A) + P(\bar{A}) \times P(B|\bar{A})}$$

Note: The denominator is when condition B happens. It happens with A and with \bar{A} .

III. Counting rules

A. The counting rule of multiple events: If one event can happen M ways and a second event can happen N ways, then the two events can happen (M)(N) ways. For 3 events, use (M)(N)(O).

B. Factorial rule for arranging all of the items of one event: N items can be arranged in N! ways.

C. Permutation rule for arranging some of the items of one event: (order is important: a, b, c and c, a, b are different)

$${}_N P_R = \frac{N!}{(N-R)!}$$

D. Combination rule for choosing some of the items of one event:

(order is not important: abc and cba are the same and are not counted twice)

$${}_N C_R = \frac{N!}{(N-R)!(R)!}$$

IV. Discrete probability distributions

A. Probability distributions

1. $P(x) = [x \cdot P(x)]$ is calculated for each value of x.

2. Mean of a probability distribution: $\mu = E(x) = \sum [x \cdot P(x)]$

3. Variance of a probability distribution: $V(x) = [\sum x^2 \cdot P(x)] - [E(x)]^2$

B. Binomial distributions

$$P(x) = \frac{n!}{x!(n-x)!} p^x q^{n-x} \quad \text{where}$$

n is number of trials	x is number of successes
p is probability of success	q, the probability of failure, is 1 - p
$\mu = np, \sigma^2 = npq \text{ and } \sigma = \sqrt{npq}$	

C. Poisson distributions

$$P(x) = \frac{\mu^x e^{-\mu}}{x!} \quad \text{where } \mu = np$$


Poisson approximation of the binomial requires $n \geq 30$ and $np < 5$ or $nq < 5$.

Business Book Mall has material to enhance your career.

V. The continuous normal probability distribution

- A. To find the probability of x being within a given range: $Z = \frac{x - \mu}{\sigma}$ Normal approximation of the binomial requires $n \geq 30$ and both np and nq are ≥ 5 . The continuity correction factor applies.
- B. To find a range for x given the probability: $\mu \pm Z\sigma$

VI. Central limit theorem



[Excel Internet Library](#) has learning materials classified by type of user.

[MBA Internet Library](#) will help with acceptance, graduation, and career advancement.

[Software Tutorial Internet Library](#) has material to help with many popular software programs.

[Free Non-business Textbooks Library](#) covers many subjects.

VII. Point estimates

- A. \bar{x} for μ B. s for σ C. \bar{p} for p D. $S_{\bar{x}}$ for $\sigma_{\bar{x}}$ where $S_{\bar{x}} = \frac{s}{\sqrt{n}}$ and $\sigma_{\bar{x}} = \frac{\sigma}{\sqrt{n}}$

VIII. Interval estimates when $n \geq 30$

- A. For a population mean $\bar{x} \pm z \frac{\sigma}{\sqrt{n}}$ or $\bar{x} \pm z \frac{s}{\sqrt{n}}$

Note: Use the finite correction factor in section VIII formulas when $n/N \geq .05$. $\sqrt{\frac{N-n}{N-1}}$

- B. For a population proportion $\bar{p} \pm z \sqrt{\frac{\bar{p}(1-\bar{p})}{n}}$ where $\bar{p} = \frac{x}{n}$

Section VIII Note: When $n < 30$ and σ is unknown, the t distribution, to be discussed in chapter 16, must be substituted for the z distribution when making interval estimates. Many statistics software programs do all interval calculations, regardless of sample size, using the t distribution.

IX. Determining sample size

- A. When estimating the population mean $n = \left(\frac{z\sigma}{E} \right)^2$
- B. When estimating the population proportion $n = \bar{p}(1-\bar{p}) \left(\frac{z}{E} \right)^2$