

Yet another results of the spectral and fragmentation study of small meteoroids

V. Vojáček, J. Borovička, P. Koten, P. Spurný, R. Štork

Astronomical institute of the Czech Academy of Sciences

IMC, 2018, Pezinok-Modra, Slovakia



Video observation of meteors

- In Ondřejov since 1990, database of faint meteors
- 2 stations, 2 direct cameras, 1 spectral camera



This work: S-VHS cameras, Mullard image intensifiers, 2004 – 2014

- Orbits, spectra
- fragmentation model




Obtained data

- 152 meteors with orbits, trajectories, spectra
- Spectral classification (Borovička et al., 2005)
 - Mg I-2, Na I-1, Fe I-15
- Application of fragmentation model (Borovička et al., 2007)
- Monochromatic light curves




SX696 - Na free, Halley type, sporadic





Yet another results...

- Study of morphology of faint meteors




- Irons on cometary orbits
- Two phases of erosion of small meteoroids



The morphology

- Low resolution of cameras (200m/px), but still...
- Three categories:
 - No wake

10406060



The morphology

- Low resolution of cameras (200m/px), but still...
- Three categories:
 - Short, faint wake (3 - 15px)


06A20013



The morphology


- Low resolution of cameras (200m/px), but still...
- Three categories:
 - Long, bright wake (15 – 100px)

06B18075



The morphology


- Wake formation?
 - Smaller grains decelerated more than larger grains (Campbell-Brown et al. 2013)
- Na free, Na poor – short faint wakes, no wakes
 - No very small grains
- Irons – no wakes, short faint wakes
 - Small number of grains



The morphology

- Elongated endings
 - Draconids
 - Three sporadic meteors

08728076



The morphology

- Elongated endings
 - Draconids
 - Three sporadic meteors
- Some parameters similar to each other:
- Velocity 25 km/s
- Higher ablation coefficient
- Grain sizes $10^{-2} - 10^{-3}$ mm
- $E_s \sim 10^6$ J/m²

Irons

- 15x irons, 4x modelled
- 10x asteroidal
- 1x Sun – approaching
- 2x Jupiter family
 - 81P/Wild 2: large range of Fe-Mg, Fe-Ni content (Zolensky et al., 2008)
- 2x Halley – inclination $\sim 60^\circ$



Irons


- 2x Halley – inclination $\sim 60^\circ$
 - ??Grand Tack model (Walsh et al. 2011): migration of big planets – planetesimals in Oort cloud

Table 1: Orbital elements of meteoroids classified as Irons. Second row for each meteor contains corresponding errors.

	a	q	Q	i	ω	Ω	v	T_J	orbit
	(AU)	(AU)	(AU)	($^\circ$)	($^\circ$)	($^\circ$)	(km/s)		
SZ2410	3.50	0.654	6.4	63.2	103.2	14.32	40.7	1.9	Halley
	0.09	0.001	0.2	0.2	0.3	-	0.1	0.1	
SX1938	3.6	0.486	7	63	277	143.30	43.4	1.7	Halley
	0.8	0.008	2	1	2	-	0.8	1.0	

Two stages of erosion

- 10% meteors with second brightening




Two stages of erosion

- 10% meteors with second brightening
- Cometary, S-A & Normal, Na poor, Draconids
- Model:
 - Different grain sizes
 - 2nd stage larger grains, less grains
 - Draconids smaller, more grains
 - Ablation coefficient same
 - Erosion coefficient
 - Brightening by Mg = smaller
 - Brightening by Na = larger, same

Two stages of erosion

- Erosion coefficient
 - Brightening by Mg = smaller
 - Brightening by Na = larger, same



Two stages of erosion

- Only cometary and Sun-approaching?
 - Material differences more common
- GAIDA – Rosetta mission:
 - mix of compact and fluffy material (Fulle et al. 2015)
- Solar heating – change in material properties?
(increased density – model Mukai & Fechtig 1983)

Conclusion

- Wake formation: highr probability for meteoroids with larger difference in grain sizes?
- Elongated ending typical for Draconid like meteoroids
- Irons on cometary orbits?
 - We can expect iron rich particles
 - Early solar system mixing of material?
- Two stages of erosion
 - Mix of material with different structure
 - Sometimes caused by brightening of individual lines