

Important declarations

Please remove this info from manuscript text if it is also present there.

Associated Data

Data not supplied by the author for this reason:

The raw data is all in the published literature.

Required Statements

Competing Interest statement:

Mathew J. Wedel is an Academic Editor at PeerJ.

Funding statement:

The research and writing of this work were supported by the svpow.com Patreon subscribers, especially Ben Byer, Gregory Fam, Christopher Heiny, Tyler Holmes, Mad Doctor Hrothgar, William Dale McInnes, Thomas Redd, Mike Traynor, and Humza Yaqoob.

Blood vessels provided the template for vertebral and costal pneumatization in sauropod dinosaurs

Mathew John Wedel ^{Corresp., 1}, Michael P Taylor ²

¹ Department of Anatomy, Western University of Health Sciences, Pomona, California, United States of America

² Department of Earth Sciences, University of Bristol, Bristol, United Kingdom

Corresponding Author: Mathew John Wedel
Email address: mathew.wedel@gmail.com

Pneumatization of the postcranial skeleton, especially the vertebral column, was a key innovation in the evolution of gigantism in sauropod dinosaurs. Despite a century and a half of study, the morphogenetic rules that govern skeletal pneumatization in sauropods — and indeed in extant birds — remain obscure. It has been suggested that pneumatic diverticula of the respiratory system follow pre-existing arteries and veins in pneumatizing the skeleton. This could explain why skeletal pneumaticity is so variable, both in birds and in their extinct outgroups, but published evidence for the “blood vessels first” hypothesis is scant. In extant amniotes, arteries and veins enter and leave the vertebrae and ribs in seven locations: (1) outer surfaces of the neural arch, (2) roof and walls of the neural canal, (3) floor of the neural canal, (4) ventral surfaces of the transverse processes and inner surfaces of the ribs, (5) lateral surfaces of the centrum, (6) ventral surface of the centrum, and (7) medial surfaces of the haemal arch. Vascular foramina are present in all of these locations in apneumatic vertebrae of sauropods, suggesting a highly conserved pattern of vertebral vascularization. Pneumatic fossae and foramina are also present in all of these locations in at least some vertebrae and ribs of sauropods, though not all locations may be pneumatized in a single individual, or even across a clade. The striking similarity of vertebral vascularization and pneumatization in sauropods is consistent with the hypothesis that pneumatic diverticula followed blood vessels, and is difficult to explain any other way. The same relationship of pneumatic features to blood vessels probably existed in non-avian theropods and pterosaurs, but has not yet been documented. The relationships of vascular and pneumatic features in sauropods and other extinct archosaurs could be elucidated by systematic surveys of existing museum collections.

Blood vessels provided the template for vertebral and costal pneumatization in sauropod dinosaurs

Mathew J. Wedel, College of Osteopathic Medicine of the Pacific and College of Podiatric Medicine, Western University of Health Sciences, Pomona, California, USA. mathew.wedel@gmail.com

Michael P. Taylor, Department of Earth Sciences, University of Bristol, Bristol BS8 1RJ, UK. dino@miketaylor.org.uk

Corresponding author: Mathew J. Wedel, mathew.wedel@gmail.com

Abstract

Pneumatization of the postcranial skeleton, especially the vertebral column, was a key innovation in the evolution of gigantism in sauropod dinosaurs. Despite a century and a half of study, the morphogenetic rules that govern skeletal pneumatization in sauropods — and indeed in extant birds — remain obscure. It has been suggested that pneumatic diverticula of the respiratory system follow pre-existing arteries and veins in pneumatizing the skeleton. This could explain why skeletal pneumaticity is so variable, both in birds and in their extinct outgroups, but published evidence for the “blood vessels first” hypothesis is scant. In extant amniotes, arteries and veins enter and leave the vertebrae and ribs in seven locations: (1) outer surfaces of the neural arch, (2) roof and walls of the neural canal, (3) floor of the neural canal, (4) ventral surfaces of the transverse processes and inner surfaces of the ribs, (5) lateral surfaces of the centrum, (6) ventral surface of the centrum, and (7) medial surfaces of the haemal arch. Vascular foramina are present in all of these locations in apneumatic vertebrae of sauropods, suggesting a highly conserved pattern of vertebral vascularization. Pneumatic fossae and foramina are also present in all of these locations in at least some vertebrae and ribs of sauropods, though not all locations may be pneumatized in a single individual, or even across a clade. The striking similarity of vertebral vascularization and pneumatization in sauropods is consistent with the hypothesis that pneumatic diverticula followed blood vessels, and is difficult to explain any other way. The same relationship of pneumatic features to blood vessels probably existed in non-avian theropods and pterosaurs, but has not yet been documented. The relationships of vascular and pneumatic features in sauropods and other extinct archosaurs could be elucidated by systematic surveys of existing museum collections.

Introduction

Pneumatization of the vertebral column was a key innovation in the evolution of gigantism in sauropod dinosaurs (Sander et al. 2010, Sander 2013). Pneumatic features in sauropods vary significantly among taxa (Wedel 2003), among individuals (Wedel and Taylor 2013), within the vertebral column of an individual (Wilson 2012), and even between the left and right sides of a single individual (Wilson 1999, Taylor and Naish 2007, Zurriaguz and Alvarez 2014, Tschopp and Upchurch 2019, Moore et al. 2020, Zurriaguz 2025), and this variation is consistent with that observed in pneumaticity in extant birds (O’Connor 2006). In extant birds, pneumatic diverticula have been reported to follow pre-existing blood vessels in pneumatizing the bones of the postcranial skeleton (Bremer 1940:200). Blood vessels are

highly variable among and within individuals, and to the extent that pneumatic diverticula follow blood vessels, pneumatization will likewise be variable (Taylor and Wedel 2021).

In the present paper we explore the locations where vascular and pneumatic features appear on vertebrae and the reasons why they appear in these particular locations.

Anatomical nomenclature

The term “transverse process” properly refers to processes projecting laterally (or dorsolaterally, posterolaterally, etc.) from the neural arch. The transverse process bears the diapophysis with which the tuberculum of the rib articulates. The term is sometimes also used for the processes that extend laterally from caudal vertebrae. In most caudal vertebrae these processes are actually coalesced costal elements (caudal ribs; Fig. 1). Gallina and Otero (2009) argued that the dorsoventrally tall, plate-like lateral processes of proximal caudal vertebrae in some sauropods consist of a combination of true transverse processes (parts of the neural arches) and fused caudal ribs. Caudal vertebrae otherwise do not have true transverse processes, which are found only in dorsal vertebrae (and in cervicals, although since they are fused to the cervical ribs they are rarely referred to as separate parts of cervical vertebrae). For clarity, we use the term “transverse process” only for the diapophysis-bearing portion of the neural arch, and not for costal elements.

Foramina in the walls of bones that transmit blood vessels have historically been referred to as nutrient foramina, vascular foramina, or neurovascular foramina. Blood vessels are always innervated by microscopic nerve fibers that provide sensory feedback and regulate vasodilation and vasoconstriction (nervi vasorum, see Standring 2008:138), and the living tissue of bones is innervated by additional nerve fibers that accompany blood vessels as they pass inside the bones (Hurrell 1937, Steverink et al. 2021), so technically all vascular foramina are in fact neurovascular. Similarly, all macroscopic nerves are accompanied by microscopic (and sometimes macroscopic) blood vessels (vasa nervorum, see Adams 1942, El-Barrany et al. 1999, and Standring 2008:53), so technically all nervous foramina are also neurovascular. Any given foramen that is said to transmit an artery, vein, or nerve, actually transmits all three, at least at the microscopic level. Gailloud (2021) found that in humans the basivertebral foramina – paired foramina in the floor of the neural canal that admit the basivertebral veins into the dorsal surface of the centrum – also admit small arteries into the centrum along with the veins. Still, it is often the case that vessels that pass into bones are accompanied only by microscopic nerves, or vice versa, so we use the term “vascular foramina” for those whose volume is mostly occupied by blood vessels.

Similarly, O'Connor (2006) found in dissecting birds that had been treated with vascular injections that the pneumatic foramina transmitted blood vessels along with pneumatic diverticula. This probably accounts for the scarcity of purely neurovascular foramina at the macroscopic level in bones that have been completely pneumatized, although neurovascular foramina are sometimes found next to and even inside pneumatic fossae in partially-pneumatized elements (e.g., Taylor and Wedel 2021: figure 6). So in the same sense that vascular and nervous foramina are all technically neurovascular, pneumatic foramina are often and possibly always neurovascular as well. For simplicity and to maintain consistency with existing literature, we refer to foramina that transmit pneumatic diverticula as “pneumatic foramina”, but the fact that such foramina also transmit blood vessels will be important in the discussion below.

Museum Abbreviations

- AMNH — American Museum of Natural History, New York, New York, USA.
- BIBE — Big Bend National Park, held at Perot Museum of Nature and Science, Dallas, Texas, USA.
- BYU — Brigham Young University Museum of Paleontology, Provo, Utah, USA.
- CM — Carnegie Museum of Natural History, Pittsburgh, Pennsylvania, USA.
- FHPR — Utah Field House of Natural History State Park Museum, Vernal, Utah, USA.
- FMNH — Field Museum of Natural History, Chicago, Illinois, USA.
- IVPP — Institute of Vertebrate Paleontology and Paleoanthropology, Beijing, China.
- LACM — Natural History Museum of Los Angeles County, Los Angeles, USA.
- MACN — Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”, Buenos Aires, Argentina.
- MB — Humboldt Museum für Naturkunde Berlin, Berlin, Germany.
- NSMT — National Science Museum, Tokyo, Japan.
- RRBP — Rukwa Rift Basin Project, Tanzanian Antiquities Unit, Dar es Salaam, Tanzania.
- YPM — Yale Peabody Museum, New Haven, Connecticut, USA.

Description

Vertebral articulation

Virtually all tetrapod vertebrae are composed of the same elements: the centrum, neural arch, haemal arch (haemal arch), and paired costal elements (Fig. 1). Not all elements are present in all vertebrae, and the costal elements may be fused (as is generally the case in the neck, sacrum and tail) or mobile (as is generally the case in the trunk), but this is the general plan.

[Figure 1 goes here]

Vertebral vascularization

The vertebral elements are served by a system of arteries and veins. Specific branching patterns are variable among lineages and individuals, but the basic plan is highly conserved. In amniotes, early development of the spine and spinal cord is supported by bilaterally paired intersegmental arteries that branch from the dorsal aorta and pass between adjacent somites (Moran et al. 2012, Moore et al. 2015). After resegmentation, each pair of arteries lies alongside a vertebral centrum, and they become the paired segmental arteries that serve each vertebral segment and its associated muscles and neural tissues (e.g., intercostal arteries and lumbar arteries). The branching of paired segmental arteries from the dorsal aorta produces a ladder-like appearance; this is beautifully illustrated in contrast-enhanced radiographs of newly-hatched chickens (Levinsohn et al. 1984).

To the best of our knowledge, there is no broad, phylogenetically-grounded synthesis of blood supply to the vertebrae in amniotes. Vascular supply to the vertebrae is best understood in mammals (Amato et al. 1959, Crock 1960, Crock et al. 1973, Smuts 1975, Crock and Yoshizawa 1976, Ratcliffe 1982, Konerding and Blank 1987, Gailloud 2019, 2021, and Travan et al. 2015), but aspects of the general pattern described below have been documented in most other amniotes. Where some clades diverge from this pattern, it is generally in the loss of vessels in one or more of these areas. For example, Amato et al. (1959) reported that in rabbits no large vessels entered the external surfaces of the vertebral centra, which were supplied entirely by blood vessels within the neural canal. In contrast, external vascular foramina are common in the centra of many lizards (Camp 1923, Mookerjee and Das 1933). In geckos, vascular foramina are prominent on the ventral centrum but not the lateral surfaces of the

centrum, and foramina are occasionally present on the neural arches as well (Holder 1960). In rare cases, vascular traces on vertebrae allow specific vessels to be reconstructed even in extinct taxa, like the intercostal vessels identified in Mesozoic crocodilians by Gunther (1924).

In birds, vascular foramina are often present on the lateral surfaces of the centra, at least where they have not been overprinted by pneumatization (O'Connor 2006, Taylor and Wedel 2021). The free (unfused) caudal vertebrae and pygostyle bear ventral grooves for the caudal aorta and caudal veins (Rashid et al. 2018, Imai et al. 2019). Blood supply to the spinal cord in birds is broadly similar to that of mammals and other animals, by segmental spinal arteries that branch from the segmental arteries (Roncali and Bosco 1980, Hasouna et al. 1995a, b). In contrast to mammals and most other amniotes, birds lack basivertebral veins, and instead have an expansive internal venous plexus within the neural canal (Burger and Estavillo 1977, O'Connor 2006). Even pneumatic vertebrae of birds may have small vascular foramina within the neural canal (Atterholt et al. 2025).

[Figure 2 goes here]

Figure 2 shows a generalized schema of vertebral vascularization in amniotes, based on the literature cited above. Seven distinct groups of arteries and veins enter or exit the vertebrae at distinctive locations (Table 1): (1) the neural arch; (2) dorsal roof or lateral walls of the neural canal (neural arch elements); (3) ventral floor of the neural canal (centrum); (4) ventral surfaces of the transverse processes, or inner surfaces of the costal elements; (5) lateral surfaces of the centrum; (6) ventral surface of the centrum; (7) medial surface of the haemal arch (Fig. 3).

[Figure 3 goes here]

We assembled this seven-site model of vertebral vascularization based on published accounts, and tested it by documenting vascular foramina in the vertebrae of extant taxa that phylogenetically bracket sauropod dinosaurs: the American alligator, *Alligator mississippiensis*, a pseudosuchian archosaur in which postcranial pneumaticity is primitively absent, and the Emperor penguin, *Aptenodytes forsteri*, a flightless diving bird in which postcranial pneumaticity is secondarily lost. We chose *Alligator* for easy availability of skeletal material, and *Aptenodytes* because of its size (meaning that vascular foramina would be larger and easier to see and photograph) and lack of pneumaticity (meaning that vascular foramina would be identifiable as such, because they had not been overprinted by pneumatization).

[Figure 4 goes here]

We found vascular foramina in all seven of the predicted sites in *Alligator* (Fig. 4), and did not observe prominent vascular foramina in any additional sites. In *Aptenodytes*, we found vascular foramina in the six vertebral sites (Fig. 5). Vascularization of the haemal arches cannot be assessed in birds, because haemal arches were lost in the evolution of birds from other dinosaurs (Rashid et al. 2014). The caudal artery (= caudal aorta) is still present in extinct birds, and it terminates distally by entering the pygostyle through a foramen on the ventral surface (Imai et al. 2019).

[Figure 5 goes here]

We also found vascular foramina at all seven of the predicted sites in the vertebrae of sauropod dinosaurs (Fig. 6). The precaudal vertebrae of most neosauropods are so extensively pneumatized that finding vascular foramina that can be clearly distinguished from pneumatic foramina is challenging. Proximal caudal vertebrae are also pneumatized in many sauropod lineages (Wedel and Taylor 2013, Beeston et al. 2026), so we focused on taxa in which caudal pneumaticity is limited or nonexistent. FHPR 13443, a caudal vertebral centrum from the Upper Jurassic Morrison Formation of Utah that is probably referable to the basal macronarian *Camarasaurus*, bears vascular foramina on the lateral and ventral surfaces of the centrum, and in the floor of the neural canal (dorsal surface of the centrum). The caudal vertebrae of USNM 1556, a partial skeleton of the titanosaur *Alamosaurus* from the Upper Cretaceous North Horn Formation of Utah, bear vascular foramina on the lateral and ventral surfaces of the centra, the lateral aspects of the neural spines, the ventral surfaces of the costal processes, and the medial surfaces of the haemal arches (the neural canals have not been prepped in these vertebrae). Small vascular foramina are visible in the roof of the neural canal in the unfused neural arches of CM 555, a partial skeleton of *Brontosaurus* from the Upper Jurassic Morrison Formation of Wyoming.

[Figure 6 goes here]

Pneumatic sites in sauropod vertebrae

The seven sites where arteries and veins enter and exit the bones are also the sites where the vertebral elements may be pneumatized by diverticula of the respiratory system. In the skeletons of sauropods, pneumatic fossae and foramina have been documented at all seven of these locations (Fig. 7, Table 1), although not all locations may be pneumatized in a single individual, or even across a clade.

[Figure 7 goes here]

The most obvious and best-documented pneumatic traces in sauropod vertebrae are the fossae and foramina on the lateral aspects of the neural arch and centrum (e.g., Wedel 2005:figure 7.2). Also widespread are peduncular fossae: the paired pneumatic fossae or foramina on the anterior or posterior aspect of the neural arch, lateral to the neural canal. These are found in pterosaurs, sauropods, and non-avian theropods, as illustrated and discussed by Taylor and Wedel (2021:figure. 4). These probably also represent pneumatization of the neural arch by diverticula following the segmental spinal arteries that serve the spinal cord.

Pneumatic openings from the neural canal into the neural arch and centrum are less well-documented, in part because the inner surface of the neural canal is difficult to prepare or observe, especially in proportionally long vertebrae. Nevertheless, a few examples of pneumatic fossae and foramina inside the neural canal have been described, both from CT scans (Schwarz and Fritsch 2006:figure 4H–J; Aureliano et al. 2021:figure 2.3) and from direct observation in broken or unfused specimens (Atterholt and Wedel 2018, Gorscak and O'Connor 2019). The centrodiapophyseal fossae of sauropod presacral vertebrae (Wilson et al. 2011) represent pneumatization of the ventral surfaces of the transverse processes and adjacent regions of the neural arches, presumably by diverticula that followed segmental arteries (= intercostal arteries in the dorsal series). The proximal portions of the dorsal ribs are pneumatized in the mamenchisaurid *Xinjiangtitan* (Zhang et al. 2022), a few diplodocids (Lovelace et al. 2008:figure 9, King et al. 2024) and in most titanosauriforms (e.g., Janensch 1950:figure 107, Goman 2005:figure 12, Curry Rogers 2009:figure 30, Taylor and Wedel 2023).

Pneumatization of the ventral surface of the centrum has been documented in multiple regions of the vertebral column. In cervical vertebrae, they are found in the holotype of *Mamenchisaurus sinocanadorum* IVPP V10603 (Moore et al. 2023), the referred *Apatosaurus ajax* specimen NSMT-PV 20375 (Upchurch et al. 2005:plate 1K), the Wyoming *Supersaurus* (Lovelace et al. 2008:figure 4A) and the *Giraffatitan brancai* lectotype MB.R.2180 (personal observation) and paralectotype MB.R.2181 (Janensch 1950:figures 40, 42) among others. We have not observed pneumatic features on ventral surfaces of dorsal or sacral vertebrae, despite having looked for them. In caudal vertebrae, ventral pneumaticity is seen in, for example, *Rocasaurus muniozi* MPCA- Pv 54 (Cerdeña et al. 2012:figure 1G), an indeterminate saltasaurine titanosaur (Zurriaguz et al. 2017:figure 5), and the diplodocoid BYU 11505 (pers. obs.).

Ventral pneumatization may be more common than currently appreciated due to the difficulty in studying the ventral aspects of large, fragile vertebrae that are often stored upright on bases (see, e.g.,

Taylor 2022: figures 5 and 11). This is a rare situation in which mounted sauropod skeletons can be more rather than less scientifically useful than individual elements in collections: they provide an opportunity to view the ventral aspects of large vertebrae, at least in those cases where the ventral aspects are not blocked by supporting armatures.

Finally, pneumatization of the haemal arches has been documented in the indeterminate saltasaurine titanosaur MACN-Pv RN 233-4 (Zurriaguz et al. 2017:figure 9). As is often the case with pneumatic features (Wedel and Taylor 2013) it seems possible that this feature also exists in other known elements, but has not been identified because it was not expected. More careful inspection of other sauropod haemal arches may present a fruitful opportunity to extend our knowledge in this area.

Discussion

Do pneumatic diverticula follow blood vessels?

In describing the process of pneumatization of the humerus in the chicken, *Gallus domesticus*, Bremer (1940: 199) argued that having a pneumatic diverticulum ('air sac' of his usage) in contact with a bone was necessary but not sufficient to lead to internal pneumatization of the bone:

The lower bay of the air sac meets the periosteum, the connective tissue of the sac fuses with it, and the sac flattens against the bone, but does not enter it. Instead, the sac wall becomes wrinkled, the epithelium thickens, and new side pockets burrow between the muscles at some distance from the bone, seeking new regions in which the sac may expand. The apposition of the air sac to the bone cortex is not in itself sufficient to ensure entrance into the bone.

Instead of burrowing directly through the cortical bone, the diverticulum induces bone resorption (through a process of cell signalling that is still not well understood) only in the vicinity of pre-existing blood vessels. As described by Bremer (1940):

The air sac, above the teres major [muscle], meets the bone at the level of the exit of one of the veins, in the region of the newly formed cortex. [...] The inner layer of the periosteum is replaced by a delicate tissue closely resembling mesenchyma. These changes are at first limited to the area immediately surrounding the vein exit (fig. I), but they extend inward chiefly along the vein, with the result that the latter soon lies in a constantly enlarging tubular compartment,

surrounded by bone trabeculae, filled by mesenchyma... Into this loose tissue, along the vein, the air sac finally grows in the form of a long tube. [...] The actual entrance of the air sac into the main marrow cavity is effected at first at the internal opening of the vein.

To our knowledge, this vivid description of the process of pneumatization is still the most detailed available for any postcranial element. (Cranial pneumatization is better studied but not really comparable, since the paranasal and paratympanic diverticula [sensu Witmer 1997] are in intimate contact with the bones of the skull from their earliest stages of development.)

As detailed as Bremer's account is, it is still just one study on the pneumatization of one element — the humerus — in a single species of bird. But other lines of evidence corroborate the hypothesis that pneumatic diverticula follow blood vessels during the process of pneumatizing the postcranial skeleton:

1. Injection studies show that 'pneumatic' foramina are often co-occupied by both pneumatic diverticula and blood vessels (O'Connor 2006:1208-1209).
2. In ducks, *Anas platyrhynchos*, not only do pneumatic fossae on the lateral aspects of the dorsal vertebrae form in places where vascular foramina are already present, but vascular foramina remain visible within the pneumatic fossae (Taylor and Wedel 2021:figure 6).
3. At the macroscopic level, purely vascular foramina are few to nonexistent in well-pneumatized elements. This is certainly true in birds. The cortical bone of sauropod vertebrae is so thick that tiny vascular foramina are often visible on the outer surfaces of pneumatic vertebral centra, especially ventrally and adjacent to the articular surfaces, but obvious vascular foramina are absent on the portions of the centra and especially the neural arches that have been remodeled by pneumaticity. This is consistent with the hypothesis that vascular foramina tend to become overprinted by pneumatic diverticula.

If blood vessels and pneumatic diverticula co-occupy the same foramina, can we say which anatomical system was present first? The vascular network of birds is substantially complete before hatching, including the segmental arteries that supply the vertebrae (Levinsohn et al. 1984). Indeed, bones cannot ossify or be remodeled without substantial blood supply, and vessels come to occupy neurovascular foramina after bone ossifies around them (Seymour et al. 2012, Hendriks and Rasamany 2020). In contrast, pneumatization of the postcranial skeleton doesn't begin until several weeks into post-hatching ontogeny (Hogg 1984). Even absent Bremer's (1940) description of pneumatic diverticula following blood vessels and exploiting existing neurovascular foramina to gain access to the interiors of

postcranial bones, we would predict this based on the facts that (1) blood vessels and pneumatic diverticula co-occupy the same foramina, and (2) blood vessels develop before pneumatic diverticula.

In extant birds, pneumatic foramina can occur on the anterior or posterior surfaces of the proximal femur (Livesey and Zusi 2006). Birds have femoral circumflex arteries that encircle the proximal femur (Baumel 1993), and presumably pneumatic diverticula follow these arteries. It would be interesting to know whether anterior versus posterior pneumatization of the femur correlates with greater blood supply to the femur on its anterior or posterior surfaces; indeed, this would make a near-perfect test of the hypothesis that pneumatic diverticula follow blood vessels.

Although further experimental validation would of course be welcome, the hypothesis that pneumatic diverticula follow blood vessels and that pneumatic foramina are created by the remodeling and expansion of vascular foramina is consistent with all available evidence.

Sites of vascularization are sites of pneumatization

In the present study, we have shown that sauropods manifest pneumatic fossae or foramina in all seven of the sites where blood vessels penetrate vertebrae, corroborating the hypothesis that pneumaticity followed vascularization. As discussed by Taylor and Wedel (2021), this developmental pattern explains why bone — usually the least variable material in the vertebrate body — shows so much variation in its pneumatic morphology: pneumatization follows one of the most developmentally labile systems of the body (see Berger 1956 on the relative variability of bones, blood vessels, and other body systems).

The same relationship of pneumatic features to blood vessels probably existed in other animals with postcranial pneumaticity: non-avian theropods and pterosaurs. The presence of pneumatic features at all seven sites of vascularization has not yet been documented in those clades, but since the known sites of pneumatization are analogous to those documented here for sauropods (e.g. O'Connor 2006, Claessens et al. 2009, Benson et al. 2012), it would not be surprising if pneumatic features were to be found in all sites within both clades.

Pneumatic sites unrelated to major blood vessels

The schema presented here does not explain all known pneumatization in sauropods. For example, according to the seven-sites plan outlined above, pneumatic openings in the ribs should be found only on their posteromedial aspects, corresponding to the location of the intercostal arteries. And indeed in

the great majority of documented pneumatic ribs, this is the case. However, there are exceptions. For example, right dorsal rib 2 of the well-preserved *Apatosaurus louisae* holotype CM 3018 has a pneumatic foramen on its anterolateral surface (Gilmore 1936:plate 24). How did this happen?

As discussed by Witmer (1997:64), pneumatic diverticula can be viewed as “opportunistic pneumatizing machines, resorbing as much bone as possible within the constraints imposed by local biomechanical loading regimes”. As such, pneumatic variability routinely results in unexpected morphology with a great deal of individual variation. In fact diverticula are even more opportunistic than suggested here by Witmer, as they can make their way through soft-tissues as well as bone (Schachner et al. 2021). For example, pelicans and gannets have extensive subcutaneous diverticula (Richardson 1939, 1943, Daoust et al. 2008:figure 3). Soft-tissue diverticula similarly unmoored from major blood vessels occasionally find their way back to bone (O’Connor 2004), enabling pneumatic features to develop even in aspects of a bone where no major blood vessel enters.

Thus the seven-sites plan discussed herein provides a framework that accounts for the great majority of sauropod pneumaticity, but the opportunistic behavior of diverticula means that this schema cannot be exhaustive. It remains possible – and perhaps inevitable – that “out of bounds” pneumatic features will be found in places the model does not predict.

Pneumatic features in unusual neural canals

As described by Wedel and Atterholt (2023), in some aberrant sauropod vertebrae the neural canal is enclosed entirely either by the arch or by the centrum. Neural canals in these positions will still carry the usual segmental spinal arteries, and as they penetrate the walls of the neural canal they will still have the capacity to carry diverticula with them, creating pneumatic foramina in the roof, walls and floor of the canal. The only difference is that different elements of the vertebra would thereby be pneumatized: either all neural-canal pneumatic features penetrate the arch, or they all penetrate the centrum. In other words, the opportunistic behavior of the diverticula remains the same, treating bone as bone without concern for the identity of the bony element that happens to be at hand.

The mundanity of sauropods

Famously, sauropods attained masses an order of magnitude greater than other terrestrial animals, and much speculation has arisen regarding anatomical novelties that may have enabled this gigantism — e.g.

pharyngeal slits for respiration (Gale 1997), multiple accessory hearts in the neck (Choy and Altman 1992), and obligate aquatic habits (as overturned by Kermack 1951, Coombs 1975, Naish 2012 and others). Such ideas are entertaining but have no basis in fossilized evidence. Instead, the conservatism of the sauropod vascular system, as inferred from patterns of vertebral pneumaticity, provides evidence that they were built substantially like other animals, rather than being special magical monsters. Vascular and pneumatic foramina give us a window on the fact that, at the most basic level, sauropod skeletons developed much like those of extant amniotes.

Directions for future research

After roughly a century of only sporadic attention to dinosaurian pneumaticity (e.g., Janensch 1947), work by Witmer (1987, 1990, 1995, 1997), Britt (1993, 1997) and Britt et al. (1998) kicked off a broad resurgence of interest in both cranial and postcranial pneumaticity in dinosaurs and pterosaurs. After the past quarter century of intense work, we now find ourselves in the curious position of knowing far more about pneumaticity in dinosaurs than we do about the underlying vasculature. A systematic survey of vascular foramina in even in a single nearly-complete dinosaur skeleton, like the work done by Smuts (1974, 1975, 1976, 1977a, 1977b) on cattle, would be a welcome advance.

In this paper we have documented that sites of vertebral pneumatization in sauropods correspond with sites of vertebral vascularization in amniotes generally. Further, we have proposed that this correspondence represents causation rather than mere correlation – pneumatic diverticula invaded vertebrae and ribs in predictable locations because they followed pre-existing blood vessels as they developed. This hypothesis can be tested by looking for counter-examples to our seven-sites model. The occasional exception, like pneumatization of the outer surface of a dorsal rib in *Apatosaurus louisae*, can be chalked up to the variability and developmental opportunism of pneumatic diverticula (see Witmer 1997), but a large number of exceptions could force a reevaluation of our hypothesis, as could further experimental work on the actual process of postcranial pneumatization, following the example of Bremer (1940).

Furthermore, the seven-sites model could provide a framework for future investigations comparing the existence or prevalence of vascular and pneumatic traces serially within an individual, ontogenetically within a species or genus, and phylogenetically among lineages. This is an area in which little is known, but rapid progress could potentially be made simply by parsing existing published information using the seven sites as an organizational framework.

Like the collections surveys of Drumheller et al. (2020), McHugh et al. (2020, 2023) and McHugh and Drumheller (2026), all of our proposed investigations — documenting vascular foramina in the skeletons of dinosaurs (and other extinct vertebrates), searching for pneumatic features in places that might contradict the “blood vessels first” hypothesis, and using the seven sites as a framework for investigating serial, ontogenetic, and phylogenetic variation — are inherently collections-based projects that could leverage existing resources and involve students and early-career researchers; and they could be done in parallel, separately or collaboratively, across many subject taxa and many institutions. Even if we are mistaken about the morphogenetic basis of pneumaticity, we hope to inspire useful work.

Acknowledgements

We thank Virginia Zurriaguz for permission to reproduce her photograph of a pneumatic haemal arch in Figure 7G. We are also grateful to all those who allowed us access to specimens in collections, in particular Brooks Britt and Rod Scheetz (Brigham Young University), Matt Lamanna and Amy Henrici (Carnegie Museum of Natural History), Daniela Schwarz (Humboldt Museum für Naturkunde Berlin), Allison Schultz and Young Ha Suh (Natural History Museum of Los Angeles County), Julia McHugh (Museums of Western Colorado — Dinosaur Journey), Ron Tykoski (Perot Museum of Nature and Science), Matt Carrano, Amanda Millhouse, and Matthew Miller (US National Museum of Natural History), and Steve Sroka and John Foster (Utah Field House of Natural History State Park Museum). We thank Andrew Moore and Virginia Zurriaguz for constructive review comments, and handling editor Brandon Hedrick for further helpful suggestions.

References

- Adams, W.E., 1942. The blood supply of nerves: I. Historical review. *Journal of Anatomy*, 76(Pt 4), p.323.
- Amato, V. P., Sliema Malta and R. Bombelli. 1959. The normal vascular supply of the vertebral column in the growing rabbit. *The Journal of Bone & Joint Surgery British Volume* 41B(4):782–795.
doi.org:10.1302/0301-620X.41B4.782
- Atterholt, Jessie, and Mathew J. Wedel. 2018. A CT-based survey of supramedullary diverticula in extant birds. *PeerJ Preprints* 6:e27201v2. doi:10.7287/peerj.preprints.27201v2

Atterholt, Jessie; Burton, M. Grace; Wedel, Mathew J.; Benito, Juan; Fricano, Ellen; and Field, Daniel J. 2025. Osteological correlates of the respiratory and vascular systems in the neural canals of Mesozoic ornithurines *Ichthyornis* and *Janavis*. *The Anatomical Record*. <http://doi.org/10.1002/ar.70070>.

Aureliano, Tito, Aline M. Ghilardi, Bruno A. Navarro, Marcelo A. Fernandes, Fresia Ricardi-Branco and Mathew J. Wedel. 2021. Exquisite air sac histological traces in a hyperpneumatized nanoid sauropod dinosaur from South America. *Scientific Reports* 11(24207). doi:10.1038/s41598-021-03689-8

Baumel, J.J. (editor) 1993. Handbook of avian anatomy: Nomina anatomica avium. Publications of the Nuttall Ornithological Club (USA), no. 23.

Beeston, S.L., Schwarz, D., Upchurch, P., Poropat, S.F., Asbach, P. and Mannion, P.D., 2026. Computed tomography reveals multiple origins of extreme caudal vertebral pneumaticity in sauropod dinosaurs. *Journal of Anatomy*.<https://doi.org/10.1111/joa.70177>

Benson, Roger B. J., Richard J. Butler, Matthew T. Carrano and Patrick M. O'Connor. 2012. Air-filled postcranial bones in theropod dinosaurs: physiological implications and the “reptile”-bird transition. *Biological Reviews* 87:168-193. doi:10.1111/j.1469-185X.2011.00190.x

Bremer, John L. 1940 The pneumatization of the humerus in the common fowl and the associated activity of the elin. *The Anatomical Record* 77(2):197–211. doi:10.1002/ar.1090770209

Britt, B. B. 1993. Pneumatic postcranial bones in dinosaurs and other archosaurs. Ph.D. dissertation, University of Calgary, Calgary, 383 pp.

Britt, Brooks B. 1997. Postcranial pneumaticity. pp. 590–593 in: Philip J. Currie and Kevin Padian (eds). *The Encyclopedia of Dinosaurs*. Academic Press, San Diego.

Britt, Brooks B., Peter J. Makovicky, Jacques Gauthier & Niels Bonde. 1998. Postcranial pneumatization in *Archaeopteryx*. *Nature* 395:374–376.

Burger, R.E. and Estavillo, J.A., 1977. Pulmonary circulation—vertebral venous interconnections in the chicken. *The Anatomical Record*, 188(1), pp.39-43.

Camp, C.L. 1923. Classification of the lizards. *Bulletin of the American Museum of Natural History* 48 : 289-481.

405 Cerda, Ignacio A., Leonardo Salgado and Jaime E. Powell. 2012. Extreme postcranial pneumaticity in
 406 sauropod dinosaurs from South America. *Paläontologische Zeitschrift* 86:441–449. doi:10.1007/s12542-
 407 012-0140

408 Choy, D. S. J., and P. Altmann. 1992. The cardiovascular system of *Barosaurus*: an educated guess. *Lancet*
 409 340(8818):534-536.

410 Claessens, Leon P. A. M., Patrick M. O'Connor, and David M. Unwin. 2009. Respiratory evolution
 411 facilitated the origin of pterosaur flight and aerial gigantism. *PLOS ONE* 4(2): e4497.
 412 doi:10.1371/journal.pone.0004497

413 Coombs Jr, W.P., 1975. Sauropod habits and habitats. *Palaeogeography, Palaeoclimatology,*
 414 *Palaeoecology*, 17(1), pp.1-33.

415 Crock, H. V. 1960. The arterial supply and venous drainage of the vertebral column of the dog. *Journal of*
 416 *Anatomy* 94(1):88–99.

417 Crock, H.V. and Yoshizawa, H., 1976. The blood supply of the lumbar vertebral column. *Clinical*
 418 *Orthopaedics and Related Research* (1976-2007), 115, pp.6-21.

419 Crock HV, Yoshizawa H, Kame SK (1973) Observations on the venous drainage of the human vertebral
 420 body. *Journal of Bone and Joint Surgery*. 55 B, 528-533.

421 Curry Rogers, Kristina. 2009. The postcranial osteology of *Rapetosaurus krausei* (Sauropoda:
 422 Titanosauria) from the Late Cretaceous of Madagascar. *Journal of Vertebrate Paleontology* 29(4):1046-
 423 1086. doi:10.1671/039.029.0432

424 Daoust, P.-Y., G. V. Dobbin, R. C. F. Ridlington Abbott and S. D. Dawson. 2008. Descriptive anatomy of
 425 the subcutaneous air diverticula in the Northern Gannet *Morus bassanus*. *Seabird* 21:64–76.

426 Drumheller, Stephanie K., Julia B. McHugh, Miriam Kane, Anja Riedel and Domenic C. D'Amore. 2020.
 427 High frequencies of theropod bite marks provide evidence for feeding, scavenging, and possible
 428 cannibalism in a stressed Late Jurassic ecosystem. *PLOS ONE* 15(5):e0233115. doi:10.1371/journal.
 429 Pone.0233115

430 El-Barrany, W.G., Marei, A.G. and Vallée, B., 1999. Anatomic basis of vascularised nerve grafts: the blood
 431 supply of peripheral nerves. *Surgical and Radiologic Anatomy*, 21(2), pp.95-102.

432 Fanti, F., Cau, A., Hassine, M., and Contessi, M. 2013. A new sauropod dinosaur from the Early
433 Cretaceous of Tunisia with extreme avian-like pneumatization. *Nature Communications* 4:2080.
434 doi:10.1038/ncomms3080

435 Gailloud, P., 2019. Spinal vascular anatomy. *Neuroimaging Clinics*, 29(4), pp.615-633.

436 Gailloud, P., 2021. Thoracolumbar Spinal Arterial Anatomy, with Special Consideration Given to Spine
437 Intervention. *Image Guided Interventions of the Spine: Principles and Clinical Applications*, pp.35-59.

438 Gale, H. H. 1997. Breathing through a long neck: sauropod lung ventilation. *Journal of Vertebrate*
439 *Paleontology* 17(Suppl. to No. 3):48A.

440 Gallina, P.A. and Otero, A., 2009. Anterior caudal transverse processes in sauropod dinosaurs:
441 morphological, phylogenetic and functional aspects. *Ameghiniana*, 46(1), pp.165-176.

442 Gilmore, Charles W. 1936. Osteology of *Apatosaurus* with special reference to specimens in the Carnegie
443 Museum. *Memoirs of the Carnegie Museum* 11:175–300 and plates XXI–XXXIV.

444 Gomani, Elizabeth M. 2005. Sauropod dinosaurs from the Early Cretaceous of Malawi, Africa.
445 *Palaeontologia Electronica* 8(1):27A (37 pp.)

446 Gorscak, Eric, and Patrick M. O'Connor. 2019, A new African titanosaurian sauropod dinosaur from the
447 middle Cretaceous Galula Formation (Mtuka Member), Rukwa Rift Basin, Southwestern Tanzania. *PLOS*
448 *ONE* 14(2):e0211412. doi:10.1371/journal.pone.0211412

449 Gunther, R.T., 1924. Intercostal blood-vessels in Mesozoic crocodiles. *Proceedings of the Geologists'*
450 *Association*, 35(3), pp.263-264.

451 Hasouna, E.M., A. K. Ahmed, and H. Badawi. 1995a. Deep distribution of spinal arteries in the cervical
452 and lumbosacral enlargements in Pigeon, Duck and Chicken. *Assiut Veterinary Medical Journal* 33(65): 1-
453 10.

454 Hasouna, E.M., Ahmed, A.K. and Badawi, H., 1995b. The arterial vasculature of the lumbo-sacral
455 enlargement of the spinal cord in Duck, Pigeon and Chicken. *Assiut Veterinary Medical Journal* 33(65):
456 11-17.

457 Hendriks, M. and Ramasamy, S.K., 2020. Blood vessels and vascular niches in bone development and
458 physiological remodeling. *Frontiers in Cell and Developmental Biology*, 8, p.602278.

459 Hogg, D.A. 1984b. The development of pneumatisation in the postcranial skeleton of the domestic fowl.
460 *Journal of Anatomy* 139:105-113.

461 Holder, L.A., 1960. The comparative morphology of the axial skeleton in the Australian Gekkonidae.
462 *Zoological Journal of the Linnean Society*, 44(297), pp.300-335.

463 Hurrell, D.J., 1937. The nerve supply of bone. *Journal of Anatomy*, 72, p.54.

464 Imai, T., Azuma, Y., Kawabe, S., Shibata, M., Miyata, K., Wang, M. and Zhou, Z., 2019. An unusual bird
465 (Theropoda, Avialae) from the Early Cretaceous of Japan suggests complex evolutionary history of basal
466 birds. *Communications Biology*, 2(1), p.399.

467 Janensch, W. 1947. Pneumatizität bei Wirbeln von Sauropoden und anderen Saurischien.
468 *Palaeontographica*, Supplement 7, 3:1–25.

469 Janensch, Werner. 1950. Die Wirbelsäule von *Brachiosaurus brancai*. *Palaeontographica* (Suppl. 7) 3:27-
470 93.

471 Kermack, Kenneth A. 1951. A note on the habits of sauropods. *Annals and Magazine of Natural History*,
472 series 12, 4:830–832.

473 King, J.L., McHugh, J.B., Wedel, M.J., and Curtice, B. 2024. A previously unreported form of dorsal rib
474 pneumaticity in *Apatosaurus* (Dinosauria: Sauropoda) and implications for pneumatic variation among
475 diplodocid dorsal ribs. *Journal of Vertebrate Paleontology*. DOI: 10.1080/02724634.2024.2316665

476 Konderding, M.A. and Blank, M., 1987. The vascularization of the vertebral column of rats. *Scanning*
477 *Microscopy*, 1(4), p.23.

478 Levinsohn, E.M., Packard Jr, D.S., West, E.M. and Hootnick, D.R., 1984. Arterial anatomy of chicken
479 embryo and hatchling. *American Journal of Anatomy*, 169(4), pp.377-405.

480 Livesey B, Zusi R (2006) Higher-order phylogeny of modern birds (Theropoda, Aves: Neornithes) based
481 on comparative anatomy. I. Methods and characters. *Bulletin of the Carnegie Museum of Natural*
482 *History* 37:1–544.

483 Lovelace, David M., Scott A. Hartman and William R. Wahl. 2008. Morphology of a specimen of
484 *Supersaurus* (Dinosauria, Sauropoda) from the Morrison Formation of Wyoming, and a re-evaluation of
485 diplodocid phylogeny. *Arquivos do Museu Nacional, Rio de Janeiro*, 65(4):527–544.

486 Marsh, Othniel. C. 1896. *The Dinosaurs of North America*. Extract from the 16th annual report of the U.
487 S. Geological Survey, 1894–95, part I, pp. 133-244 and plates II–LXXXV. doi:10.5962/bhl.title.60562

488 McHugh, J.B. and Drumheller, S.K., 2026. Inter-site and inter-analyst variation in reported frequencies of
489 modified bone material from the upper jurassic morrison formation of Colorado and Wyoming: New
490 Mexico Museum of Natural History and Science Bulletin 102: 375-385.

491 McHugh, Julia B., Stephanie K. Drumheller, Anja Riedel and Miriam Kane. 2020. Decomposition of
492 dinosaurian remains inferred by invertebrate traces on vertebrate bone reveal new insights into Late
493 Jurassic ecology, decay, and climate in western Colorado. *PeerJ* 8:e9510. doi:10.7717/peerj.9510

494 McHugh, J.B., Drumheller, S.K., Kane, M., Riedel, A. and Nestler, J.H., 2023. Assessing paleoecological
495 data retention among disparate field collection regimes: a case study at the Mygatt-Moore Quarry
496 (Morrison Formation). *Palaaios*, 38(5), pp.233-239.

497 Moran, C.M., Myers, C.T., Lewis, C.M. and Krieg, P.A., 2012. Hedgehog regulates angiogenesis of
498 intersegmental vessels through the VEGF signaling pathway. *Developmental Dynamics*, 241(6), pp.1034-
499 1042.

500 Mookerjee, H.K. and Das, G.M., 1933, June. 17. On the peculiar apertures in the vertebral centra of
501 *Typhlops braminus*. *Proceedings of the Zoological Society of London* 103(2): 283-283.

502 Moore, A.J., Upchurch, P., Barrett, P.M., Clark, J.M. and Xing, X., 2020. Osteology of *Klamelisaurus*
503 *gobiensis* (Dinosauria, Eusauropoda) and the evolutionary history of Middle–Late Jurassic Chinese
504 sauropods. *Journal of Systematic Palaeontology*, 18(16), pp.1299-1393.

505 Moore, Andrew J., Paul M. Barrett, Paul Upchurch, Chun-Chi Liao, Yong Ye, Baoqiao Hao & Xing Xu
506 (2023) Re-assessment of the Late Jurassic eusauropod *Mamenchisaurus sinocanadorum* Russell and
507 Zheng, 1993, and the evolution of exceptionally long necks in mamenchisaurids, *Journal of Systematic*
508 *Palaeontology*, 21:1, 2171818, DOI:10.1080/14772019.2023.2171818

509 Moore, Keith L.; Persaud, T. V. N.; Torchia, Mark G. 2015. *The Developing Human: Clinically Oriented*
510 *Embryology*. Elsevier Health Sciences. 288 pp.

511 Naish, Darren. 2012. Palaeontology bites back. *Lab News*, May 2012:31–32.

O'Connor, Patrick M. 2004. Pulmonary pneumaticity in the postcranial skeleton of extant Aves: a case study examining Anseriformes. *Journal of Morphology* 261:141–161.

O'Connor, Patrick M. 2006. Postcranial pneumaticity: an evaluation of soft-tissue influences on the postcranial skeleton and the reconstruction of pulmonary anatomy in archosaurs. *Journal of Morphology* 267(10):1199–1226. doi: 10.1002/jmor.10470

Osborn, Henry Fairfield, and Charles C. Mook. 1921. *Camarasaurus, Amphicoelias* and other sauropods of Cope. *Memoirs of the American Museum of Natural History*, new series, 3:247-387, and plates LX-LXXXV.

Rashid, D.J., Surya, K., Chiappe, L.M., Carroll, N., Garrett, K.L., Varghese, B., Bailleul, A., O'Connor, J.K., Chapman, S.C. and Horner, J.R., 2018. Avian tail ontogeny, pygostyle formation, and interpretation of juvenile Mesozoic specimens. *Scientific reports*, 8(1), p.9014.

Ratcliffe, J. F. 1982. An evaluation of the intra-osseous arterial anastomoses in the human vertebral body at different ages. A microarteriographic study. *Journal of Anatomy* 134(2):373–382.

Richardson, F., 1939. Functional aspects of the pneumatic system of the California brown pelican. *The Condor*, 41(1), pp. 13-17.

Richardson, F., 1943. Pneumaticity of the white pelican. *The Condor*, 45(5), pp. 37-38.

Riggs, Elmer S. 1904. Structure and relationships of opisthocoelian dinosaurs. Part II, the Brachiosauridae. *Field Columbian Museum, Geological Series* 2(6):229–247, plus plates LXXI–LXXV.

Roncali, L. and Bosco, L., 1980. Relationships between vasculogenesis and morphogenesis in the chick embryo spinal ganglia. *Cells Tissues Organs*, 106(4), pp.473-486.

Sander, P. Martin. 2013. An evolutionary cascade model for sauropod dinosaur gigantism – overview, update and tests. *PLOS ONE* 8(10):e78573. doi:10.1371/journal.pone.0078573

Sander, P. Martin, Andreas Christian, Marcus Clauss, Regina Fechner, Carole T. Gee, Eva-Maria Griebeler, Hanns-Christian Gunga, Jürgen Hummel, Heinrich Mallison, Steven F. Perry, Holger Preuschoft, Oliver W. M. Rauhut, Kristian Remes, Thomas Tütken, Oliver Wings and Ulrich Witzel. 2010. Biology of the sauropod dinosaurs: the evolution of gigantism. *Biological Reviews* 86:117–155. doi:10.1111/j.1469-185X.2010.00137.x

Schachner, E.R., Hedrick, B.P., Richbourg, H.A., Hutchinson, J.R. and Farmer, C.G., 2021. Anatomy, ontogeny, and evolution of the archosaurian respiratory system: A case study on Alligator mississippiensis and Struthio camelus. *Journal of Anatomy*, 238(4), pp.845-873.

Schwarz, Daniela, and Guido Fritsch. 2006. Pneumatic structures in the cervical vertebrae of the Late Jurassic Tendaguru sauropods *Brachiosaurus brancai* and *Dicraeosaurus*. *Eclogae Geologicae Helvetiae* 99 (1):65–78. doi:10.1007/s00015-006-1177-x

Seymour, Roger S., Sarah L. Smith, Craig R. White, Donald M. Henderson, Daniela Schwarz-Wings; Blood flow to long bones indicates activity metabolism in mammals, reptiles and dinosaurs. *Proceedings of the Royal Society B* 279 (1728): 451–456.

Smuts, Malie. M. S. 1974. The foramina of the cervical vertebrae of the ox. Part I: Atlas and Axis. *Anatomia, Histologia, Embryologia* 3:296–307.

Smuts, Malie M. S. 1975. The foramina of the cervical vertebrae of the ox, part II: cervical vertebrae 3–7. *Anatomia, Histologia, Embryologia* 4(1):24–37.

Smuts, Malie. M. S. 1976. Areas of muscular attachment and their correlation with foraminous areas of the cervical vertebrae of the ox (*Bos taurus* L.). *Anatomia, Histologia, Embryologia* 5:253–266.

Smuts, Malie M. S. 1977a. The arterial blood supply of the cervical vertebrae of the ox (*Bos taurus* L.). *Onderstepoort Journal of Veterinary Research* 44(2):95–106.

Smuts, Malie M. S. 1977b. The venous drainage of the cervical vertebrae of the ox (*Bos taurus* L.). *Onderstepoort Journal of Veterinary Research* 44(4):233–248.

Standring, S. (editor). 2008. *Gray's Anatomy*, 40th edition. 2008. Churchill Livingstone Elsevier, London.

Steverink, J.G., Oostinga, D., van Tol, F.R., van Rijen, M.H., Mackaaij, C., Verlinde-Schellekens, S.A., Oosterman, B.J., Van Wijck, A.J., Roeling, T.A. and Verlaan, J.J., 2021. Sensory innervation of human bone: an immunohistochemical study to further understand bone pain. *The Journal of Pain*, 22(11), pp.1385-1395.

Taylor MP. 2022. Almost all known sauropod necks are incomplete and distorted. *PeerJ* 10:e12810 <https://doi.org/10.7717/peerj.12810>

565 Taylor, Michael P., and Darren Naish. 2007. An unusual new neosauropod dinosaur from the Lower
566 Cretaceous Hastings Beds Group of East Sussex, England. *Palaeontology* 50(6):1547–1564. doi:
567 10.1111/j.1475-4983.2007.00728.x

568 Taylor, Michael P., and Mathew J. Wedel. 2021. Why is vertebral pneumaticity in sauropod dinosaurs so
569 variable? (version 5) *Qeios* 1G6J3Q.5. doi: 10.32388/1G6J3Q.5 <https://www.qeios.com/read/1G6J3Q.5>

570 Taylor, Michael P., and Matthew J. Wedel. 2023. Novel pneumatic features in the ribs of the sauropod
571 dinosaur *Brachiosaurus altithorax*. *Acta Palaeontologica Polonica* 68(4): 709–718.
572 doi:10.4202/app.01105.2023

573 Travan, Luciana, Paola Saccheri, Giorgia Gregoraci, Chiara Mardegan and Enrico Crivellato. 2015. Normal
574 anatomy and anatomic variants of vascular foramina in the cervical vertebrae: a paleo-osteological
575 study and review of the literature. *Anatomical Science International* 90:308–323. doi:10.1007/s12565-
576 014-0270-x

577 Upchurch, Paul, Yukimitsu Tomida, and Paul M. Barrett. 2005. A new specimen of *Apatosaurus ajax*
578 (Sauropoda: Diplodocidae) from the Morrison Formation (Upper Jurassic) of Wyoming, USA. *National*
579 *Science Museum Monographs* 26. Tokyo. ISSN 1342-9574.

580 Wedel, Mathew J. 2003. The evolution of vertebral pneumaticity in sauropod dinosaurs. *Journal of*
581 *Vertebrate Paleontology* 23(2):344–357. doi:10.1671/0272-
582 4634(2003)023%5B0344:TEOVPI%5D2.0.CO;2

583 Wedel, Mathew J. 2005. Postcranial skeletal pneumaticity in sauropods and its implications for mass
584 estimates. pp. 201–228 in Wilson, J. A., and Curry Rogers, K. (eds.), *The Sauropods: Evolution and*
585 *Paleobiology*. University of California Press, Berkeley.

586 Wedel, Mathew J., and Jessie Atterholt. 2023. Expanded neurocentral joints in the vertebrae of
587 sauropod dinosaurs. pp. 256–257 in: Hunt-Foster, Rebecca K., James I. Kirkland and Mark A. Loewen
588 (eds), *14th Symposium on Mesozoic Terrestrial Ecosystems and Biota. The Anatomical Record* 306(S1).

589 Wedel, Mathew J., and Michael P. Taylor 2013. Caudal pneumaticity and pneumatic hiatuses in the
590 sauropod dinosaurs *Giraffatitan* and *Apatosaurus*. *PLOS ONE* 8(10):e78213. 14 pages. doi:
591 10.1371/journal.pone.0078213

592 Wilson, J. A. 1999. A nomenclature for vertebral laminae in sauropods and other saurischian dinosaurs.
593 *Journal of Vertebrate Paleontology* 19, 639-653.

594 Willson, J.A., 2012. New vertebral laminae and patterns of serial variation in vertebral laminae of
595 sauropod dinosaurs. *Contributions from the Museum of Paleontology, University of Michigan*, 32(7),
596 pp.91-110.

597 Wilson, Jeffrey A, Michael D. D'Emic, Takehito Ikejiri, Emile M. Moacdieh and John A. Whitlock. 2011. A
598 Nomenclature for vertebral fossae in sauropods and other saurischian dinosaurs. *PLOS ONE* 6(2):e17114.
599 doi:10.1371/journal.pone.0017114

600 Witmer, Lawrence M. 1987. The nature of the antorbital fossa of archosaurs: shifting the null
601 hypothesis. pp. 230–235 in: Philip J. Currie and E. H. Koster (eds). *Fourth Symposium on Mesozoic*
602 *Terrestrial Ecosystems, Short Papers*. Occasional Paper of the Tyrrell Museum of Palaeontology #3,
603 Drumheller, Alberta.

604 Witmer, L.M., 1990. The craniofacial air sac system of Mesozoic birds (Aves). *Zoological Journal of the*
605 *Linnean Society*, 100(4), pp.327-378.

606 Witmer, L.M., 1995. Homology of facial structures in extant archosaurs (birds and crocodilians), with
607 special reference to paranasal pneumaticity and nasal conchae. *Journal of morphology*, 225(3), pp.269-
608 327.

609 Witmer, Lawrence M. 1997. The evolution of the antorbital cavity of archosaurs: a study in soft-tissue
610 reconstruction in the fossil record with an analysis of the function of pneumaticity. *Journal of Vertebrate*
611 *Paleontology* 17(S1):1–76. doi:10.1080/02724634.1997.10011027

612 Zhang, X.-Q., Li, N., Xie, Y., Li, D.-Q., & You, H.-L. (2022). Redescription of the dorsal vertebrae of the
613 mamenchisaurid sauropod *Xinjiangtitan shanshanensis* Wu et al. 2013. *Historical Biology*, 1–27.
614 <https://doi.org/10.1080/08912963.2022.2147428>

615 Zurriaguz, V.L., 2025. Variation in the postcranial pneumaticity in derived titanosaurs (Dinosauria:
616 Sauropoda). *Historical Biology*, 37(6), pp.1465-1479.

617 Zurriaguz, Virginia L., and A. Álvarez. 2014. Shape variation in presacral vertebrae of saltasaurine
618 titanosaurs (Dinosauria, Sauropoda). *Historical Biology* 26(6):801–809.
619 doi:10.1080/08912963.2013.858248

Zurriaguz, Virginia, Agustín Martinelli, Guillermo W. Rougier and Martín D. Ezcurra. 2017. A saltasaurine titanosaur (Sauropoda: Titanosauriformes) from the Angostura Colorada Formation (upper Campanian, Cretaceous) of northwestern Patagonia, Argentina. *Cretaceous Research* 75:101–114. doi:10.1016/j.cretres.2017.03.016

Figure Captions

Figure 1. Two schematic sauropod vertebrae, showing the homologous ossifications that compose them. **Left:** dorsal vertebra and free ribs. **Right:** caudal vertebra including fused ribs. Note that there are no haemal arches in precaudal (cervical, dorsal and sacral) vertebrae. Image composed from dorsal vertebra 4 of *Camarasaurus supremus* AMNH 5760'/D-X-131 in anterior view, modified from Osborn and Mook 1921:plate 70; left rib 4 (mirrored) of *Camarasaurus supremus* AMNH 5761/R-A-24 in anterior view, modified from Osborn and Mook 1921:figure 71; caudal vertebra 2 of *Camarasaurus lentus* (YPM collection, specimen number unknown) in anterior view, modified from Marsh 1896:plate 34, part 4; and haemal arch of *Camarasaurus grandis* (YPM collection, specimen number unknown) in posterior view, modified from Marsh 1896:plate 39: part 3c.

Figure 2. The major arteries and veins that serve each vertebral segment in amniotes. These vessels are largely conserved across Amniota; when a clade diverges from this pattern, it is usually through the loss of specific vessels, such as equatorial arteries in rabbits (Amato et al. 1959) and basivertebral veins in birds (O'Connor 2006).

Figure 3. The seven sites where arteries and veins enter and exit the bones (see Table 1).

Figure 4. Examples of vascular foramina in vertebrae of the American alligator, *Alligator mississippiensis*. **A.** The seven-site model from Fig. 3. **B.** Vascular foramina in the roof of the neural canal, going into the neural arch. **C1.** A vascular foramen on the lateral aspect of the neural arch. **C2.** A vascular foramen on the lateral aspect of the centrum. **D.** Vascular foramina in the floor of the neural canal, going into the centrum. **E.** A vascular foramen on the underside of the transverse process. **F.** A vascular foramen on the ventral aspect of the centrum. **G.** A vascular foramen on the medial aspect of the haemal arch. Specimens kindly provided by Andrew Farke.

Figure 5. Examples of vascular foramina in vertebrae of the Emperor penguin, *Aptenodytes forsteri*, LACM (Ornithology) 99854. **A.** The seven-site model from Fig. 3. **B.** Vascular foramina in the roof of the

neural canal, going into the neural arch. **C1.** Vascular foramina on the lateral aspect of the neural arch. **C2.** A vascular foramen on the lateral aspect of the centrum. **D.** Vascular foramina in the floor of the neural canal, going into the centrum. **E.** Vascular foramina on the underside of the transverse process. **F.** Vascular foramina on the ventral aspect of the centrum.

Figure 6. Examples of vascular foramina in the vertebrae of sauropods. **A.** The seven-site model from Fig. 3. **B.** Vascular foramina in the roof of the neural canal in an unfused cervical neural arch of *Brontosaurus* CM 555. **C1.** Vascular foramen on the lateral aspect of the neural arch in a proximal caudal of *Alamosaurus* USNM 15556. **C2.** A large vascular foramen on the lateral aspect of the centrum in a proximal caudal of *Alamosaurus* USNM 15556. **D.** Vascular foramina in the floor of the neural canal, going into the centrum in a caudal vertebra of cf. *Camarasaurus* FHPR 13443. **E.** A vascular foramen on the underside of the transverse process in a proximal caudal of *Alamosaurus* USNM 15556. **F.** A vascular foramen on the ventral aspect of the centrum in a caudal vertebra of cf. *Camarasaurus* FHPR 13443. **G.** Vascular foramina on the medial surface of a haemal arch of *Alamosaurus* USNM 15556.

Figure 7. Examples of pneumatic structures in sauropod vertebrae matching each of the seven sites of vascularization. **A.** The seven-site model from Fig. 3. **B.** Cervical neural arch of the titanosaur *Alamosaurus sanjuanensis* unnumbered BIBE specimen in ventral view (anterior to right), showing pneumatic fossae in the roof of the neural canal. Modified from slide 46 in supplementary information to Atterholt and Wedel (2018). **C.** Cervical vertebra 8 of the brachiosaurid *Giraffatitan brancai* paralectotype MB.R.2181, in left lateral view, showing extensive pneumatic sculpting in the neural spine and a deep pneumatic fossa in the side of the centrum. Photograph by authors. **D.** Dorsal vertebral centrum of the titanosaur *Mnyamawamtuka moyowamkia* holotype RRBP 05834 in dorsal view, anterior to top, showing pneumatic fossa in the floor of the neural canal. Modified from Gorscak and O'Connor (2019:figure 5C). **E.** Dorsal rib of the brachiosaurid *Brachiosaurus altithorax* holotype FMNH PR 25107, showing pneumatic foramen in the medial aspect. Modified from Riggs (1904:plate 75:figure 5). **F.** Cervical vertebra 6 of the brachiosaurid *Giraffatitan brancai* paralectotype MB.R.2181, in ventral view with anterior to the top, showing a distinct pneumatic foramen on the bottom of the centrum. Photograph by authors. **G.** haemal arch of indeterminate saltasaurine titanosaur MACN-Pv RN 233-4 in left posterolateral view, showing pneumatic foramen on medial aspect of right portion. Photograph courtesy of Virginia Zurriaguz.

Figure 1

Two schematic sauropod vertebrae, showing the homologous ossifications that compose them.

Left: dorsal vertebra and free ribs. Right: caudal vertebra including fused ribs. Note that there are no haemal arches in precaudal (cervical, dorsal and sacral) vertebrae. Image composed from dorsal vertebra 4 of *Camarasaurus supremus* AMNH 5760'/D-X-131 in anterior view, modified from Osborn and Mook 1921:plate 70; left rib 4 (mirrored) of *Camarasaurus supremus* AMNH 5761/R-A-24 in anterior view, modified from Osborn and Mook 1921:figure 71; caudal vertebra 2 of *Camarasaurus lentus* (YPM collection, specimen number unknown) in anterior view, modified from Marsh 1896:plate 34, part 4; and haemal arch of *Camarasaurus grandis* (YPM collection, specimen number unknown) in posterior view, modified from Marsh 1896:plate 39: part 3c.





Figure 2

The major arteries and veins that serve each vertebral segment in amniotes.

These vessels are largely conserved across Amniota; when a clade diverges from this pattern, it is usually through the loss of specific vessels, such as equatorial arteries in rabbits (Amato et al. 1959) and basivertebral veins in birds (O'Connor 2006).




Figure 3

The seven sites where arteries and veins enter and exit the bones (see Table 1).

In life, most bones are surrounded by a dense vascular network, and microscopic vessels may enter and leave bones at many points. Only the most important vascularization sites are shown here.




Figure 4

Examples of vascular foramina in vertebrae of the American alligator, *Alligator mississippiensis*.

(A) The seven-site model from Fig. 3. (B) Vascular foramina in the roof of the neural canal, going into the neural arch. (C1) A vascular foramen on the lateral aspect of the neural arch. (C2) A vascular foramen on the lateral aspect of the centrum. (D) Vascular foramina in the floor of the neural canal, going into the centrum. (E) A vascular foramen on the underside of the transverse process. (F) A vascular foramen on the ventral aspect of the centrum. (G) A vascular foramen on the medial aspect of the haemal arch. Specimens kindly provided by Andrew Farke.




Figure 5

Examples of vascular foramina in vertebrae of the Emperor penguin, *Aptenodytes forsteri*, LACM (Ornithology) 99854.

(A) The seven-site model from Fig. 3. (B) Vascular foramina in the roof of the neural canal, going into the neural arch. (C1) Vascular foramina on the lateral aspect of the neural arch. (C2) A vascular foramen on the lateral aspect of the centrum. (D) Vascular foramina in the floor of the neural canal, going into the centrum. (E) Vascular foramina on the underside of the transverse process. (F) Vascular foramina on the ventral aspect of the centrum.




Figure 6

Examples of vascular foramina in the vertebrae of sauropod dinosaurs.

(A) The seven-site model from Fig. 3. (B) Vascular foramina in the roof of the neural canal in an unfused cervical neural arch of *Brontosaurus* CM 555. (C1) Vascular foramen on the lateral aspect of the neural arch in a proximal caudal of *Alamosaurus* USNM 15556. (C2) A large vascular foramen on the lateral aspect of the centrum in a proximal caudal of *Alamosaurus* USNM 15556. (D) Vascular foramina in the floor of the neural canal, going into the centrum in a caudal vertebra of cf. *Camarasaurus* FHPR 13443. (E) A vascular foramen on the underside of the transverse process in a proximal caudal of *Alamosaurus* USNM 15556. (F) A vascular foramen on the ventral aspect of the centrum in a caudal vertebra of cf. *Camarasaurus* FHPR 13443. (G) Vascular foramina on the medial surface of a haemal arch of *Alamosaurus* USNM 15556.




Figure 7

Examples of pneumatic structures in sauropod vertebrae matching each of the seven sites of vascularization.

(A) The seven-site model from Fig. 3. (B) Cervical neural arch of the titanosaur *Alamosaurus sanjuanensis* unnumbered BIBE specimen in ventral view (anterior to right), showing pneumatic fossae in the roof of the neural canal. Modified from slide 46 in supplementary information to Atterholt and Wedel (2018). (C) Cervical vertebra 8 of the brachiosaurid *Giraffatitan brancai* paralectotype MB.R.2181, in left lateral view, showing extensive pneumatic sculpting in the neural spine and a deep pneumatic fossa in the side of the centrum. Photograph by authors. (D) Dorsal vertebral centrum of the titanosaur *Mnyamawamtuka moyowamkia* holotype RRBP 05834 in dorsal view, anterior to top, showing pneumatic fossa in the floor of the neural canal. Modified from Gorscak and O'Connor (2019:figure 5C). (E) Dorsal rib of the brachiosaurid *Brachiosaurus altithorax* holotype FMNH PR 25107, showing pneumatic foramen in the medial aspect. Modified from Riggs (1904:plate 75:figure 5). (F) Cervical vertebra 6 of the brachiosaurid *Giraffatitan brancai* paralectotype MB.R.2181, in ventral view with anterior to the top, showing a distinct pneumatic foramen on the bottom of the centrum. Photograph by authors. (G) haemal arch of indeterminate saltasaurine titanosaur MACN-Pv RN 233-4 in left posterolateral view, showing pneumatic foramen on medial aspect of right portion. Photograph courtesy of Virginia Zurriaguz.

