

# The Characteristic Equation

**Geometric Algorithms**

**Lecture 18**

# Introduction

# Recap Problem

$$\begin{bmatrix} 5 & 2 & 3 & 0 \\ -1 & 2 & -3 & 1 \\ 2 & 4 & 10 & 0 \\ 1 & 2 & 3 & 5 \end{bmatrix}$$

*Determine the dimension of the eigenspace of  $A$  for the eigenvalue 4.*

*(try not to do any row reductions)*

**Answer: 2**

$$\begin{bmatrix} 5 & 2 & 3 & 0 \\ -1 & 2 & -3 & 1 \\ 2 & 4 & 10 & 0 \\ 1 & 2 & 3 & 5 \end{bmatrix}$$

# Objectives

1. Briefly recap eigenvalues and eigenvectors.
2. Get a primer on determinants.
3. Determine how to find eigenvalues (not just verify them).

# Keyword

eigenvectors

eigenvalues

eigenspaces

eigenbases

determinant

characteristic equation


polynomial roots

triangular matrices

multiplicity

# Recap

# Recall: Eigenvalues/vectors




A *nonzero* vector  $\mathbf{v}$  in  $\mathbb{R}^n$  and real number  $\lambda$  are an **eigenvector** and **eigenvalue** for a  $n \times n$  matrix  $A$  if

$$A\mathbf{v} = \lambda\mathbf{v}$$




# Recall: Eigenvalues/vectors



A *nonzero* vector  $\mathbf{v}$  in  $\mathbb{R}^n$  and real number  $\lambda$  are an **eigenvector** and **eigenvalue** for a  $n \times n$  matrix  $A$  if

$$A\mathbf{v} = \lambda\mathbf{v}$$

# Recall: Eigenvalues/vectors




A *nonzero* vector  $\mathbf{v}$  in  $\mathbb{R}^n$  and real number  $\lambda$  are an **eigenvector** and **eigenvalue** for a  $n \times n$  matrix  $A$  if

$$A\mathbf{v} = \lambda\mathbf{v}$$

$\mathbf{v}$  is "just scaled" by  $A$ , not rotated

# Recall: The Picture



# Recall: Verifying Eigenvectors

# Recall: Verifying Eigenvectors

**Question.** Determine if  $\mathbf{v}$  is an eigenvector of  $A$  and determine the corresponding eigenvalues.

# Recall: Verifying Eigenvectors

**Question.** Determine if  $\mathbf{v}$  is an eigenvector of  $A$  and determine the corresponding eigenvalues.

**Solution.** Easy. Work out the matrix–vector multiplication.

# Recall: Verifying Eigenvectors

**Question.** Determine if  $\mathbf{v}$  is an eigenvector of  $A$  and determine the corresponding eigenvalues.

**Solution.** Easy. Work out the matrix–vector multiplication.

Example.

$$\begin{bmatrix} 1 & 6 \\ 5 & 2 \end{bmatrix} \begin{bmatrix} 6 \\ -5 \end{bmatrix} = \begin{bmatrix} -24 \\ 20 \end{bmatrix} = -4 \begin{bmatrix} 6 \\ -5 \end{bmatrix} \quad \checkmark$$

$$\begin{bmatrix} 1 & 6 \\ 5 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 13 \\ 9 \end{bmatrix} \quad \times$$

# Recall: Verifying Eigenvalues



# Recall: Verifying Eigenvalues

**Question.** Find an eigenvector of  $A$  whose corresponding eigenvalue is  $\lambda$ .

# Recall: Verifying Eigenvalues

**Question.** Find an eigenvector of  $A$  whose corresponding eigenvalue is  $\lambda$ .

**Solution.** Find a nontrivial solution to

$$(A - \lambda I)\mathbf{x} = \mathbf{0}$$

# Recall: Verifying Eigenvalues

**Question.** Find an eigenvector of  $A$  whose corresponding eigenvalue is  $\lambda$ .

**Solution.** Find a nontrivial solution to

$$(A - \lambda I)\mathbf{x} = \mathbf{0}$$

*If we don't need the vector we can just show that  $A - \lambda I$  is **not** invertible (by IMT).*

# Recall: Finding Eigenspaces

# Recall: Finding Eigenspaces

**Question.** Find a basis for the eigenspace of  $A$  corresponding to  $\lambda$ .

# Recall: Finding Eigenspaces

**Question.** Find a basis for the eigenspace of  $A$  corresponding to  $\lambda$ .

**Solution.** Find a basis for  $\text{Nul}(A - \lambda I)$ .

# Recall: Finding Eigenspaces

**Question.** Find a basis for the eigenspace of  $A$  corresponding to  $\lambda$ .

**Solution.** Find a basis for  $\text{Nul}(A - \lambda I)$ .

(we did this for our recap problem)

# Finding Eigenvalues



# Finding Eigenvalues

**Question.** Determine the eigenvalues of  $A$ , along with their associated eigenspaces.

# Finding Eigenvalues

**Question.** Determine the eigenvalues of  $A$ , along with their associated eigenspaces.

**Solution (Idea).** Can we somehow "solve for  $\lambda$ " in the equation

$$(A - \lambda I)\mathbf{x} = \mathbf{0}$$

# Determinants

# An Aside: Determinants are Mysterious


Determinants are  
strangely polarizing

Some people love them,  
some people hate them

We'll only scratch the  
surface...

Down with Determinants!

Sheldon Axler



102 (1995), 139-154.

ry writing from the Mathematical Association of America.

without determinants. The standard proof that a square matrix of complex numbers has an eigenvalue uses d  
eterminants, this allows us to define the multiplicity of an eigenvalue and to prove that the number of eigenval  
characteristic and minimal polynomials and then prove that they behave as expected. This leads to an easy p  
determinants, this paper gives a simple proof of the finite-dimensional spectral theorem.

n this paper. The book is intended to be a text for a second course in linear algebra.

# **What kind of thing is the determinant?**

# What kind of thing is the determinant?

A determinant is a number associated with a matrix.

# What kind of thing is the determinant?

A determinant is a number associated with a matrix.

**Notation.** We will write  $\det(A)$  for the determinant of  $A$ .

# What kind of thing is the determinant?

A determinant is a number associated with a matrix.

**Notation.** We will write  $\det(A)$  for the determinant of  $A$ .

In broad strokes, it's a big sum of products of entries of  $A$ .



# A Scary-Looking Definition (we won't use)

$$\det(A) = \sum_{\sigma \in S_n} (-1)^{\text{sgn}(\sigma)} A_{1\sigma(1)} A_{2\sigma(2)} \cdots A_{n\sigma(n)}$$

We can think of this function as a procedure:

# A Scary-Looking Definition (we won't use)

$$\det(A) = \sum_{\sigma \in S_n} (-1)^{\text{sgn}(\sigma)} A_{1\sigma(1)} A_{2\sigma(2)} \cdots A_{n\sigma(n)}$$

We can think of this function as a procedure:

```
1 FUNCTION det(A):  
2   total = 0  
3   FOR all matrix B we can get by swapping a bunch of rows of A:  
4     s = 1 IF (# of swaps necessary) is even ELSE -1  
5     total += s * (product of the diagonal entries of B)  
6   RETURN total
```

# The Determinant of $2 \times 2$ Matrices

$$\det \begin{bmatrix} a & b \\ c & d \end{bmatrix} = ad - bc$$

# The Determinant of $2 \times 2$ Matrices

$$\det \begin{bmatrix} a & b \\ c & d \end{bmatrix} = ad - bc$$

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \rightarrow^0 \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

$$(-1)^0 ad$$

# The Determinant of $2 \times 2$ Matrices

$$\det \begin{bmatrix} a & b \\ c & d \end{bmatrix} = ad - bc$$

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \rightarrow^1 \begin{bmatrix} c & d \\ a & b \end{bmatrix}$$

$$(-1)^1 cb$$

# The Determinant of $3 \times 3$ matrices

$$\det \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = aei + bfg + cdh - ceg - bdi - afh$$

# The Determinant of $3 \times 3$ matrices

$$\det \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = aei + bfg + cdh - ceg - bdi - afh$$

$$\begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \rightarrow^0 \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$$

$$(-1)^0 aei$$

# The Determinant of $3 \times 3$ matrices

$$\det \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = aei + bfg + cdh - ceg - bdi - afh$$

$$\begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \rightarrow^2 \begin{bmatrix} g & h & i \\ a & b & c \\ d & e & f \end{bmatrix}$$

$$(-1)^2 gbf$$



# The Determinant of $3 \times 3$ matrices

$$\det \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = aei + bfg + cdh - ceg - bdi - afh$$

$$\begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \rightarrow^2 \begin{bmatrix} d & e & f \\ g & h & i \\ a & b & c \end{bmatrix}$$

$$(-1)^2 dhc$$

# The Determinant of $3 \times 3$ matrices

$$\det \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = aei + bfg + cdh - ceg - bdi - afh$$

$$\begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \rightarrow^1 \begin{bmatrix} g & h & i \\ d & e & f \\ a & b & c \end{bmatrix}$$

$$(-1)^1 gec$$

# The Determinant of $3 \times 3$ matrices

$$\det \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = aei + bfg + cdh - ceg - bdi - afh$$

$$\begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \rightarrow^1 \begin{bmatrix} d & e & f \\ a & b & c \\ g & h & i \end{bmatrix}$$

$$(-1)^1 dbi$$

# The Determinant of $3 \times 3$ matrices

$$\det \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = aei + bfg + cdh - ceg - bdi - afh$$

$$\begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \rightarrow^1 \begin{bmatrix} a & b & c \\ g & h & i \\ d & e & f \end{bmatrix}$$

$$(-1)^1 ahf$$

# Another Perspective

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

Let's row reduce an arbitrary  $2 \times 2$  matrix:

# Another Perspective

Let's row reduce an arbitrary  $3 \times 3$  matrix:

$$\begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$$

# Determinants and Invertibility

# Determinants and Invertibility

**Theorem.** A matrix is invertible if and only if  $\det(A) \neq 0$ .



# Determinants and Invertibility

**Theorem.** A matrix is invertible if and only if  $\det(A) \neq 0$ .

So we can yet again extend the IMT:

# Determinants and Invertibility

**Theorem.** A matrix is invertible if and only if  $\det(A) \neq 0$ .

So we can yet again extend the IMT:

- »  $A$  is invertible
- »  $\det(A) \neq 0$
- » 0 is not an eigenvalue

*These must be all true or all false.*

# Determinants (the definition we'll use)

$$\det(A) = \frac{(-1)^s}{c} U_{11} U_{22} \cdots U_{nn}$$

# Determinants (the definition we'll use)

$$\det(A) = \frac{(-1)^s}{c} U_{11} U_{22} \cdots U_{nn}$$

**Defintion.** The **determinant** of a matrix  $A$  is given by the above equation, where

# Determinants (the definition we'll use)

$$\det(A) = \frac{(-1)^s}{c} U_{11} U_{22} \cdots U_{nn}$$

**Defintion.** The **determinant** of a matrix  $A$  is given by the above equation, where

- $U$  is an echelon form of  $A$

# Determinants (the definition we'll use)

$$\det(A) = \frac{(-1)^s}{c} U_{11} U_{22} \cdots U_{nn}$$

**Defintion.** The **determinant** of a matrix  $A$  is given by the above equation, where

- $U$  is an echelon form of  $A$
- $s$  is the number of row swaps used to get  $U$

# Determinants (the definition we'll use)

$$\det(A) = \frac{(-1)^s}{c} U_{11} U_{22} \cdots U_{nn}$$

**Defintion.** The **determinant** of a matrix  $A$  is given by the above equation, where

- $U$  is an echelon form of  $A$
- $s$  is the number of row swaps used to get  $U$
- $c$  is the product of all scalings used to get  $U$

# Determinants (the definition we'll use)

$$\det(A) = \frac{(-1)^s}{c} \text{product of diagonal entries } U_{11}U_{22}\cdots U_{nn}$$

**Defintion.** The **determinant** of a matrix  $A$  is given by the above equation, where

- $U$  is an echelon form of  $A$
- $s$  is the number of row swaps used to get  $U$
- $c$  is the product of all scalings used to get  $U$



# Determinants (the definition we'll use)

$$\det(A) = \frac{(-1)^s}{c} \underbrace{U_{11}U_{22}\cdots U_{nn}}_{\text{product of diagonal entries}}$$

$c = 0$  if  $A$  is not invertible

**Defintion.** The **determinant** of a matrix  $A$  is given by the above equation, where

- $U$  is an echelon form of  $A$
- $s$  is the number of row swaps used to get  $U$
- $c$  is the product of all scalings used to get  $U$

# Example

$$\begin{bmatrix} 1 & 5 & 0 \\ 2 & 4 & -1 \\ 0 & -2 & 0 \end{bmatrix}$$

Let's find the determinant of this matrix:

## Example (Again)

$$\begin{bmatrix} 1 & 5 & 0 \\ 2 & 4 & -1 \\ 0 & -2 & 0 \end{bmatrix}$$

Let's find the determinant of this matrix again  
but with a different sequence of row operations:

The definition holds no matter  
which sequence of row  
operations you use.

# How To: Determinants

# How To: Determinants

**Question.** Determine the determinant of a matrix  $A$ .

# How To: Determinants

**Question.** Determine the determinant of a matrix  $A$ .

**Solution.**

# How To: Determinants

**Question.** Determine the determinant of a matrix  $A$ .

**Solution.**

1. Convert  $A$  to an echelon form  $U$ .



# How To: Determinants

**Question.** Determine the determinant of a matrix  $A$ .

**Solution.**

1. Convert  $A$  to an echelon form  $U$ .
2. Keep track of the number of row swaps you used, call this  $s$ , and the product of all scalings, call this  $c$

# How To: Determinants

**Question.** Determine the determinant of a matrix  $A$ .

**Solution.**

1. Convert  $A$  to an echelon form  $U$ .
2. Keep track of the number of row swaps you used, call this  $s$ , and the product of all scalings, call this  $c$
3. Determine the product of entries along the diagonal of  $U$ , call this  $P$ .

# How To: Determinants

**Question.** Determine the determinant of a matrix  $A$ .

**Solution.**

1. Convert  $A$  to an echelon form  $U$ .
2. Keep track of the number of row swaps you used, call this  $s$ , and the product of all scalings, call this  $c$ .
3. Determine the product of entries along the diagonal of  $U$ , call this  $P$ .
4. The determinant of  $A$  is  $\frac{sP}{c}$ .

# The Shorter Version

Beyond small matrices, we'll often just use computers.

**With NumPy:**

*`numpy.linalg.det(A)`*

# Properties of Determinants

# Properties of Determinants (1)

$$\det(AB) = \det(A) \det(B)$$

It follows that  $AB$  is invertible if and only if  $A$  and  $B$  are invertible

(we won't verify this)

# Question

*Use the fact that  $\det(AB) = \det(A)\det(B)$  to give an expression for  $\det(A^{-1})$  in terms of  $\det(A)$ .*

*Hint. What is  $\det(I)$ ?*

**Answer:**  $1/\det(A)$



# Properties of Determinants (2)

$$\det(A^T) = \det(A)$$

It follows that  $A^T$  is invertible if and only if  $A$  is invertible.

(we also won't verify this)

# Question

*If  $A^{-1} = A^T$ , then what are the possible values of  $\det(A)$ ?*

**Answer:  $\pm 1$**

# Properties of Determinants (3)

**Theorem.** If  $A$  is triangular, then  $\det(A)$  is the product of entries along the diagonal.

Verify:

# Question

$$\begin{bmatrix} 1 & 5 & -4 \\ -1 & -5 & 5 \\ -2 & -8 & 7 \end{bmatrix}$$

*Find the determinant of the above matrix.*

# Answer

# Characteristic Equation

**What kind of thing is the determinant, really?**



# What kind of thing is the determinant, really?

The determinant of a matrix  $A$  is an arithmetic expression written in terms of the entries of  $A$ .

# What kind of thing is the determinant, really?

The determinant of a matrix  $A$  is an arithmetic expression written in terms of the entries of  $A$ .

**But a matrix may not have numbers as entries.**

# What kind of thing is the determinant, really?

The determinant of a matrix  $A$  is an arithmetic expression written in terms of the entries of  $A$ .

**But a matrix may not have numbers as entries.**

We might think of the matrix  $A - \lambda I$  as having *polynomials* as entries.

# What kind of thing is the determinant, really?


The determinant of a matrix  $A$  is an arithmetic expression written in terms of the entries of  $A$ .

**But a matrix may not have numbers as entries.**

We might think of the matrix  $A - \lambda I$  as having *polynomials* as entries.


Then  $\det(A - \lambda I)$  is a **polynomial**.

# Reminder: Polynomial Roots



# Reminder: Polynomial Roots


A **root** of a polynomial  $p(x)$  is a value  $r$  such that  $p(r) = 0$ .



# Reminder: Polynomial Roots

A **root** of a polynomial  $p(x)$  is a value  $r$  such that  $p(r) = 0$ .

(A polynomial may have many roots)




# Reminder: Polynomial Roots

A **root** of a polynomial  $p(x)$  is a value  $r$  such that  $p(r) = 0$ .

(A polynomial may have many roots)

If  $r$  is a root of  $p(x)$ , then it is possible to find a polynomial  $q(x)$  such that

$$p(x) = (x - r)q(x)$$





# Characteristic Polynomial

# Characteristic Polynomial

**Definition.** The characteristic polynomial of a matrix  $A$  is  $\det(A - \lambda I)$  viewed as a polynomial in the variable  $\lambda$ .

# Characteristic Polynomial

**Definition.** The characteristic polynomial of a matrix  $A$  is  $\det(A - \lambda I)$  viewed as a polynomial in the variable  $\lambda$ .

**This is a polynomial with the eigenvalues of  $A$  as roots.**

# Characteristic Polynomial

**Definition.** The characteristic polynomial of a matrix  $A$  is  $\det(A - \lambda I)$  viewed as a polynomial in the variable  $\lambda$ .

**This is a polynomial with the eigenvalues of  $A$  as roots.**

So we can "solve" for the eigenvalues in the equation

$$\det(A - \lambda I) = 0$$

# Example: $2 \times 2$ Matrix\*

$$\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$$

Let's find the characteristic polynomial of this matrix:

\*we won't deal explicitly with matrices beyond  $2 \times 2$ , though there may be conceptual questions about larger matrices

## **Example: $2 \times 2$ Matrix\***

$$\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$$

Let's find the characteristic polynomial of this matrix:

**An Aside: What is this matrix?**

\*we won't deal explicitly with matrices beyond  $2 \times 2$ , though there may be conceptual questions about larger matrices

# A Special Linear Dynamical System

$$\mathbf{v}_{k+1} = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \mathbf{v}_k \quad \mathbf{v}_0 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

Consider the system given by the above matrix.

What does this system represent?:


# Fibonacci Numbers

$$F_0 = 0$$

$$F_1 = 1$$

$$F_k = F_{k-1} + F_{k-2}$$

```
define fib(n):  
    curr, next ← 0, 1  
    repeat n times:  
        curr, next ← next, curr + next  
    return curr
```




The Fibonacci numbers are defined in terms of a recurrence relation.

They seem to crop-up in nature, engineering, etc.



# Recall: The Fibonacci Matrix



The largest  
eigenvalue is  
the slope of  
this line

# Golden Ratio

$$\varphi = \frac{1 + \sqrt{5}}{2} \quad \frac{F_{k+1}}{F_k} \rightarrow \varphi \text{ as } k \rightarrow \infty$$

**This is the largest eigenvalue of  $\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$ .**

**To Come.** The "long term behavior" is the Fibonacci sequence is defined by the golden ratio.

# Example: Triangular matrix

$$\begin{bmatrix} 1 & -3 & 0 & 6 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 4 \end{bmatrix}$$

The characteristic polynomial of a triangular matrix comes pre-factored:

# How To: Finding Eigenvalues

# How To: Finding Eigenvalues

**Question.** Find all eigenvalues of the matrix  $A$ .

# How To: Finding Eigenvalues

**Question.** Find all eigenvalues of the matrix  $A$ .

**Solution.** Find the roots of the characteristic polynomial of  $A$ .

# How To: Finding Eigenvalues

**Question.** Find all eigenvalues of the matrix  $A$ .

**Solution.** Find the roots of the characteristic polynomial of  $A$ .

*In Reality.* We'll mostly just use

*`numpy.linalg.eig(A)`*

# An Observation: Multiplicity

$$\lambda^1(\lambda - 1)^2(\lambda - 4)^1 \text{ multiplicities}$$

In the examples so far, we've seen a number appear as a root multiple times.

This is called the **multiplicity** of the root.

**Is the multiplicity meaningful in this context?**



# Multiplicity and Dimension

**Theorem.** The dimension of the eigenspace of  $A$  for the eigenvalue  $\lambda$  is at most the multiplicity of  $\lambda$  in  $\det(A - \lambda I)$ .

The multiplicity is an upper bound on "how large" the eigenspace is.

# Example

Let  $A$  be a  $5 \times 5$  matrix with characteristic polynomial  $(x - 1)^3(x - 3)(x + 5)$ .

» What is  $\text{rank}(A)$ ?

» What is the minimum possible rank of  $A - I$ ?

# Application: Similar Matrices

**Definition.** Two square matrices  $A$  and  $B$  are **similar** if there is an invertible matrix  $P$  such that

$$A = P^{-1}BP$$

# Application: Similar Matrices

**Theorem.** Similar matrices have the same eigenvalues.

Verify:

# Summary

The determinant of a matrix is an arithmetic expression of its entries.

The characteristic polynomial is the determinant of  $A - \lambda I$  viewed as a polynomial of  $\lambda$ , and it tells us what the eigenvalues of a matrix are.