

# Analysis of variance

(Module 8)

Statistics (MAST20005) & Elements of Statistics (MAST90058)

Semester 2, 2020

## Contents

<b>1</b>	<b>Analysis of variance (ANOVA)</b>	<b>1</b>
1.1	Introduction . . . . .	1
1.2	One-way ANOVA . . . . .	2
1.3	Two-way ANOVA . . . . .	7
1.4	Two-way ANOVA with interaction . . . . .	10
<b>2</b>	<b>Hypothesis testing in regression</b>	<b>13</b>
2.1	Analysis of variance approach . . . . .	15
<b>3</b>	<b>Likelihood ratio tests</b>	<b>16</b>

## Aims of this module

- Introduce the **analysis of variance** technique, which builds upon the variance decomposition ideas in previous modules.
- Revisit linear regression and apply the ideas of hypothesis testing and analysis of variance.
- Discuss ways to derive optimal hypothesis tests.

## Overview

- **Analysis of variance (ANOVA).** Comparisons of more than two groups
- **Regression.** Hypothesis testing for simple linear regression
- **Likelihood ratio tests.** A method for deriving the best test for a given problem


## 1 Analysis of variance (ANOVA)

### 1.1 Introduction


#### Analysis of variance: introduction

- Initial aim: compare the means of **more than two** populations
- Broader and more advanced aims:
  - Explore components of variation
  - Evaluate the fit a (general) linear model
- Formulated as hypothesis tests
- Referred to as *analysis of variance*, or *ANOVA* for short
- Involves comparing different summaries of variation
- Related to the ‘analysis of variance’ and ‘variance decomposition’ formulae we derived previously

### Example: large variation between groups



### Example: smaller variation between groups



## 1.2 One-way ANOVA

### ANOVA: setting it up

- We have random samples from  $k$  populations, each having a normal distribution
- We sample  $n_i$  iid observations from the  $i$ th population, which has mean  $\mu_i$
- All populations assumed have the **same** variance,  $\sigma^2$
- Question of interest: do the populations all have the same mean?
- Hypotheses:

$$H_0: \mu_1 = \mu_2 = \dots = \mu_k = \mu \quad \text{versus} \quad H_1: \bar{H}_0$$

( $\bar{H}_0$  means 'not  $H_0$ ')

- This model is known as a *one-way ANOVA*, or *single-factor ANOVA*

### Notation

Population	Sample	Statistics	
$N(\mu_1, \sigma^2)$	$X_{11}, X_{12}, \dots, X_{1n_1}$	$\bar{X}_{1.}$	$S_1^2$
$N(\mu_2, \sigma^2)$	$X_{21}, X_{22}, \dots, X_{2n_2}$	$\bar{X}_{2.}$	$S_2^2$
$\vdots$	$\vdots$	$\vdots$	$\vdots$
$N(\mu_k, \sigma^2)$	$X_{k1}, X_{k2}, \dots, X_{kn_k}$	$\bar{X}_{k.}$	$S_k^2$
Overall		$\bar{X}_{..}$	

$$n = n_1 + \dots + n_k \quad (\text{total sample size})$$

$$\bar{X}_{i\cdot} = \frac{1}{n_i} \sum_{j=1}^{n_i} X_{ij} \quad (\text{group means})$$

$$\bar{X}_{..} = \frac{1}{n} \sum_{i=1}^k \sum_{j=1}^{n_i} X_{ij} = \frac{1}{n} \sum_{i=1}^k n_i \bar{X}_{i\cdot} \quad (\text{grand mean})$$

### Sum of squares (definitions)

- We now define statistics each called a *sum of squares* (*SS*)
- The *total SS* is:

$$SS(TO) = \sum_{i=1}^k \sum_{j=1}^{n_i} (X_{ij} - \bar{X}_{..})^2$$

- The *treatment SS*, or *between groups SS*, is:

$$SS(T) = \sum_{i=1}^k \sum_{j=1}^{n_i} (\bar{X}_{i\cdot} - \bar{X}_{..})^2 = \sum_{i=1}^k n_i (\bar{X}_{i\cdot} - \bar{X}_{..})^2$$

- The *error SS*, or *within groups SS*, is:

$$SS(E) = \sum_{i=1}^k \sum_{j=1}^{n_i} (X_{ij} - \bar{X}_{i\cdot})^2 = \sum_{i=1}^k (n_i - 1) S_i^2$$

### Analysis of variance decomposition

- It turns out that:

$$SS(TO) = SS(T) + SS(E)$$

- This is similar to the analysis of variance formulae we derived earlier, in simpler scenarios (iid model, regression model)
- We will use this relationship as a basis to derive a hypothesis test
- Let's first prove the relationship. . .
- Start with the 'add and subtract' trick:

$$\begin{aligned} SS(TO) &= \sum_{i=1}^k \sum_{j=1}^{n_i} (X_{ij} - \bar{X}_{..})^2 \\ &= \sum_{i=1}^k \sum_{j=1}^{n_i} (X_{ij} - \bar{X}_{i\cdot} + \bar{X}_{i\cdot} - \bar{X}_{..})^2 \\ &= \sum_{i=1}^k \sum_{j=1}^{n_i} (X_{ij} - \bar{X}_{i\cdot})^2 + \sum_{i=1}^k \sum_{j=1}^{n_i} (\bar{X}_{i\cdot} - \bar{X}_{..})^2 \\ &\quad + 2 \sum_{i=1}^k \sum_{j=1}^{n_i} (X_{ij} - \bar{X}_{i\cdot})(\bar{X}_{i\cdot} - \bar{X}_{..}) \\ &= SS(E) + SS(T) + CP \end{aligned}$$

- The cross-product term is:

$$\begin{aligned}
CP &= 2 \sum_{i=1}^k \sum_{j=1}^{n_i} (X_{ij} - \bar{X}_{i.})(\bar{X}_{i.} - \bar{X}_{..}) \\
&= 2 \sum_{i=1}^k (\bar{X}_{i.} - \bar{X}_{..}) \sum_{j=1}^{n_i} (X_{ij} - \bar{X}_{i.}) \\
&= 2 \sum_{i=1}^k (\bar{X}_{i.} - \bar{X}_{..})(n_i \bar{X}_{i.} - n_i \bar{X}_{i.}) \\
&= 0
\end{aligned}$$

- Thus, we have:

$$SS(TO) = SS(T) + SS(E)$$

### Sampling distribution of $SS(E)$

- The sample variance from the  $i$ th group,  $S_i^2$ , is an unbiased estimator of  $\sigma^2$  and we know that  $(n_i - 1)S_i^2/\sigma^2 \sim \chi_{n_i-1}^2$
- The samples from each group are independent, so we can usefully combine them,

$$\sum_{i=1}^k \frac{(n_i - 1)S_i^2}{\sigma^2} = \frac{SS(E)}{\sigma^2} \sim \chi_{n-k}^2$$

- Note that:  $(n_1 - 1) + (n_2 - 1) + \dots + (n_k - 1) = n - k$
- This also gives us an unbiased pooled estimator of  $\sigma^2$ ,

$$\hat{\sigma}^2 = \frac{SS(E)}{n - k}$$

- These results are true irrespective of whether  $H_0$  is true or not

### Null sampling distribution of $SS(TO)$

- If we assume  $H_0$ , we can derive simple expressions for the sampling distributions of the other sums of squares
- The combined data would be a sample of size  $n$  from  $N(\mu, \sigma^2)$ . Hence  $SS(TO)/(n - 1)$  is an unbiased estimator of  $\sigma^2$  and

$$\frac{SS(TO)}{\sigma^2} \sim \chi_{n-1}^2$$

### Null sampling distribution of $SS(T)$

- Under  $H_0$ , we have  $\bar{X}_{i.} \sim N(\mu, \frac{\sigma^2}{n_i})$
- $\bar{X}_{1.}, \bar{X}_{2.}, \dots, \bar{X}_{k.}$  are independent
- (Can think of this as a sample of sample means, and then think about what its variance estimator is)
- It is possible to show that (proof not shown):

$$\sum_{i=1}^k \frac{n_i(\bar{X}_{i.} - \bar{X}_{..})^2}{\sigma^2} = \frac{SS(T)}{\sigma^2} \sim \chi_{k-1}^2$$

and that this is independent of  $SS(E)$

## Null sampling distributions


In summary, under  $H_0$ :

$$\frac{SS(TO)}{\sigma^2} = \frac{SS(E)}{\sigma^2} + \frac{SS(T)}{\sigma^2}$$


$$\frac{SS(TO)}{\sigma^2} \sim \chi_{n-1}^2, \quad \frac{SS(E)}{\sigma^2} \sim \chi_{n-k}^2, \quad \frac{SS(T)}{\sigma^2} \sim \chi_{k-1}^2,$$

$SS(E)$  and  $SS(T)$  are independent

$H_0$  is true



$H_1$  is true



$SS(T)$  under  $H_1$

- What happens if  $H_1$  is true?
- The population means differ, which will make  $SS(T)$  larger
- Let's make this precise...

- Let  $\bar{\mu} = n^{-1} \sum_{i=1}^k n_i \mu_i$ , and then,

$$\begin{aligned}
\mathbb{E}[SS(T)] &= \mathbb{E} \left[ \sum_{i=1}^k n_i (\bar{X}_{i.} - \bar{X}_{..})^2 \right] = \mathbb{E} \left[ \sum_{i=1}^k n_i \bar{X}_{i.}^2 - n \bar{X}_{..}^2 \right] \\
&= \sum_{i=1}^k n_i \mathbb{E}(\bar{X}_{i.}^2) - n \mathbb{E}(\bar{X}_{..}^2) \\
&= \sum_{i=1}^k n_i [\text{var}(\bar{X}_{i.}) + \mathbb{E}(\bar{X}_{i.})^2] - n [\text{var}(\bar{X}_{..}) + \mathbb{E}(\bar{X}_{..})^2] \\
&= \sum_{i=1}^k n_i \left[ \frac{\sigma^2}{n_i} + \mu_i^2 \right] - n \left[ \frac{\sigma^2}{n} + \bar{\mu}^2 \right] \\
&= (k-1)\sigma^2 + \sum_{i=1}^k n_i (\mu_i - \bar{\mu})^2
\end{aligned}$$

- Under  $H_0$  the second term is zero and we have,

$$\frac{\mathbb{E}(SS(T))}{k-1} = \sigma^2$$

- Otherwise (under  $H_1$ ), the second term is positive and gives,

$$\frac{\mathbb{E}(SS(T))}{k-1} > \sigma^2$$

- In contrast, we always have,

$$\frac{E(SS(E))}{n-k} = \sigma^2$$

### F-test statistic

- This motivates using the following as our test statistic:

$$F = \frac{SS(T)/(k-1)}{SS(E)/(n-k)}$$

- Under  $H_0$ , we have  $F \sim F_{k-1, n-k}$ , since it is the ratio of independent  $\chi^2$  random variables
- Under  $H_1$ , the numerator will tend to be larger
- Therefore, reject  $H_0$  if  $F > c$
- This is known as an *F-test*

### ANOVA table

The test quantities are often summarised using an *ANOVA table*:


Source	df	SS	MS	F
Treatment	$k-1$	$SS(T)$	$MS(T) = \frac{SS(T)}{k-1}$	$\frac{MS(T)}{MS(E)}$
Error	$n-k$	$SS(E)$	$MS(E) = \frac{SS(E)}{n-k}$	
Total	$n-1$	$SS(TO)$		

Notes:

- MS = ‘Mean square’
- $\hat{\sigma}^2 = MS(E)$  is an unbiased estimator

## Example (one-way ANOVA)

Force required to pull out window studs in 5 positions on a car window.



```
> head(data1)
  Position Force
1         1    92
2         1    90
3         1    87
4         1   105
5         1    86
6         1    83

> table(data1$Position)

1 2 3 4 5
7 7 7 7 7

> model1 <- lm(Force ~ factor(Position), data = data1)
> anova(model1)
Analysis of Variance Table

Response: Force
          Df Sum Sq Mean Sq F value    Pr(>F)
factor(Position)  4 16672.1   4168.0   44.202 3.664e-12 ***
Residuals       30  2828.9     94.3
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Notes:

- Need to use `factor()` to denote categorical variables
- R doesn't provide a 'Total' row, but we don't need it
- `Residuals` is the 'Error' row
- `Pr(>F)` is the p-value for the F-test

We conclude that the mean force required to pull out the window studs varies between the 5 positions on the car window (e.g. p-value < 0.01)

This was obvious from the boxplots: positions 1 & 2 are quite different from 3, 4 & 5

## 1.3 Two-way ANOVA

### Two factors

- In one-way ANOVA, the observations were partitioned into  $k$  groups
- In other words, they were defined by a single categorical variable ('factor')
- What if we had two such variables?

- We can extend the procedure to give *two-way ANOVA*, or *two-factor ANOVA*
- For example, the fuel consumption of a car may depend on type of petrol and the brand of tyres

### Two-way ANOVA: setting it up

- Factor 1 has  $a$  levels, Factor 2 has  $b$  levels
- Suppose we have exactly one observation per factor combination
- Observe  $X_{ij}$  with factor 1 at level  $i$  and factor 2 at level  $j$
- Gives a total of  $n = ab$  observations
- Assume  $X_{ij} \sim N(\mu_{ij}, \sigma^2)$ ,  $i = 1, \dots, a$ ,  $j = 1, \dots, b$ , and that these are independent
- Consider the model:

$$\mu_{ij} = \mu + \alpha_i + \beta_j$$

$$\text{with } \sum_{i=1}^a \alpha_i = 0, \sum_{j=1}^b \beta_j = 0$$

- $\mu$  is an overall effect,  $\alpha_i$  is the effect of the  $i$ th row and  $\beta_j$  the effect of the  $j$ th column.
- For example,  $a = 4$  and  $b = 4$ ,

	1	2	3	4
1	$\mu + \alpha_1 + \beta_1$	$\mu + \alpha_1 + \beta_2$	$\mu + \alpha_1 + \beta_3$	$\mu + \alpha_1 + \beta_4$
2	$\mu + \alpha_2 + \beta_1$	$\mu + \alpha_2 + \beta_2$	$\mu + \alpha_2 + \beta_3$	$\mu + \alpha_2 + \beta_4$
3	$\mu + \alpha_3 + \beta_1$	$\mu + \alpha_3 + \beta_2$	$\mu + \alpha_3 + \beta_3$	$\mu + \alpha_3 + \beta_4$
4	$\mu + \alpha_4 + \beta_1$	$\mu + \alpha_4 + \beta_2$	$\mu + \alpha_4 + \beta_3$	$\mu + \alpha_4 + \beta_4$

- We are usually interested in  $H_{0A}: \alpha_1 = \alpha_2 = \dots = \alpha_a = 0$  or  $H_{0B}: \beta_1 = \beta_2 = \dots = \beta_b = 0$
- Let

$$\bar{X}_{..} = \frac{1}{ab} \sum_{i=1}^a \sum_{j=1}^b X_{ij}, \quad \bar{X}_{i.} = \frac{1}{b} \sum_{j=1}^b X_{ij}, \quad \bar{X}_{.j} = \frac{1}{a} \sum_{i=1}^a X_{ij}$$

- Arguing as before,

$$\begin{aligned} SS(TO) &= \sum_{i=1}^a \sum_{j=1}^b (X_{ij} - \bar{X}_{..})^2 \\ &= \sum_{i=1}^a \sum_{j=1}^b [(\bar{X}_{i.} - \bar{X}_{..}) + (\bar{X}_{.j} - \bar{X}_{..}) \\ &\quad + (X_{ij} - \bar{X}_{i.} - \bar{X}_{.j} + \bar{X}_{..})]^2 \\ &= b \sum_{i=1}^a (\bar{X}_{i.} - \bar{X}_{..})^2 + a \sum_{j=1}^b (\bar{X}_{.j} - \bar{X}_{..})^2 \\ &\quad + \sum_{i=1}^a \sum_{j=1}^b (X_{ij} - \bar{X}_{i.} - \bar{X}_{.j} + \bar{X}_{..})^2 \\ &= SS(A) + SS(B) + SS(E) \end{aligned}$$

- If both  $\alpha_1 = \dots = \alpha_a = 0$  and  $\beta_1 = \dots = \beta_b = 0$ , then we have  $SS(A)/\sigma^2 \sim \chi_{a-1}^2$ ,  $SS(B)/\sigma^2 \sim \chi_{b-1}^2$  and  $SS(E)/\sigma^2 \sim \chi_{(a-1)(b-1)}^2$  and these variables are independent (proof not shown)
- Reject  $H_{0A}: \alpha_1 = \dots = \alpha_a = 0$  at significance level  $\alpha$  if:

$$F_A = \frac{SS(A)/(a-1)}{SS(E)/((a-1)(b-1))} > c$$

where  $c$  is the  $1 - \alpha$  quantile of  $F_{a-1, (a-1)(b-1)}$



- Reject  $H_{0B}: \beta_1 = \dots = \beta_b = 0$  at significance level  $\alpha$  if:

$$F_B = \frac{SS(B)/(b-1)}{SS(E)/((a-1)(b-1))} > c$$

where  $c$  is the  $1 - \alpha$  quantile of  $F_{b-1, (a-1)(b-1)}$


### ANOVA table

Source	df	SS	MS	F
Factor A	$a - 1$	$SS(A)$	$MS(A) = \frac{SS(A)}{a-1}$	$\frac{MS(A)}{MS(E)}$
Factor B	$b - 1$	$SS(B)$	$MS(B) = \frac{SS(B)}{b-1}$	$\frac{MS(B)}{MS(E)}$
Error	$(a - 1)(b - 1)$	$SS(E)$	$MS(E) = \frac{SS(E)}{(a-1)(b-1)}$	
Total	$ab - 1$	$SS(TO)$		

### Example (two-way ANOVA)

Data on fuel consumption for three types of car ( $A$ ) and four types of fuel ( $B$ ).

```
> head(data2)
  Car Fuel Consumption
1   1    1          16
2   1    2          18
3   1    3          21
4   1    4          21
5   2    1          14
6   2    2          15
```



```
> model2 <- lm(Consumption ~ factor(Car) + factor(Fuel),
+             data = data2)
> anova(model2)
Analysis of Variance Table
```

```
Response: Consumption
          Df Sum Sq Mean Sq F value    Pr(>F)
factor(Car)  2     24  12.0000     18 0.002915 **
factor(Fuel)  3     30  10.0000     15 0.003401 **
Residuals    6      4   0.6667
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

From this we conclude there is a clear difference in fuel consumption between cars (we reject  $H_{0A}: \alpha_1 = \alpha_2 = \alpha_3$ ) and also between fuels (we reject  $H_{0B}: \beta_1 = \beta_2 = \beta_3 = \beta_4$ ).

## 1.4 Two-way ANOVA with interaction

### Interaction terms

- In the previous example we assumed an additive model:

$$\mu_{ij} = \mu + \alpha_i + \beta_j$$

- This assumes, for example, that the relative effect of petrol 1 is the same for all cars.
- If it is not true, then there is a *statistical interaction* (or simply an *interaction*) between the factors
- A more general model, which includes interactions, is:

$$\mu_{ij} = \mu + \alpha_i + \beta_j + \gamma_{ij}$$

where  $\gamma_{ij}$  is the *interaction term* associated with combination  $(i, j)$ .

- In addition to our previous assumptions, we also impose:

$$\sum_{i=1}^a \gamma_{ij} = 0, \quad \text{and} \quad \sum_{j=1}^b \gamma_{ij} = 0$$

- The terms  $\alpha_i$  and  $\beta_j$  are called *main effects*
- When written out as a table they are also often referred to as the *row effects* and *column effects* respectively
- Writing this out as a table:

	1	2	3	4
1	$\mu + \alpha_1 + \beta_1 + \gamma_{11}$	$\mu + \alpha_1 + \beta_2 + \gamma_{12}$	$\mu + \alpha_1 + \beta_3 + \gamma_{13}$	$\mu + \alpha_1 + \beta_4 + \gamma_{14}$
2	$\mu + \alpha_2 + \beta_1 + \gamma_{21}$	$\mu + \alpha_2 + \beta_2 + \gamma_{22}$	$\mu + \alpha_2 + \beta_3 + \gamma_{23}$	$\mu + \alpha_2 + \beta_4 + \gamma_{24}$
3	$\mu + \alpha_3 + \beta_1 + \gamma_{31}$	$\mu + \alpha_3 + \beta_2 + \gamma_{32}$	$\mu + \alpha_3 + \beta_3 + \gamma_{33}$	$\mu + \alpha_3 + \beta_4 + \gamma_{34}$
4	$\mu + \alpha_4 + \beta_1 + \gamma_{41}$	$\mu + \alpha_4 + \beta_2 + \gamma_{42}$	$\mu + \alpha_4 + \beta_3 + \gamma_{43}$	$\mu + \alpha_4 + \beta_4 + \gamma_{44}$

- We are now interested in testing whether:
  - the row effects are zero
  - the column effects are zero
  - the interactions are zero (do this first!)
- To make inferences about the interactions we need more than one observation per cell
- Let  $X_{ijk}$ ,  $i = 1, \dots, a$ ,  $j = 1, \dots, b$ ,  $k = 1, \dots, c$  be the  $k$ th observation for combination  $(i, j)$
- Let

$$\bar{X}_{ij\cdot} = \frac{1}{c} \sum_{k=1}^c X_{ijk}$$

$$\bar{X}_{i\cdot\cdot} = \frac{1}{bc} \sum_{j=1}^b \sum_{k=1}^c X_{ijk}$$

$$\bar{X}_{\cdot j\cdot} = \frac{1}{ac} \sum_{i=1}^a \sum_{k=1}^c X_{ijk}$$

$$\bar{X}_{\dots} = \frac{1}{abc} \sum_{i=1}^a \sum_{j=1}^b \sum_{k=1}^c X_{ijk}$$

- and as before

$$\begin{aligned}
SS(TO) &= \sum_{i=1}^a \sum_{j=1}^b \sum_{k=1}^c (X_{ijk} - \bar{X}_{...})^2 \\
&= bc \sum_{i=1}^a (\bar{X}_{i..} - \bar{X}_{...})^2 + ac \sum_{j=1}^b (\bar{X}_{.j.} - \bar{X}_{...})^2 \\
&\quad + c \sum_{i=1}^a \sum_{j=1}^b (\bar{X}_{ij.} - \bar{X}_{i..} - \bar{X}_{.j.} + \bar{X}_{...})^2 \\
&\quad + \sum_{i=1}^a \sum_{j=1}^b \sum_{k=1}^c (X_{ijk} - \bar{X}_{ij.})^2 \\
&= SS(A) + SS(B) + SS(AB) + SS(E)
\end{aligned}$$

### Test statistics

- Familiar arguments show that to test

$$H_{0AB}: \gamma_{ij} = 0, \quad i = 1, \dots, a, \quad j = 1, \dots, b$$

we may use the statistic

$$F = \frac{SS(AB)/[(a-1)(b-1)]}{SS(E)/[ab(c-1)]}$$

which has a  $F$  distribution with  $(a-1)(b-1)$  and  $ab(c-1)$  degrees of freedom.

- To test

$$H_{0A}: \alpha_i = 0, \quad i = 1, \dots, a$$

we may use the statistic

$$F = \frac{SS(A)/[(a-1)]}{SS(E)/[ab(c-1)]}$$

which has a  $F$  distribution with  $(a-1)$  and  $ab(c-1)$  degrees of freedom.

- To test

$$H_{0B}: \beta_j = 0, \quad j = 1, \dots, b$$

we may use the statistic

$$F = \frac{SS(B)/[(b-1)]}{SS(E)/[ab(c-1)]}$$

which has a  $F$  distribution with  $(b-1)$  and  $ab(c-1)$  degrees of freedom.

### ANOVA table

Source	df	SS	MS	$F$
Factor A	$a - 1$	$SS(A)$	$MS(A) = \frac{SS(A)}{a-1}$	$\frac{MS(A)}{MS(E)}$
Factor B	$b - 1$	$SS(B)$	$MS(B) = \frac{SS(B)}{b-1}$	$\frac{MS(B)}{MS(E)}$
Factor AB	$(a-1)(b-1)$	$SS(AB)$	$MS(AB) = \frac{SS(AB)}{(a-1)(b-1)}$	$\frac{MS(AB)}{MS(E)}$
Error	$ab(c-1)$	$SS(E)$	$MS(E) = \frac{SS(E)}{ab(c-1)}$	
Total	$abc - 1$	$SS(TO)$		

### Example (two-way ANOVA with interaction)

- Six groups of 18 people
- Each person takes an arithmetic test: the task is to add three numbers together
- The numbers are presented either in a down array or an across array; this defines 2 levels of factor  $A$
- The numbers have either one, two or three digits; this defines 3 levels of factor  $B$

- The response variable,  $X$ , is the average number of problems completed correctly over two 90-second sessions
- Example of adding **one-digit** numbers in an **across** array:

$$2 + 5 + 1 = ?$$

- Example of adding **two-digit** numbers in an **down** array:

$$\begin{array}{r} 13 \\ 87 \\ + 51 \\ \hline ? \end{array}$$

```
> head(data3)
      A B    X
1 down 1 19.5
2 down 1 18.5
3 down 1 32.0
4 down 1 21.5
5 down 1 28.5
6 down 1 33.0

> table(data3[, 1:2])
      B
A      1  2  3
down   18 18 18
across 18 18 18

> model3 <- lm(X ~ factor(A) * factor(B), data = data3)
> anova(model3)
Analysis of Variance Table


Response: X
          Df Sum Sq Mean Sq  F value    Pr(>F)
factor(A)    1   48.7    48.7    2.8849 0.09246 .
factor(B)    2 8022.7   4011.4  237.7776 < 2e-16 ***
factor(A):factor(B) 2  185.9    93.0   5.5103 0.00534 **
Residuals   102 1720.8    16.9
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Note the use of '\*' in the model formula.

The interaction is significant at a 5% level (or even at 1%).

## Interaction plot

```
with(data3, interaction.plot(B, A, X, col = "blue"))
```



## Beyond the F-test

- We have rejected the null... now what?
- This is often only the beginning of a statistical analysis of this type of data
- Will be interested in more detailed inferences, e.g. CIs/tests about individual parameters
- You know enough to be able to work some of this out...
- ...and later subjects will go into this in more detail (e.g. MAST30025)

## 2 Hypothesis testing in regression

### Recap of simple linear regression

- $Y$  a response variable, e.g. student's grade in first-year calculus
- $x$  a predictor variable, e.g. student's high school mathematics mark
- Data: pairs  $(x_1, y_1), \dots, (x_n, y_n)$
- Linear regression model:

$$Y_i = \alpha + \beta(x_i - \bar{x}) + \epsilon_i$$

where  $\epsilon_i \sim N(0, \sigma^2)$  is a random error

- **Note:**  $\alpha$  here plays the same role as  $\alpha_0$  from Module 5. We have dropped the '0' subscript for convenience, and also to avoid confusion with its use to denote null hypotheses.
- The MLE (and OLS) estimators are:

$$\hat{\alpha} = \bar{Y}, \quad \hat{\beta} = \frac{\sum_{i=1}^n Y_i(x_i - \bar{x})}{\sum_{i=1}^n (x_i - \bar{x})^2}$$

- and

$$\hat{\sigma}^2 = \frac{1}{n-2} \sum_{i=1}^n [Y_i - \hat{\alpha} - \hat{\beta}(x_i - \bar{x})]^2$$

- We also derived:

$$\begin{aligned}\hat{\alpha} &\sim N\left(\alpha, \frac{\sigma^2}{n}\right) \\ \hat{\beta} &\sim N\left(\beta, \frac{\sigma^2}{\sum_{i=1}^n (x_i - \bar{x})^2}\right)\end{aligned}$$

- and

$$\frac{(n-2)\hat{\sigma}^2}{\sigma^2} = \frac{\sum_{i=1}^n [Y_i - \hat{\alpha} - \hat{\beta}(x_i - \bar{x})]^2}{\sigma^2} \sim \chi_{n-2}^2$$

- From these we obtain,

$$\begin{aligned}T_\alpha &= \frac{\hat{\alpha} - \alpha}{\hat{\sigma}/\sqrt{n}} \sim t_{n-2} \\ T_\beta &= \frac{\hat{\beta} - \beta}{\hat{\sigma}/\sqrt{\sum_{i=1}^n (x_i - \bar{x})^2}} \sim t_{n-2}\end{aligned}$$

- We used these previously to construct confidence intervals
- We can also use them to construct hypothesis tests
- For example, to test  $H_0: \beta = \beta_0$  versus  $H_1: \beta \neq \beta_0$  (or  $\beta > \beta_0$  or  $\beta < \beta_0$ ), we use  $T_\beta$  as the test statistic

### Example: testing the slope parameter ( $\beta$ )

- Data: 10 pairs of scores on a preliminary test and a final exam
- Estimates:  $\hat{\alpha} = 81.3$ ,  $\hat{\beta} = 0.742$ ,  $\hat{\sigma}^2 = 27.21$
- Test  $H_0: \beta = 0$  versus  $H_1: \beta \neq 0$  with a 1% significance level
- Reject  $H_0$  if:

$$|T_\beta| \geq 3.36 \quad (0.995 \text{ quantile of } t_8)$$

- For the observed data,

$$t_\beta = \frac{0.742 - 0}{\sqrt{27.21/756.1}} = 3.91$$

so we reject  $H_0$ , concluding there is sufficient evidence that the slope differs from zero.

### Note regarding the intercept parameter ( $\alpha$ )

- Software packages (such as R) will typically fit the model:

$$Y_i = \alpha + \beta x_i + \epsilon_i$$

- This is equivalent to

$$Y_i = \alpha^* + \beta(x_i - \bar{x}) + \epsilon_i$$

where  $\alpha = \alpha^* - \beta\bar{x}$

- The formulation  $Y_i = \alpha^* + \beta(x_i - \bar{x}) + \epsilon$  is easier to examine theoretically.
- We saw that

$$\hat{\alpha}^* = \bar{Y}, \quad \text{and} \quad \hat{\alpha} = \bar{Y} - \hat{\beta}\bar{x}$$

- $\hat{\alpha}$  or  $\hat{\alpha}^*$  are rarely of direct interest

### Using R

Use R to fit the regression model for the slope example:

```
> m1 <- lm(final_exam ~ prelim_test)
> summary(m1)
```

Call:

```
lm(formula = final_exam ~ prelim_test)
```

Residuals:

Min	1Q	Median	3Q	Max
-6.883	-3.264	-0.530	3.438	8.470

Coefficients:

	Estimate	Std. Error	t value	Pr(> t )
(Intercept)	30.6147	13.0622	2.344	0.04714 *
prelim_test	0.7421	0.1897	3.912	0.00447 **

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 5.217 on 8 degrees of freedom

Multiple R-Squared: 0.6567, Adjusted R-squared: 0.6137

F-statistic: 15.3 on 1 and 8 DF, p-value: 0.004471

The t-value and the p-value are for testing  $H_0: \alpha = 0$  and  $H_0: \beta = 0$  respectively.

## Interpreting the R output

- Usually most interested in testing  $H_0: \beta = 0$  versus  $H_1: \beta \neq 0$
- If we reject  $H_0$  then we conclude there is sufficient evidence of (at least) a linear relationship between the mean response and  $x$
- In the example,

$$t = \frac{0.7421}{0.1897} = 3.912$$

- This test statistic has a  $t$ -distribution with  $10 - 2 = 8$  degrees of freedom, and the associated p-value is  $0.00447 < 0.05$  so at the 5% level of significance we reject  $H_0$
- It is also possible to represent this test using an ANOVA table

## 2.1 Analysis of variance approach

### Deriving the variance decomposition formula

- Independent pairs  $(x_1, Y_1), \dots, (x_n, Y_n)$
- Parameter estimates,

$$\hat{\alpha} = \bar{Y}, \quad \hat{\beta} = \frac{\sum_{i=1}^n Y_i(x_i - \bar{x})}{\sum_{i=1}^n (x_i - \bar{x})^2}$$

- Fitted value (estimated mean),

$$\hat{Y}_i = \bar{Y} + \hat{\beta}(x_i - \bar{x})$$

- Do the ‘add and subtract’ trick again:

$$\begin{aligned} \sum_{i=1}^n (Y_i - \bar{Y})^2 &= \sum_{i=1}^n (Y_i - \hat{Y}_i + \hat{Y}_i - \bar{Y})^2 \\ &= \sum_{i=1}^n (Y_i - \hat{Y}_i)^2 + \sum_{i=1}^n (\hat{Y}_i - \bar{Y})^2 \\ &\quad + 2 \sum_{i=1}^n (Y_i - \hat{Y}_i)(\hat{Y}_i - \bar{Y}) \end{aligned}$$

- Deal with the cross-product term,

$$\begin{aligned} \sum_{i=1}^n (Y_i - \hat{Y}_i)(\hat{Y}_i - \bar{Y}) &= \sum_{i=1}^n \left[ Y_i - \bar{Y} - \hat{\beta}(x_i - \bar{x}) \right] \hat{\beta}(x_i - \bar{x}) \\ &= \hat{\beta} \sum_{i=1}^n \left[ Y_i - \bar{Y} - \hat{\beta}(x_i - \bar{x}) \right] (x_i - \bar{x}) \\ &= \hat{\beta} \left[ \sum_{i=1}^n (Y_i - \bar{Y})(x_i - \bar{x}) - \hat{\beta} \sum_{i=1}^n (x_i - \bar{x})^2 \right] \\ &= \hat{\beta} \left[ \sum_{i=1}^n Y_i(x_i - \bar{x}) - \hat{\beta} \sum_{i=1}^n (x_i - \bar{x})^2 \right] \\ &= 0 \end{aligned}$$

- That gives us,

$$\sum_{i=1}^n (Y_i - \bar{Y})^2 = \sum_{i=1}^n (Y_i - \hat{Y}_i)^2 + \sum_{i=1}^n (\hat{Y}_i - \bar{Y})^2$$

- We can write this as follows,

$$SS(TO) = SS(E) + SS(R)$$

where  $SS(R)$  is the *regression SS* or *model SS*

- The regression SS quantifies the variation **due to** the straight line

- The error SS quantifies the variation **around** the straight line
- To complete the specification,

$$MS(E) = \frac{SS(E)}{n-2} = \frac{1}{n-2} \sum_{i=1}^n (Y_i - \hat{Y}_i)^2 = \hat{\sigma}^2$$

$$MS(R) = \frac{SS(R)}{1} = \sum_{i=1}^n (\hat{Y}_i - \bar{Y})^2$$

- Then we have the test statistic,

$$F = \frac{MS(R)}{MS(E)} \sim F_{1,n-2}$$

### ANOVA table

Source	df	SS	MS	F
Model	1	$SS(R)$	$MS(R) = \frac{SS(R)}{1}$	$\frac{MS(R)}{MS(E)}$
Error	$n-2$	$SS(E)$	$MS(E) = \frac{SS(E)}{n-2}$	
Total	$n-1$	$SS(TO)$		

### Using R

```
> anova(m1)
Analysis of Variance Table

Response: final_exam
          Df Sum Sq Mean Sq F value    Pr(>F)
prelim_test  1 416.39   416.39   15.301 0.004471 **
Residuals    8 217.71    27.21
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Notes:
```

- The F-statistic tests the ‘significance of the regression’
- That is,  $H_0: \beta = 0$  versus  $H_1: \beta \neq 0$

## 3 Likelihood ratio tests

### Is there a ‘best’ test?

- We have examined a variety of commonly used tests
- We used test statistics that:
  - Seemed useful
  - We were familiar with
- Did we use the ‘best’ one?
- Is there a general procedure for finding a good/best test statistic?
- We will introduce a general procedure now, and discuss why it is optimal later in the semester

### Likelihood ratio test

- The *likelihood ratio test* (LRT) is a general procedure that can find the best test for a given problem



- Suppose we have  $H_0$  and  $H_1$  and both are composite and of the form:

$$H_0: \theta \in A_0 \quad \text{versus} \quad H_1: \theta \in A_1$$

where  $A_0$  and  $A_1$  are sets of possible parameter values consistent with each of the hypotheses.

- Note: we have mostly dealt with  $A_0$  that has only one element (simple null hypothesis)
- The *likelihood ratio* is:

$$\lambda = \frac{L_0}{L_1} = \frac{\max_{\theta \in A_0} L(\theta)}{\max_{\theta \in A_1} L(\theta)}$$

- $L$  is the likelihood function
- Clearly  $\lambda \geq 0$
- Large  $\lambda \Rightarrow$  more support for  $H_0$  over  $H_1$
- $\lambda$  near zero  $\Rightarrow$  more support for  $H_1$  over  $H_0$
- Therefore, we want a critical region of the form,

$$\lambda \leq k$$

- Choose  $k$  to give the desired significance level

### Example 1 (likelihood ratio test)

- $X_i \sim N(\mu, \sigma^2 = 5)$ , i.e.  $\sigma$  is known
- $H_0: \mu = 162$  versus  $H_1: \mu \neq 162$
- When  $H_0$  is true,  $\mu = 162$  so  $L_0 = L(162)$
- When  $H_1$  is true, need to maximise the likelihood,  $L_1 = L(\hat{\theta}) = L(\bar{x})$
- The likelihood ratio is,

$$\begin{aligned} \lambda &= \frac{L_0}{L_1} = \frac{L(162)}{L(\bar{x})} = \frac{(10\pi)^{-n/2} \exp \left[ -\frac{1}{10} \sum_{i=1}^n (x_i - 162)^2 \right]}{(10\pi)^{-n/2} \exp \left[ -\frac{1}{10} \sum_{i=1}^n (x_i - \bar{x})^2 \right]} \\ &= \exp \left[ -\frac{n}{10} (\bar{x} - 162)^2 \right] \end{aligned}$$

- $\lambda \leq k$  same as

$$\frac{|\bar{x} - 162|}{\sigma/\sqrt{n}} \geq c$$

- A critical region for a size  $\alpha$  test is

$$\frac{|\bar{x} - 162|}{\sigma/\sqrt{n}} \geq \Phi^{-1}(1 - \alpha/2)$$

- Note: this required knowledge of the distribution of  $\bar{X}$ !

### Example 2 (likelihood ratio test)

- $X_i \sim N(\mu, \sigma^2)$ , i.e.  $\sigma$  is unknown
- $H_0: \mu = \mu_0$  versus  $H_1: \mu \neq \mu_0$
- Under  $H_0$  we have  $\mu = \mu_0$ , and under  $H_1$  we need to use its MLE
- Under either hypothesis,  $\sigma^2$  is unspecified, so in both cases we need its MLE (conditional on the specified value of  $\mu$ ).
- So, under  $H_0$  we use:

$$\hat{\mu} = \mu_0, \quad \hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \mu_0)^2$$

- And under  $H_1$  we use:

$$\hat{\mu} = \bar{x}, \quad \hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2$$

- Some simplification yields

$$\lambda = \left[ \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{\sum_{i=1}^n (x_i - \mu_0)^2} \right]^{n/2}$$

- and

$$\sum_{i=1}^n (x_i - \mu_0)^2 = \sum_{i=1}^n (x_i - \bar{x} + \bar{x} - \mu_0)^2 = \sum_{i=1}^n (x_i - \bar{x})^2 + n(\bar{x} - \mu_0)^2$$

- Substitute and rearrange to get

$$\lambda = \left[ \frac{1}{1 + \frac{n(\bar{x} - \mu_0)^2}{\sum_{i=1}^n (x_i - \bar{x})^2}} \right]^{n/2}$$

- Therefore, we have  $\lambda \leq k$  when,

$$\frac{n(\bar{x} - \mu_0)^2}{\frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2} \geq c$$

- When  $H_0$  is true,  $\sqrt{n}(\bar{X} - \mu_0)/\sigma \sim N(0, 1)$  and  $\sum_{i=1}^n (X_i - \bar{X})^2/\sigma^2 \sim \chi_{n-1}^2$ , and is independent of  $\bar{X}$ .
- Therefore,

$$\begin{aligned} T &= \frac{\sqrt{n}(\bar{X} - \mu_0)/\sigma}{\sqrt{\frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2/\sigma^2}} \\ &= \frac{\sqrt{n}(\bar{X} - \mu_0)}{\sqrt{\frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2}} = \frac{\bar{X} - \mu_0}{S/\sqrt{n}} \sim t_{n-1} \end{aligned}$$

- So we reject  $H_0$  when  $|T|$  is too large, with the following critical region for a test with significance level  $\alpha$ ,

$$|T| \geq d, \quad \text{where } d \text{ is the } 1 - \frac{\alpha}{2} \text{ quantile of } t_{n-1}$$

## Remarks

- Usually easy to find the **form** of the test
- What is harder is to find the corresponding sampling distribution
- Manipulating  $\lambda$  until we have something whose distribution we know can be tricky!
- Many of the standard tests arise from the likelihood ratio

## Asymptotic distribution & optimality

- The likelihood ratio itself is a statistic and therefore has a sampling distribution.
- For large sample sizes, this approaches a known distribution
- Also, the LRT gives the optimal test
- We will cover this theory later in the semester