

Question 4 Solution:

Part a:

Given information:

Let,

$x_1, x_2, x_3 \sim (N(\mu, \sigma^2))$ be a sequence of independent normal random variables,

$$\bar{x} = \frac{x_1 + x_2 + x_3}{3}$$

$$\mathbf{x}^T = (x_1, x_2, x_3)^T$$

Supposed to be x^T as noted!

$$\mathbf{y} = (x_1 - \bar{x}, x_2 - \bar{x}, x_3 - \bar{x})^T$$

To solve A from:

$$\mathbf{y} = A\mathbf{x}$$

$$\begin{bmatrix} x_1 - \bar{x} \\ x_2 - \bar{x} \\ x_3 - \bar{x} \end{bmatrix} = \frac{1}{3} \begin{bmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

$$A = \frac{1}{3} \begin{bmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{bmatrix}$$

Where A is symmetric and idempotent!

$$A^2 = \frac{1}{9} \begin{bmatrix} 6 & -3 & -3 \\ -3 & 6 & -3 \\ -3 & -3 & 6 \end{bmatrix} = \frac{1}{3} \begin{bmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{bmatrix} = A$$

Part b: Finding the rank of A

Proof: that there is a linear combination for any columns?

```

```{r}
A = matrix(c(2,-1,-1,-1,2,-1,-1,-1,2)/3,3,3)
```

```

```

      [,1]      [,2]      [,3]
[1,] 0.6666667 -0.3333333 -0.3333333
[2,] -0.3333333 0.6666667 -0.3333333
[3,] -0.3333333 -0.3333333 0.6666667

```

```

# Finding rank of A

```

```

```{r}
rankMatrix(A)[1]
```

```

```

[1] 2

```

Each column all added up together gives us 0!.

$$x_1 + x_2 + x_3 = 0$$

Can be written as,

$$x_1 = -x_2 - x_3$$

That are linearly dependant and similar for x_2 and x_3

Hence $r(A) = 2$

Part c: Computing $E[y^T y]$

Using Theorem 3.5:

$$E[y^T A y] = \text{tr}(AV) + \mu^T A \mu$$

since $A = I$,

$$= \text{tr}(V) + \mu^T \mu$$

$$V = \text{var} y = \text{var} Ax = A \text{var} x A^T$$

since A is symmetric and idempotent!!

$$\text{var}(x_i) = \begin{bmatrix} \sigma^2 & 0 & 0 \\ 0 & \sigma^2 & 0 \\ 0 & 0 & \sigma^2 \end{bmatrix}$$

$$V = \frac{1}{3} \begin{bmatrix} 2\sigma^2 & -1 & -1 \\ -1 & 2\sigma^2 & -1 \\ -1 & -1 & 2\sigma^2 \end{bmatrix}$$

$$\mu = E[y] = E[Ax] = AE[x]$$

$$= \frac{1}{3} \begin{bmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{bmatrix} \begin{bmatrix} \mu \\ \mu \\ \mu \end{bmatrix} = 0$$

$$E[y^T y] = \text{tr}\left(\frac{1}{3} \begin{bmatrix} 2\sigma^2 & -1 & -1 \\ -1 & 2\sigma^2 & -1 \\ -1 & -1 & 2\sigma^2 \end{bmatrix}\right) + 0$$

$$= 2\sigma^2$$

Part d:

Using Theorem 3.5:

Proof:

Assuming that A is idempotent and has rank k . Because it is symmetric, it can be diagonalised. Let the (orthogonal) diagonalising matrix be P .

$$D = P^T A P = \begin{bmatrix} \lambda_1 & \dots & 0 \\ \dots & \lambda_2 & \dots \\ 0 & \dots & \lambda_k \end{bmatrix}$$

since A is symmetric and idempotent, all eigenvalues are either 0 or 1. We know from definition:

$$\text{tr}(A) = r(A) = k$$

$$A = \frac{1}{3} \begin{bmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{bmatrix}$$

$$A^2 = A = \frac{1}{3} \begin{bmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{bmatrix}$$

from Part 4b, we find out the rank and trace of matrix A we found in Part 4a. Is also is the same number of degrees of freedom for the chi squared distribution.

$$tr(A) = r(A) = 2$$

Therefore, A must have two eigenvalues of 1 and one eigenvalue of 0.

Using Theorem 3.5 and Corollary 3.7:

with our non central parameter λ !

$$\lambda = \frac{1}{2} \mu^T A \mu$$

$$= \frac{1}{2} [\mu \quad \mu \quad \mu] \frac{1}{3} \begin{bmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{bmatrix} \begin{bmatrix} \mu \\ \mu \\ \mu \end{bmatrix}$$

$$= 0$$

\iff : if and only if

$$E[y] = E \begin{bmatrix} x_1 - \mu \\ x_2 - \mu \\ x_3 - \mu \end{bmatrix}$$

Since x_1, x_2 and x_3 is identically independently distributed! and taking the expectation of the expectation is the expectation itself!

$$E[y] = E \begin{bmatrix} \mu - \mu \\ \mu - \mu \\ \mu - \mu \end{bmatrix} = 0$$

NOTE: $\mu = \bar{x}$

In which case,

$$\frac{y^T y}{\sigma^2}$$

is just the sum of two independent standard normal's. This is just an ordinary (central) chi squared distribution χ_2^2 .

with expectation of 2 and variance of 4 with 2 degrees of freedom. In which A is symmetric and idempotent!