

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/51525403>

Review: Lead Exposure in Battery Manufacturing and Recycling in Developing Countries and Among Children in Nearby Communities

Article in *Journal of Occupational and Environmental Hygiene* · September 2011

DOI: 10.1080/15459624.2011.601710 · Source: PubMed

CITATIONS

84

READS

2,069

2 authors, including:



[Amod Pokhrel](#)

University of California, Berkeley

26 PUBLICATIONS 561 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Nepal TB Study [View project](#)


This article was downloaded by: [Optimised: University of California, Berkeley]

On: 28 July 2011

Access details: Access Details: [subscription number 935379560]

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Occupational and Environmental Hygiene

Publication details, including instructions for authors and subscription information:

<http://oeh.informaworld.com/soeh/title~content=t713657996>

Review: Lead Exposure in Battery Manufacturing and Recycling in Developing Countries and Among Children in Nearby Communities

Perry Gottesfeld^a; Amod K. Pokhrel^b

^a Occupational Knowledge International, San Francisco, California ^b School of Public Health, University of California, Berkeley, California

First published on: 27 July 2011

To cite this Article Gottesfeld, Perry and Pokhrel, Amod K.(2011) 'Review: Lead Exposure in Battery Manufacturing and Recycling in Developing Countries and Among Children in Nearby Communities', Journal of Occupational and Environmental Hygiene, 8: 9, 520 — 532, First published on: 27 July 2011 (iFirst)

To link to this Article: DOI: 10.1080/15459624.2011.601710

URL: <http://dx.doi.org/10.1080/15459624.2011.601710>

PLEASE SCROLL DOWN FOR ARTICLE

The American Conference of Governmental Industrial Hygienists (<http://www.acgih.org/>) and the American Industrial Hygiene Association (<http://www.aiha.org/>) have licensed the Taylor & Francis Group to publish this article and other materials. To join the American Conference of Governmental Industrial Hygienists visit <http://www.acgih.org/Members/>. To join the American Industrial Hygiene Association visit <http://www.aiha.org/Content/BecomeMember/becomemember-splash.htm>.

Full terms and conditions of use: <http://oeh.informaworld.com/terms-and-conditions-of-access.pdf>

This article maybe used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Review: Lead Exposure in Battery Manufacturing and Recycling in Developing Countries and Among Children in Nearby Communities

Perry Gottesfeld¹ and Amod K. Pokhrel²

¹Occupational Knowledge International, San Francisco, California

²School of Public Health, University of California, Berkeley, California

The battery industry is the largest consumer of lead, using an estimated 80% of the global lead production. The industry is also rapidly expanding in emerging market countries. A review of published literature on exposures from lead-acid battery manufacturing and recycling plants in developing countries was conducted. The review included studies from 37 countries published from 1993 to 2010 and excluded facilities in developed countries, such as the United States and those in Western Europe, except for providing comparisons to reported findings. The average worker blood lead level (BLL) in developing countries was 47 µg/dL in battery manufacturing plants and 64 µg/dL in recycling facilities. Airborne lead concentrations reported in battery plants in developing countries averaged 367 µg/m³, which is 7-fold greater than the U.S. Occupational Safety and Health Administration's 50 µg/m³ permissible exposure limit. The geometric mean BLL of children residing near battery plants in developing countries was 19 µg/dL, which is about 13-fold greater than the levels observed among children in the United States. The blood lead and airborne lead exposure concentrations for battery workers were substantially higher in developing countries than in the United States. This disparity may worsen due to rapid growth in lead-acid battery manufacturing and recycling operations worldwide. Given the lack of regulatory and enforcement capacity in most developing countries, third-party certification programs may be the only viable option to improve conditions.

Keywords battery, blood lead level, developing countries, lead, recycling, third-party certification

Correspondence to: Perry Gottesfeld, Occupational Knowledge International, 4444 Geary Blvd., Suite 300, San Francisco, CA 94118; e-mail: pgottesfeld@okinternational.org.

INTRODUCTION

Lead poisoning remains a significant occupational disease and a ubiquitous environmental health threat to children. Lead causes numerous adverse health effects, including damage to the nervous system, the kidneys, the cardiovascular

system, the hematopoietic system, and the reproductive system.^(1–8) In children, blood lead concentration is associated with a significant decrease in academic performance and lower standardized test scores (including IQ test scores) and is linked with hyperactive and violent behavior.⁽⁹⁾ The International Agency for Research on Cancer (IARC) has classified lead and inorganic lead compounds in Group 2A as probable human carcinogens.⁽¹⁰⁾

Occupational and environmental lead exposure in the developing world arguably may have a more profound effect than in developed countries. Poor nutrition, common in developing countries, increases lead absorption through the gastrointestinal tract.⁽¹¹⁾ Lead poisoning imposes a range of hidden costs on developing countries. One study estimated that a 50% decrease in childhood blood lead levels in Nigeria could save the country \$1 billion annually; the health care cost of lead exposure for adults is estimated to total \$7 billion.⁽¹²⁾ Lead-induced decrements to health status may further decrease productivity, which will lead to less investment and the continuation of the cycle of poverty.⁽¹³⁾

The Lead Battery Industry

The battery industry is the principal consumer of lead and uses an estimated 80% of annual primary lead (mined) and secondary lead (recycled) production.⁽¹⁴⁾ Approximately 50% of global lead production is derived from recycling lead batteries.⁽¹⁵⁾ These batteries are used primarily in vehicles for starting, lighting, and ignition purposes, but are also used in photovoltaic solar installations and telecommunications systems to store energy. In developing countries where the power supplies are often inconsistent, lead batteries are routinely used in homes and businesses to back up computer systems, lights, and appliances when outages occur. Electric vehicles are becoming an important market for lead batteries; electric bikes in China presently account for more than 20% of the country's lead demand.^(16,17) The expected increase in automobile, solar, telecommunications, and computer sales in the

developing world will increase the production and recycling of lead batteries.

It is estimated that between 60,000 and 70,000 people are employed globally in lead battery manufacturing, in addition to a similar number working in mining, smelting, and refining.⁽¹⁵⁾ The vast majority of these workers live in the developing world. The efficiency of lead recovery from battery recycling can vary greatly, and the crude methods employed in most developing countries result in the release of millions of tons of lead into the environment.⁽¹⁸⁾ The Basel Convention, which came into force in 1992, restricts the export of used lead batteries to developing countries where environmentally sound recycling cannot be ensured. The United States exports a substantial number of used lead batteries to Mexico, South Korea, India, and other countries, which underlines the importance of investigating lead exposures in these countries.^(19,20)

Given forecasts that the lead battery industry will double in size in many developing countries over the next 5 to 10 years, we performed this review of lead exposures among workers in lead battery manufacturing and recycling and among children living in nearby communities.^(21–23) We discuss barriers and opportunities to better control these exposures and environmental emissions.

METHODS

We reviewed the published literature on lead exposures due to lead-acid battery manufacturing and recycling operations (which include reconditioning used batteries) in developing countries. We used the terms “lead, blood lead levels (BLLs), battery manufacturing, recycling, and children,” and searched databases of the National Library of Medicine, MEDLINE service, and Web of Science and Global Health. More than 232 studies appeared on these search terms, among which 98 were relevant for this review. Pertinent references cited in the identified studies were also reviewed.

We limited our review to studies with an abstract published in English from 1993 to 2010 and excluded facilities from our search if they were located in countries considered “developed,” (Table I). Information on relevant measures from developed countries was used to compare aggregate results obtained in this review. Given the improving economies in many countries during this time period and the lack of a universal classification system, we used the term “developing countries” to distinguish the latter from the “developed countries” listed in Table I. We used the 1993 start point for three reasons. First, there are few pertinent studies prior to 1993. Second, similar manufacturing technologies have been employed around the world since 1993. Third, the start point follows the implementation of the Basel Convention mentioned previously.

From the 98 selected studies identified in our search, 84 provided summary measures of BLLs (the arithmetic mean, geometric mean, or median) among exposed worker cohorts and, in some cases, airborne lead concentration inside the plants.

TABLE I. List of Developed Countries Excluded from the Literature Search

Region	Countries
Europe	Andorra Austria Belgium Denmark Finland France Germany Iceland Ireland Italy Liechtenstein Luxembourg Monaco Netherlands Northern Ireland Norway Scotland Spain Sweden Switzerland United Kingdom
North America	Canada United States
Oceania	Australia New Zealand
Asia	Japan

Ten studies reported summary measures of BLLs in children living in the vicinity of battery manufacturing and recycling facilities. We attempted to contact the authors of studies that presented incomplete statistics (e.g., only the range of BLLs) and did not provide a summary cohort exposure.^(24–31) If no additional information could be obtained, we excluded the study.

For the BLL data, we grouped those studies reporting the arithmetic mean BLL and grouped separately those studies reporting either the geometric mean or median BLL. For individual studies that reported separate arithmetic mean BLLs for two or more subgroups, we computed an average value weighted by the number fraction in each subgroup. For individual studies that reported a geometric mean (GM) and a geometric standard deviation (GSD) for the BLLs, we estimated the cohort arithmetic unweighted mean (μ) by the equation: $\mu = \text{GM} \times \exp[0.5 \times (\ln \text{GSD})^2]$.⁽³²⁾ Because the distribution of BLLs tends to be right skewed, we judged that the GM BLL was approximately equal to the median BLL and therefore grouped studies reporting only the GM with those reporting only the median value.⁽³³⁾ For each group, we calculated the mean or median of all the individual study values in the group.

We evaluated the BLL trend from 1993 to 2010 among the 61 studies that reported arithmetic mean BLLs by regressing the individual study results against the year of study publication. Blood lead level samples were almost certainly collected before the publication dates listed in the tables. The Theil-Sen median slope for trend in Stata software (version 10; StataCorp LLC, College Station, Texas) was used to evaluate these cohort BLLs over time.⁽³⁴⁾ This statistical analysis is a nonparametric estimate of slope, identified as the Kendall-Theil slope (also known as the Kendall-Sen).^(34–36) The method is resistant to the effects of outliers and non-normality in residuals, as commonly observed in BLL data.⁽³⁷⁾ We also used the Student's t-test to compare average BLLs among lead battery plant workers reported in the period 1993–2002 versus the period 2003–2010. We chose 2002/2003 as the cut point to divide the data set into approximately equal numbers of BLL reports. Five countries, accounting for 40 of the studies, were represented in both groups allowing for comparisons over these time periods.

RESULTS

Table II lists arithmetic mean BLLs among workers at lead battery plants in 20 countries as reported in 61 studies; a total of 8350 workers are represented in the table. Arithmetic mean cohort values ranged from 22 to 128 $\mu\text{g/dL}$, and the mean and median of the 61 cohort arithmetic mean values are 47 $\mu\text{g/dL}$ and 42 $\mu\text{g/dL}$, respectively. Table III lists median BLLs among workers at 12 lead battery plants in six countries as reported in seven studies; a total of 1189 workers are represented in the table. The median of the cohort median values is 37 $\mu\text{g/dL}$. Table IV lists arithmetic mean BLLs among workers at 14 lead battery recycling/reconditioning plants in 13 countries as reported in 13 studies; a total of 479 workers are represented in the table. Arithmetic mean cohort levels range from 43 to 113 $\mu\text{g/dL}$, and the mean and median of the cohort arithmetic mean values are 64 $\mu\text{g/dL}$ and 60 $\mu\text{g/dL}$, respectively.

Table V lists the arithmetic mean or GM value for lead concentrations in air (based on area and personal samples) in 12 battery manufacturing plants, and the arithmetic mean lead concentration in air (based on area samples) in one battery recycling facility. One study that summarized airborne lead exposure in Chinese lead battery industries was excluded because of discrepancies in the results.⁽³⁸⁾ Due to the lack of detail regarding workplace conditions associated with the samples in most studies, we combined the area and personal sample results within the same industry. Arithmetic mean airborne lead concentrations ranged from 39 to 1260 $\mu\text{g/m}^3$, and geometric mean airborne lead concentrations ranged from 33 to 355 $\mu\text{g/m}^3$. Among the studies that provided an arithmetic mean airborne lead concentration, the average and median value of the eight means are 367 $\mu\text{g/m}^3$ and 264 $\mu\text{g/m}^3$, respectively. Among the six studies that provided a GM airborne lead concentration, the median value of the five GMs is 103 $\mu\text{g/m}^3$.

Table VI lists 10 studies that reported the arithmetic mean or median BLLs for groups of children residing near lead battery manufacturing and recycling plants in developing countries; a total of 2284 children are represented in the table. These studies included different age ranges (up to 15 years) and distances from the subject plants (0.18 to 5 km). The arithmetic mean BLL in groups of children range from 9 to 71 $\mu\text{g/dL}$; the average and median values of the nine group arithmetic means are 29 $\mu\text{g/dL}$ and 19 $\mu\text{g/dL}$. Reported median BLLs for two other groups of children were 7.3 and 8.1 $\mu\text{g/dL}$.

As seen in Figure 1, there appears to be a modest decline in worker BLLs during the period 1993 to 2010, although the trend is not statistically significant (Theil-Sen median slope = -0.69 , with two-sided 95% confidence interval $[-1.54; 0.16]$). When the cohort arithmetic mean BLLs are grouped by publication date with 2002/2003 as the cut point, the group average BLL is higher during 1993–2002 (50 $\mu\text{g/dL}$) than during 2003–2010 (44 $\mu\text{g/dL}$), although the difference is not statistically significant ($p = 0.29$ for a two-tailed test).

DISCUSSION

Tables II–IV demonstrate that lead battery industry workers in developing countries, at both the front and back end of the battery life cycle, experience BLLs that far exceed health protection guidelines. Based on a comprehensive review of lead toxicity in adults, Kosnett et al.⁽¹⁾ recommended that individuals be removed from occupational lead exposure if a single blood lead concentration exceeds 30 $\mu\text{g/dL}$, or if two successive blood lead concentrations measured over a 4-week interval are ≥ 20 $\mu\text{g/dL}$. The ACGIH[®] has established a biological exposure index of 30 $\mu\text{g/dL}$.⁽³⁹⁾ The Centers for Disease Control and Prevention (CDC) recommends that BLLs should be maintained at < 5 $\mu\text{g/dL}$ for occupationally exposed women who are or may become pregnant.⁽⁴⁰⁾

BLLs among U.S. and U.K. workers in lead battery manufacturing facilities are substantially lower than the levels reported in this review.^(41–43) Although no comprehensive U.S. occupational lead registry exists, several states have compiled extensive data on BLLs from workers in the lead battery industry via laboratory reporting requirements. In California, BLLs above 25 $\mu\text{g/dL}$ must be reported to the state. In practice, most blood lead test results (at all levels) are voluntarily reported. In 1999 (the last year that data was reported), among 1931 workers from this industry, no reported worker BLLs exceeded 60 $\mu\text{g/dL}$, and only 36 (1.9%) worker BLLs exceeded 40 $\mu\text{g/dL}$.⁽⁴⁴⁾ From 1993 through 2001 in Washington State, only five workers from the lead battery manufacturing industry reported BLLs exceeding 60 $\mu\text{g/dL}$.⁽⁴⁵⁾

The U.S. National Institute for Occupational Safety and Health (NIOSH) had compiled BLL report data for 2007 from states that collect and summarize laboratory testing reports. Among 1743 production workers in storage battery industries (NAICS 335911) in California, Iowa, and Oregon, 325 (18.6%) had BLLs ≥ 25 $\mu\text{g/dL}$, and 21 (1.2%) had reported BLLs ≥ 40 $\mu\text{g/dL}$ (personal communication from Walter A. Alarcon,

TABLE II. Arithmetic Mean BLLs of Workers in Lead Battery Manufacturing Plants

Year	Author	Country	n	Arithmetic Mean BLL ($\mu\text{g/dL}$)
1993	Zhang ⁽⁷⁰⁾	China	128	125.7 ^A
1993	Far et al. ⁽⁷¹⁾	Singapore	25	48.9
1994	Ibiebele ⁽⁵²⁾	Barbados	20	35.3
1994	Makino et al. ⁽⁷²⁾	Philippines	199	64.5
1994	Singh et al. ⁽⁷³⁾	India	15	52.0
1995	Kim et al. ⁽⁷⁴⁾	South Korea	66	45.7
1995	Schwartz, S. et al. ⁽⁷⁵⁾	South Korea	308	29.1
1996	Solliway et al. ⁽⁷⁶⁾	Israel	34	40.7
1996	Cordeiro et al. ⁽⁷⁷⁾	Brazil	20	49.4
1996	Chang et al. ⁽⁷⁸⁾	Taiwan	27	48.6
1997	Lai et al. ⁽⁷⁹⁾	Taiwan	219	56.9
1997	Yucesoy et al. ⁽⁸⁰⁾	Turkey	20	59.4
1997	Schwartz et al. ⁽⁸¹⁾	South Korea	57	25.4
1997	Kuo et al. ⁽⁸²⁾	Taiwan	5	43.6
1998	Ho et al. ⁽⁸³⁾	Singapore	50	32.5
1998	Ehrlich et al. ⁽⁸⁴⁾	South Africa	382	53.5
1998	Vaglenov et al. ⁽⁸⁵⁾	Bulgaria	22	60.9 ^A
1998	Williams et al. ⁽⁸⁶⁾	Trinidad-West Indies	22	23.0
1998	Undeger and Basaran ⁽⁸⁷⁾	Turkey	25	74.8
1999	Froom et al. ⁽⁸⁸⁾	Israel	94	38.1
1999	Chuang et al. ⁽⁶³⁾	Taiwan	392	23.9
2000	Caldeira et al. ⁽⁸⁹⁾	Brazil	17	63.8
2000	Restrepo et al. ⁽⁹⁰⁾	Colombia	43	98.5
2000	Roh et al. ⁽⁹¹⁾	South Korea	49	42.6
2000	Basaran and Undeger ⁽⁹²⁾	Turkey	25	74.8
2000	Ratzon et al. ⁽⁹³⁾	Israel	63	42.5
2000	Chuang et al. ⁽⁹⁴⁾	Taiwan	206	31.8
2001	Duydu et al. ⁽⁹⁵⁾	Turkey	31	36.3
2001	Vaglenov et al. ⁽⁹⁶⁾	Bulgaria	103	55.9 ^A
2002	Hwang ⁽⁹⁷⁾	Taiwan	96	29.6
2002	Wang et al. ⁽⁵³⁾	Taiwan	229	58.6 ^B
2002	Sonmez et al. ⁽⁹⁸⁾	Turkey	13	25.3
2003	Mishra et al. ⁽⁹⁹⁾	India	34	128.1
2003	Nusier et al. ⁽²⁹⁾	Jordan	83	51.4
2003	Palus et al. ⁽¹⁰⁰⁾	Poland	44	50.4
2003	Suzen et al. ⁽¹⁰¹⁾	Turkey	71	34.5
2003	Pizent et al. ⁽¹⁰²⁾	Croatia	143	41.4
2003	Duydu and Suzen ⁽¹⁰³⁾	Turkey	71	34.5
2003	Lormphongs et al. ⁽¹⁰⁴⁾	Thailand	182	21.7 ^C
2004	Gurer-Orhan et al. ⁽¹⁰⁵⁾	Turkey	20	54.6
2004	Chuang et al. ⁽¹⁰⁶⁾	Taiwan	544	23.2
2004	Bagc et al. ⁽¹⁰⁷⁾	Turkey	22	36.8
2004	Heo et al. ⁽²⁵⁾	South Korea	1123	22.9 ^D
2005	Duydu et al. ⁽¹⁰⁸⁾	Turkey	50	40.1
2005	Chuang et al. ⁽¹⁰⁹⁾	Taiwan	855	30.4
2005	Ravichandran et al. ⁽³¹⁾	India	171	32.3 ^D
2005	Karakaya et al. ⁽¹¹⁰⁾	Turkey	23	72.7
2006	Patil et al. ⁽¹¹¹⁾	India	28	53.6

(Continued on next page)

TABLE II. Arithmetic Mean BLLs of Workers in Lead Battery Manufacturing Plants (Continued)

Year	Author	Country	n	Arithmetic Mean BLL ($\mu\text{g/dL}$)
2006	Kuruville et al. ⁽¹¹²⁾	India	52	42.4
2006	Li et al. ⁽¹¹³⁾	Taiwan	597	27.1
2006	Engin et al. ⁽¹¹⁴⁾	Turkey	30	63.5
2006	Chen et al. ⁽¹¹⁵⁾	China	25	32.0
2006	Wananukul et al. ⁽²⁴⁾	Thailand	389	35.5 ^{A,D}
2007	Lin and J. Tai-y ⁽¹¹⁶⁾	China	135	42.2
2007	Chia ⁽¹¹⁷⁾	Vietnam	276	25.3 ^C
2008	Chuang et al. ⁽¹¹⁸⁾	Taiwan	120	37.5 ^E
2008	Raviraja et al. ⁽⁵⁹⁾	India	5	81.1 ^A
2009	Kašuba et al. ⁽¹¹⁹⁾	Croatia	15	43.6
2009	Hsu et al. ⁽¹²⁰⁾	Taiwan	80	40.2
2010	Nsheiwat et al. ⁽¹²¹⁾	Jordan	22	27.5
2010	Gao et al. ⁽¹²²⁾	China	135	42.5 ^F
			Total = 8350	Mean = 47 Median = 42

Note: n = number of samples.

^A $\mu\text{g/dL}$ conversion from $\mu\text{mol/L}$.

^BThe number-weighted average of male and female workers reported.

^CArithmetic mean computed based on the reported GM and GSD.

^DAggregate average provided by authors of the study.

^EThe number-weighted average of Taiwanese and Thai nationals reported.

^FThe number-weighted average of ALAD11 and ALAD12 genotypes reported.

CDC/NIOSH Adult Blood Epidemiology and Surveillance (ABLES) Program project officer, February 2, 2011).^(46,47) In an Environmental Assessment filed in 2010, the average BLLs for employees at two lead battery plants at Bristol, Tennessee, and Columbus, Georgia, were reported as 13 and 15 $\mu\text{g/dL}$, respectively.⁽⁴¹⁾ Since 1993, occupational lead exposures in

battery manufacturing declined substantially in the United States due to improvements in ventilation and work practices; environmental lead emission also decreased.⁽⁴⁸⁾

In the U.K., summary data from 2003/2004 indicated that of 11,011 male lead workers undergoing medical surveillance, about 5% had BLLs $\geq 50 \mu\text{g/dL}$, and less than 1% had BLLs

TABLE III. Median BLLs of Workers in Lead Battery Manufacturing Plants

Year	Author	Country	n	Median BLL ($\mu\text{g/dL}$)
1994	Dos Santos et al. ⁽¹²³⁾	Brazil	166	36.8
1996	Chia et al. ⁽¹²⁴⁾	Singapore	72	39.2
1997	Chia ⁽¹²⁵⁾	Singapore	50	37.1
1997	Sithisarankul et al. ⁽¹²⁶⁾	South Korea	65	27.9
1997	Chia et al. ⁽¹²⁷⁾	Singapore	28	42.6
1998	Bergdahl et al. ⁽¹²⁸⁾	Russia	42	27.0
1998	Jakubowski et al. ⁽¹²⁹⁾	Poland	460	34.9 ^A
1998	Jakubowski et al. ⁽¹²⁹⁾	Poland	8	48.2 ^A
1998	Jakubowski et al. ⁽¹²⁹⁾	Poland	26	44.9 ^A
1998	Jakubowski et al. ⁽¹²⁹⁾	Poland	70	49.6 ^A
1998	Jakubowski et al. ⁽¹²⁹⁾	Poland	10	50.3 ^A
2008	Sun et al. ⁽¹³⁰⁾	China	155	20.2 ^{A,B}
2008	Sun et al. ⁽¹³⁰⁾	China	37	15.5 ^{A,C}
			Total = 1189	Median = 37

^AGeometric mean considered to approximate the median value.

^BMale.

^CFemale.

TABLE IV. Arithmetic Mean Blood Lead Levels of Workers in Lead Battery Recycling and Reconditioning Plants

Year	Author	Country	n	BLL ($\mu\text{g/dL}$)
1995	Yeh et al. ⁽¹³¹⁾	South Korea	31	63.0
1998	Wang et al. ⁽¹³²⁾	Taiwan	64	66.5
1998	Corzo and Naveda ⁽¹³³⁾	Venezuela	11	45.8
2000	Mehdi et al. ⁽¹³⁴⁾	Iraq	37	60.6
2000	Suplido et al. ⁽¹³⁵⁾	Philippines	40	54.2
2001	Cardenas-Bustamante et al. ⁽¹³⁶⁾	Colombia	94	88.0
2002	Kumar et al. ⁽¹³⁷⁾	India	36	55.6
2002	Ayatollahi ⁽¹³⁸⁾	Iran	21	46.8
2008	Minozzo et al. ⁽¹³⁹⁾	Brazil	53	59.4
2008	Jiang et al. ⁽¹⁴⁰⁾	China	7	63.5
2008	Ramirez ⁽²⁸⁾	Peru	41	37.7 ^A
2009	Peter ⁽¹⁴¹⁾	Nigeria-Lagos	20	112.5 ^B
2009	Peter ⁽¹⁴¹⁾	Nigeria-Ibadan	20	93.9 ^B
2010	Nsheiwat et al. ⁽¹²¹⁾	Israel	4	43.4
			Total = 479	Mean = 64 Median = 60

^AAggregate average provided by authors of the study.^BTwo values are from different factories.**TABLE V. Workplace Airborne Lead Concentrations (Arithmetic and Geometric Mean) in Lead Battery Manufacturing and Recycling Plants**

Year	Author	Country	Sample Duration, n	Arithmetic Mean Airborne Lead ($\mu\text{g/m}^3$)	Geometric Mean Airborne Lead ($\mu\text{g/m}^3$)
1993	Far et al. ⁽⁷¹⁾	Singapore	6 hr, n = 25	NA	98 ^A
1994	Ibiebele ⁽⁵²⁾	Barbados	8 hr, n = 80	NA	33.1 ^B
1997	Lai et al. ⁽⁷⁹⁾	Taiwan	NR, 219	190	NA
1998	Vaglenov et al. ⁽⁸⁵⁾	Bulgaria	7 hr, n = 22	447	NA
1998	Ho et al. ⁽⁸³⁾	Singapore	6 hr, n = 50	88.6 ^A	NA
1998	Ehrlich et al. ⁽⁸⁴⁾	South Africa	NR, n = 30	NA	145 ^C
2000	Hwang et al. ⁽¹⁴²⁾	Taiwan	7 hr, n = 81	NA	107 ^A
2002	Wang et al. ⁽⁵³⁾	China	8 hr, n = 229	206 ^A	NA
2002	Hwang et al. ⁽⁹⁷⁾	Taiwan	4 hr, n = 96	NA	27.7
2004	Donguk and Namwon ⁽¹⁴³⁾	South Korea	8 hr, n = 44	NA	354.8 ^A
2005	Ravichandran et al. ⁽³¹⁾	India	8 hr, n = 22	384	NA
2006	Chen et al. ⁽¹¹⁵⁾	China	NR, n = 24	1260	NA
2007	Dyosi ⁽⁵⁰⁾	South Africa	8 hr (NR)	321 ^{D,E}	NA
2008	Sun et al. ⁽¹³⁰⁾	China	NR	39	NA
Mean				367	NA
Median				264	103

Note: NA = not applicable, NR = not reported.

^APersonal air samples worn on workers.^BCombined average values for wet ($31.9 \mu\text{g/m}^3$) and dry season ($34.2 \mu\text{g/m}^3$).^CMedian value.^DCombined average values for 2001 and 2002.^ELead battery recycling plant.

TABLE VI. BLLs Reported in Children Residing in Proximity to Lead Battery Manufacturing and Recycling Facilities

Year	Author	Country	n	Age	Distance	Mean BLL ($\mu\text{g/dL}$)	Median BLL ($\mu\text{g/dL}$)
1998	Morales Bonilla et al. ⁽¹⁴⁴⁾	Nicaragua	97	6 mo–13 years	200 m	17.2	NA
1998	Tabaku et al. ⁽¹⁴⁷⁾	Albania	84	2–5 years	< 2 km	43.4	NA
1998	Tabaku et al. ⁽¹⁴⁷⁾	Albania	145	2–15 years	<2 km	23.5	NA
1999	Kaul and Mukerjee ⁽¹⁴⁵⁾	Dominican Republic	116	6 mo–10 years	180 m	71.0	NA
1999	Saraci and K. Ziegler- Skylakakis ⁽¹⁴⁶⁾	Albania	194	10–15 years	Not reported	19.3	NA
2000	Suplido and Ong ⁽¹³⁵⁾	Philippines	10	6 mo–12 years	20 m	49.9	NA
2003	Cortes-Maramba et al. ⁽¹⁴⁸⁾	Philippines	40	10 years ^A	1–5 km	12.9	NA
2006	Safi et al. ⁽¹⁴⁹⁾	Palestinian Authority (Gaza)	435	2–6 years	Not reported	8.6	NA
2007	De Freitas et al. ⁽¹⁵⁰⁾	Brazil	850	6 years ^A	<1 km	NA	7.3
2009	Ahmed et al. ⁽¹⁵¹⁾	Pakistan	190	1–12 years	1 km	12.0	
2010	Khan et al. ⁽⁵⁷⁾	Pakistan	123	1–6 years	Not specified	NA	8.1
Total = 2284						Mean = 29 Median = 19	NA Median = 8

^AAverage age reported.

$\geq 60 \mu\text{g/dL}$.⁽⁴³⁾ Earlier U.K. summary data from 2000–2001 showed that among 3384 male lead battery workers, 11.5% had BLLs $\geq 50 \mu\text{g/dL}$, and 2.1% had BLLs $\geq 60 \mu\text{g/dL}$.⁽⁴²⁾

Inhalation is considered the most important lead intake route, in general, in occupational settings.^(49–51) Ingestion is also a significant source of exposure due to poor hygiene and eating and smoking at the workplace, and from swallowed particles that are trapped in sputum.⁽⁴⁹⁾ Studies have documented a strong correlation between airborne lead concentrations with BLL in workers in lead battery factories.^(52,53) Consistent with the BLL data reported in Tables II–IV, Table V indicates that airborne lead exposures in lead battery plants in developing countries average more than 7-fold the $50 \mu\text{g/m}^3$ permissible exposure limit (PEL) as an 8-hr time-weighted average (TWA) established by the U.S. Occupational Safety and Health Administration (OSHA). In addition, the airborne concentrations reported in developing counties in Table V are substantially higher than those reported among lead battery plants in the United States and Italy (range of 21–45 $\mu\text{g/m}^3$).^(54,55) In addition, among the 250 air samples collected by U.S. OSHA in this industry from 2003–2008, the mean and median airborne concentrations were 57 $\mu\text{g/m}^3$ and 28 $\mu\text{g/m}^3$, respectively.⁽⁵⁴⁾

The 10 studies summarized in Table VI suggest that children in communities surrounding lead battery manufacturing and recycling in developing countries had elevated BLLs compared with U.S. children. The average BLL of 29 $\mu\text{g/dL}$ in Table VI is approximately 3-fold higher than the current level of concern for children (10 $\mu\text{g/dL}$) as established by the CDC. In addition, the geometric mean level in Table VI is approximately 13-fold that reported among U.S. children in the most recent National Health and Nutrition Survey (NHANES) data from 2005–2008.⁽⁵⁶⁾ Less dramatic evidence of this pattern was presented in a study conducted in Pakistan that reported a statistically significant higher average BLL in children living closer to lead battery recycling plants compared with controls of similar age living 30 km away (8.1 versus 6.7 $\mu\text{g/dL}$).⁽⁵⁷⁾ It is possible that some of the observed BLLs in these studies were a result of selecting children of employees in these industries with significant contributions due to take-home exposures.

A comparison of the data in Tables II–IV suggests that lead exposures among workers in recycling and reconditioning facilities are higher than those in battery manufacturing operations. In addition, the former operations may also have a greater impact on environmental exposures. Based on the authors' observations, few environmentally sound recycling plants have

been built in developing countries, in part because there are no large-scale collection programs to ensure an adequate battery supply to justify the initial capital investment.

In many areas, battery recycling work is done outdoors. Airborne lead emissions settle on soil and result in widespread environmental contamination.⁽⁵⁸⁾ In addition, the authors observed that many “backyard” recyclers are located in urban areas with high population densities. Wastewater from battery industries and the acid dumped from recycling operations are major sources of lead contamination in the environment. Many urban residents consume untreated water polluted with lead from these industries, which contributes to the overall population exposure.^(57,59)

In considering our finding that the workers in lead battery manufacturing and recycling plants in developing countries have substantially higher BLLs than those in the United States, it is important to consider alternative explanations. First, it is possible the literature was subject to a selection bias in that studies reporting low BLLs were not published because those results were less interesting than studies reporting high BLLs. It is not possible for us to assess the extent of such a publication bias. On the other hand, it is equally plausible that the smallest battery manufacturing/recycling facilities were not surveyed because of the difficulty gaining entry and limitations in sample size, and that these smaller operations had the highest lead exposure levels. This circumstance would

impart a bias toward the null in the BLLs reported in the literature.

Second, we excluded four studies from battery manufacturing plants because of incomplete data. A study by Nusier et al.⁽²⁹⁾ in 2003 reported BLLs by “direct” and “indirect” exposure group in a battery manufacturing facility, but the distinction between these exposure groups was not clear; mean BLL values were 51.4 and 23.9 $\mu\text{g/dL}$, respectively. Two studies that examined differences in BLLs by genotype did not provide summary measures of exposure.⁽³⁰⁾ A study by Bhagwat et al.⁽²⁷⁾ in 2008 provided a combined BLL of 53.6 $\mu\text{g/dL}$ for both battery manufacturing and recycling workers. Because the BLL values in these excluded studies were within the range of reported values in this review, it is unlikely that excluding these studies had much effect on our analysis.

Third, bias may arise due to BLL collection and measurement errors. Samples may be contaminated, as they are generally collected at battery plants. Because BLL tests were conducted in different laboratories in countries without laboratory accreditation or standard reference materials, there is a likelihood of some analytical bias as well as considerable inter- and intra-laboratory variability.⁽⁶⁰⁾ However, we are not able to assess the direction and magnitude of this bias. In the future, measurement error can be minimized by introducing standard reference materials and inter-laboratory blood lead proficiency testing programs in these countries.⁽⁶⁰⁾




FIGURE 1. The Theil-Sen median trend line for arithmetic mean BLLs among battery manufacturing workers for the years 1993 to 2010. The circles correspond to the 61 individual cohort mean BLLs listed in Table II.

An important limitation of our review is that the majority of studies measured a single blood lead level per subject. The BLL is influenced by recent exposure over several weeks as well as long-term exposures through efflux of lead from bone stores.^(61,62) Two or more BLLs measured at different times for each study subject would better reflect the subject's average lead exposure level. On the other hand, in occupational settings with a stable work force and chronic lead exposure, repeated blood lead measurements are likely not highly variable due to the ongoing release of lead into the blood from bone stores.

Similarly, the studies reporting BLLs among children residing near lead battery manufacturing may reflect an episode of acute exposure. Cumulative lead exposures measured in bone would provide the most accurate assessment of lead burden. Unfortunately, there are very few studies with repeated measurements of blood or bone lead in such settings.^(63,64) We note that during the time period covered by our review, leaded gasoline was banned or phased out in most countries, with the exception of a small number of African and some Middle Eastern countries.^(65,66) Lead exposures from gasoline became less significant over this time, and background BLLs declined.

CONCLUSION

The evidence reviewed from studies published on lead battery manufacturing and recycling in developing countries indicates that exposures are considerably higher than in comparable facilities in the United States and the U.K. Average BLLs reported in developing countries were 47 $\mu\text{g}/\text{dL}$ in battery manufacturing plants and 64 $\mu\text{g}/\text{dL}$ in recycling facilities. Airborne lead concentrations reported in lead battery manufacturing and recycling in developing countries were approximately 7-fold greater than those observed in the United States. Children living in proximity to lead battery manufacturing and recycling plants in developing countries had exposures that ranged from 4 to 40 times the geometric mean BLLs for children in the United States. Trend analysis for BLLs in the lead battery manufacturing industry over the time period covered by this review suggests that improvements were being made, but such gains were not statistically significant.

The manufacture and recycling of lead batteries can result in lead exposures sufficient to cause chronic and acute health effects. These lead exposures are not evenly distributed across different societies because substantially higher BLLs have been documented in developing countries than in the United States for the past two decades. At the same time, the lower BLLs among U.S. workers show that high lead exposures in this industry are avoidable without the need to develop any new battery processing, ventilation, or furnace technologies. Unless resources are devoted to reduce lead exposures in battery production and recycling, these activities will lead to increased occupational and environmental lead exposure in the developing world.

The battery industry is the largest consumer of lead and will become a larger consumer as the worldwide demand for automobiles increases and substitutes for lead in other products (e.g., paints) are found. The demand for lead is expected to increase disproportionately in developing countries, including China and India, because of several factors. First, environmentally preferable technologies including solar and wind energy are currently reliant on lead batteries for backup power. In China, 75% of all solar installations are linked to lead batteries.⁽⁶⁷⁾ Second, lead batteries are used in nearly every cell phone tower to provide backup power. Third, as computers are introduced into rural areas to narrow the "digital divide," more lead batteries will be used for backup power.⁽⁶⁸⁾ Consolidation and automation may help reduce the number of workers exposed, but given the fragmented nature of the industry in most developing countries, progress in this respect may be slow.

Fourth, poor manufacturing quality and tropical climates in many developing countries result in a shorter average life of lead batteries and more frequent recycling. Used batteries are broken apart and the metal plates are removed and melted down, often in small fires on the side of a road or in crude smelters, to sell for scrap metal. Because of the poor quality of the material produced by these rudimentary recycling processes, the lead must be melted a second time to remove impurities before it can be used to manufacture new lead batteries.

Many developing countries lack regulations and/or the enforcement capacity to adequately reduce occupational and environmental lead exposures. Alternative means to encourage improvements in the industry should be implemented, including certification programs based on specific occupational, environmental and product stewardship standards verified with independent third-party audits. For example, the Better Environmental Sustainability Targets (BEST) Certification Standard is a comprehensive certification standard developed with involvement of key stakeholders, including industry representatives. The BEST Standard outlines performance measures for workplace exposures, emissions, and extended producer responsibility to take back used batteries for proper recycling.⁽⁶⁹⁾ Companies that meet this standard are eligible to become certified and are permitted to place an eco-label on their products to gain recognition for their efforts. Environmental certifications provide opportunities for companies to expand the market for their products, increase prices, and gain competitive advantage.

ACKNOWLEDGMENTS

This research was supported in part by a generous grant from the Richard and Rhoda Goldman Fund. We would like to thank Shannon Selerowski, Carlos Cabrera, Walter A. Alarcon, and Michael Bates for their assistance.

REFERENCES

- Kosnett, M.J., R.P. Wedeen, S.J. Rothenberg, et al.: Recommendations for medical management of adult lead exposure. *Environ. Health Perspect.* 115(3):463–471 (2007).
- Haefliger, P., M. Mathieu-Nolf, S. Locicero, et al.: Mass lead intoxication from informal used lead-acid battery recycling in Dakar, Senegal. *Environ. Health Perspect.* 117(10):1535–1540 (2009).
- Sanders, T., Y. Liu, V. Buchner, et al.: Neurotoxic effects and biomarkers of lead exposure: A review. *Rev. Environ. Health* 24(1):15–45 (2009).
- Agency for Toxic Substances and Disease Registry (ATSDR): *Toxicological Profile for Lead*. Atlanta: U.S. Department of Health and Human Services, Public Health Service, 2007.
- World Health Organization (WHO): *Inorganic Lead. Environmental Health Criteria* 165. Geneva: WHO, 1995.
- Shih, R.A., H. Hu, M.G. Weisskopf, et al.: Cumulative lead dose and cognitive function in adults: A review of studies that measured both blood lead and bone lead. *Environ. Health Perspect.* 115(3):483–492 (2007).
- Fewtrell, L.J., A. Pruss-Ustun, P. Landrigan, et al.: Estimating the global burden of disease of mild mental retardation and cardiovascular diseases from environmental lead exposure. *Environ. Res.* 94(2):120–133 (2004).
- Weisskopf, M.G., J. Weuve, H. Nie, et al.: Association of cumulative lead exposure with Parkinson's disease. *Environ. Health Perspect.* 118(11):1609–1613 (2010).
- Needleman, H.L., J.A. Riess, M.J. Tobin, et al.: Bone lead levels and delinquent behavior. *JAMA* 275(5):363–369 (1996).
- International Agency for Research on Cancer (IARC): Inorganic and Organic Lead Compounds, Vol. 87. *International Agency for Research on Cancer Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans*. Lyon, France: IARC, 2006.
- Mahaffey, K.R.: Nutrition and lead: Strategies for public health. *Environ. Health Perspect.* 103(Suppl 6):191–196 (1995).
- Ogunseitan, O.A., and T.R. Smith: The cost of environmental lead (Pb) poisoning in Nigeria. *African J. Environ. Sci. Technol.* 1(2):027–036 (2007).
- Bloom, D.E., and D. Canning: The health and wealth of nations. *Science* 287(5456):1207–1209 (2000).
- International Lead and Zinc Study Group: "End Uses of Lead." [Online] Available at <http://www.ilszg.org/static/enduses.aspx?from=2> (Accessed January 12, 2011).
- IC Consultants Ltd.: *Lead: The Facts*. London: IC Consultants Ltd., 2001. pp. 65–71.
- Chen, H.Y., A.J. Li, and D.E. Finlow: The lead and lead-acid battery industries during 2002 and 2007 in China. *J. Power Sources* 191(1):22–27 (2009).
- Chanjaroen, C.: "Zinc Gains on Speculation Lower Prices Will Force Mine Closures." *Bloomberg News* (2008, June 18).
- Mao, J.S., J. Dong, and T.E. Graedel: The multilevel cycle of anthropogenic lead II. Results and discussion. *Resour. Conserv. Recycl.* 52(8–9):1050–1057 (2008).
- "Lead in July 2010." [Online] Available at <http://minerals.usgs.gov/minerals> (Accessed November 18, 2010).
- "Waste & Scrap of Primary Cells, Primary Batteries and Electric Accumulators, Spent Primary Cells, Spent Primary Batteries & Spent Electric Accumulators" [Online] Available at <http://comtrade.un.org/db/dqBasicQueryResults.aspx?cc=854810&px=HS&r=842&y=2009,2008,2007,2006,2005&p=699,%20484,%20410&rg=2&so=8> (Accessed November 18, 2010).
- Hur, J., and I. Suzuki: "Lead-Battery Demand for Cars to Rise 2.6% on China, India." [Online] Available at <http://www.businessweek.com/news/2011-02-25/lead-battery-demand-for-cars-to-rise-2-6-on-china-india.html> (Accessed April 7, 2011).
- "Battery Industry: Consolidation Required to Boost Efficiency." [Online] Available at <http://bit.ly/gHmDV0> (Accessed April 7, 2011).
- Chaowei Power Holdings Limited: "Global Offerings." Zhicheng, Changxing: Chaowei Power Holdings Limited, 2010.
- Wananukul, W., T. Sura, and P. Salaitanawatwong: Polymorphism of delta-aminolevulinic acid dehydratase and its effect on blood lead levels in Thai workers. *Arch. Environ. Occup. Health* 61(2):67–72 (2006).
- Heo, Y., B.K. Lee, K.D. Ahn, et al.: Serum IgE elevation correlates with blood lead levels in battery manufacturing workers. *Hum. Exp. Toxicol.* 23(5): 209–213 (2004).
- Shaik, A.P., and K. Jamil: Individual susceptibility and genotoxicity in workers exposed to hazardous materials like lead. *J. Hazard Mater.* 168(2–3):918–924 (2009).
- Bhagwat, V.R., A.J. Patil, J.A. Patil, et al.: Occupational lead exposure and liver functions in battery manufacture workers around Kolhapur (Maharashtra). *Al Ameen J. Med. Sci.* 1(1):2–9 (2008).
- Ramirez, A.V.: [Workers' lead exposure in informal batteries factories.] *An. Fac. Med.* 69(2):104–107 (2008). [In Spanish]
- Nusier, M.K., Z.J. El-Akawi, and I. Bani-Hani: Blood lead levels and gamma-glutamyl transferase activity in battery factory workers in Jordan. *Saudi Med. J.* 24(9):941–944 (2003).
- Xi-Biao, Y., W. Cui-E, F. Hua, et al.: Associations of blood lead levels, kidney function, and blood pressure with delta-aminolevulinic acid dehydratase and vitamin D receptor gene polymorphisms. *Toxicol. Mech. Methods* 13(2):139–146 (2003).
- Ravichandran, B., K. Ravibabu, S. Raghavan, et al.: Environmental and biological monitoring in a lead acid battery manufacturing unit in India. *J. Occup. Health* 47(4):350–353 (2005).
- Department of Health, Education, and Welfare (DHEW), Public Health Service (PHS), Centers for Disease Control (CDC), National Institute for Occupational Safety and Health (NIOSH): *Occupational Exposure Sampling Strategy Manual*. DHEW (NIOSH) 77-173, by N.A. Leidel, K.A. Busch, and J.R. Lynch. Cincinnati, Ohio: DHEW, PHS, CDC, NIOSH, 1977.
- Muntner, P., A. Menke, K.B. DeSalvo, et al.: Continued decline in blood lead levels among adults in the United States: The National Health and Nutrition Examination Surveys. *Arch. Intern. Med.* 165(18):2155–2161 (2005).
- Newson, R.: Confidence intervals for rank statistics: Percentile slopes, differences, and ratios. *Stata J.* 6(4):497–520 (2006).
- Theil, H.: A rank-invariant method of linear and polynomial regression analysis. III. *Proceedings of Koninklijke Nederlandse Akademie van Wetenschappen Part 3(A.53)*:1397–1412 (1950).
- Sen, P.K.: Estimates of the regression coefficient based on Kendall's Tau. *J. Am. Stat. Assoc.* 63:1379–1389 (1968).
- U.S. Department of the Interior, U.S. Geological Survey: *Statistical Methods in Water Resources Techniques of Water Resources Investigations, Book 4*, by D.R. Helsel and R.M. Hirsch. Reston, Va.: USGS National Center, 2002.
- Ye, X., and O. Wong: Lead exposure, lead poisoning, and lead regulatory standards in China, 1990–2005. *Regul. Toxicol. Pharmacol.* 46(2):157–162 (2006).
- ACGIH: *Threshold Limit Values for Chemical Substances and Physical Agents and Biological Exposure Indices*. Cincinnati, Ohio: ACGIH, 2004.
- Centers for Disease Control and Prevention (CDC): *Guidelines for the Identification and Management of Lead Exposure in Pregnant and Lactating Women*, A.S. Ettinger and A.G. Wengrovitz (eds.). Atlanta, Ga.: National Center for Environmental Health/Division of Emergency and Environmental Health Services, 2010.
- U.S. Department of Energy (DOE): *Environmental Assessment for Exide Technologies Electric Drive Vehicle Battery and Component Manufacturing Initiative Application, Bristol, TN, and Columbus, GA*. Morgantown, W.Va.: DOE National Energy Technology Laboratory, 2010.
- Health and Safety Executive (HSE): *Exposure to Lead at Work Blood-Lead Levels of Workers Exposed to Lead at Work in Great Britain 1996/97–2000/2001*. National Statistics. London: HSE, 2002.

43. **Health and Safety Executive (HSE):** *Occupational Health Statistics Bulletin 2003/04*. Bootle, Merseyside, U.K.: HSE, Epidemiology and Medical Statistic Unit, HSE, 2004. pp. 1–14.
44. **California Department of Health Services, Occupational Health Branch:** *Blood Lead Levels in California Workers 1995–1999, Report of the California Occupational Blood Lead Registry*. Sacramento, Calif.: California Department of Public Health, 2002. pp. 42–46.
45. **Washington State Department of Labor and Industries (DLI):** *Surveillance for Occupational Lead Poisoning, 1993–2001*, by S.G. Whittaker and C.C. Curwick (Technical Report 44-3-2001). Olympia, Wash.: Washington State Department of Labor and Industries, 2001.
46. “Adult Blood Lead Epidemiology and Surveillance (ABLES).” [Online] Available at <http://www.cdc.gov/niosh/topics/ABLES/ables.html> (Accessed November 23, 2010).
47. “2007 Economic Census, Sector 31: EC0731A1: Manufacturing: Geographic Area Series: Industry Statistics for the States, Metropolitan and Micropolitan Statistical Areas, Counties, and Places.” [Online] Available at <http://bit.ly/ei4Zlh> (Accessed February 6, 2011).
48. “Protecting People: Lead-Acid Battery Safety.” [Online] Available at <http://www.batterycouncil.org/LeadAcidBatteries/EnvironmentalRegulations/LeadAcidBatterySafety/tabid/96/Default.aspx> (Accessed January 12, 2011).
49. “Battery Manufacturing.” [Online] Available at <http://www.osha.gov/SLTC/batterymanufacturing/index.html> (Accessed April 4, 2011).
50. **Dyosi, S.:** Evaluation of preventive and control measures for lead exposure in a South African lead-acid battery recycling smelter. *J. Occup. Environ. Hyg.* 4:762–769 (2007).
51. **Centers for Disease Control (CDC):** Occupational and environmental lead poisoning associated with battery repair shops: Jamaica. *MMWR* 38(27):474–481 (1989).
52. **Ibiebele, D.D.:** Air and blood lead levels in a battery factory. *Sci. Total Environ.* 152(3):269–273 (1994).
53. **Wang, V.-S., M.-T. Lee, J.-Y. Chiou, et al.:** Relationship between blood lead levels and renal function in lead battery workers. *Int. Arch. Occup. Environ. Health* 75(8):569–575 (2002).
54. **Henn, S.A., A.L. Sussell, J. Li, et al.:** Characterization of lead in US workplaces using data from OSHA’s integrated management information system. *Am. J. Ind. Med.* 54(5):356–365 (2011).
55. **Fenga, C., A. Cacciola, L.B. Martino, et al.:** Relationship of blood lead levels to blood pressure in exhaust battery storage workers. *Ind. Health* 44(2):304–309 (2006).
56. “Table B1: Concentrations of Lead in the Blood of Children Ages 5 and under.” [Online] Available at http://www.epa.gov/envirohealth/children/body_burdens/b1-table.html (Accessed May 19, 2011)
57. **Khan, D.A., S. Qayyum, S. Saleem, et al.:** Lead exposure and its adverse health effects among occupational worker’s children. *Toxicol. Ind. Health* 26(8):497–504 (2010).
58. **Renberg, I., M. L. Brännvall, R. Bindler, et al.:** Atmospheric lead pollution history during four millennia (2000 BC to 2000 AD) in Sweden. *AMBIO* 29(3):150–156 (2000).
59. **Raviraja, A., G.N. Babu, A.R. Bijoor, et al.:** [Lead toxicity in a family as a result of occupational exposure.] *Arh. Hig. Rada. Toksikol.* 59(2):127–133 (2008). [In Croatian, English]
60. **Pidetcha, P., L. Suwanthorn, A. Preechavuth, et al.:** Interlaboratory results for blood lead proficiency testing program in Thailand. *J. Med. Assoc. Thai.* 82(12):1247–1253 (1999).
61. **Hu, H., R. Shih, S. Rothenberg, et al.:** The epidemiology of lead toxicity in adults: Measuring dose and consideration of other methodologic issues. *Environ. Health Perspect.* 115(3):455–462 (2007).
62. **Menke, A., P. Muntner, V. Batuman, et al.:** Blood lead below 0.48 micromol/L (10 microg/dL) and mortality among US adults. *Circulation* 114(13):1388–1394 (2006).
63. **Chuang, H.Y., M.L.T. Lee, K.Y. Chao, et al.:** Relationship of blood lead levels to personal hygiene habits in lead battery workers: Taiwan, 1991–1997. *Am. J. Ind. Med.* 35(6):595–603 (1999).
64. **Hsiao, C.Y., H.D. Wu, J.S. Lai, et al.:** A longitudinal study of the effects of long-term exposure to lead among lead battery factory workers in Taiwan (1989–1999). *Sci. Total Environ.* 279(1–3):151–158 (2001).
65. **United Nations Environment Programme (UNEP):** *Target 2008: Global Elimination of Leaded Petrol*. Nairobi: UNEP Partnership for Clean Fuels and Vehicles (PCFV) 2008.
66. **Organisation for Economic Co-Operation and Development (OECD), U.N. Environment Programme (UNEP):** “Phasing Lead out of Gasoline: An Examination of Policy Approaches in Different Countries”. Paris: OECD, UNEP, 1999.
67. **Chang, Y., X. Mao, Y. Zhao, et al.:** Lead-acid battery use in the development of renewable energy systems in China. *J. Power Sources* 191(1):176–183 (2009).
68. **Cherry, C.R., and P. Gottesfeld:** Plans to distribute the next billion computers by 2015 creates lead pollution risk. *J. Cleaner Production* 17:1620–1628 (2009).
69. “Better Environmental Sustainability Targets (BEST) For Lead Battery Manufacturers.” [Online] Available at <http://www.okinternational.org/lead-batteries/Background> (Accessed April 4, 2011).
70. **Zhang, J.:** Investigation and evaluation of zinc protoporphyrin as a diagnostic indicator in lead-intoxication. *Am. J. Ind. Med.* 24(6):707–712 (1993).
71. **Far, H.S., N.T. Pin, C.Y. Kong, et al.:** An evaluation of the significance of mouth and hand contamination for lead absorption in lead-acid-battery workers. *Int. Arch. Occup. Environ. Health* 64(6):439–443 (1993).
72. **Makino, S., K. Matsuno, N. Hisanaga, et al.:** Medical examination of workers exposed to lead in the Philippines. *Jap. J. Ind. Health* 36(2):115–123 (1994).
73. **Singh, B., D. Dhawan, B. Nehru, et al.:** Impact of lead pollution on the status of other trace-metals in blood and alterations in hepatic functions. *Biol. Trace Elem. Res.* 40(1):21–29 (1994).
74. **Kim, Y., K. Harada, S. Ohmori, et al.:** Evaluation of lead exposure in workers at a lead-acid battery factory in Korea: With focus on activity of erythrocyte pyrimidine 5'-nucleotidase (P5N). *Occup. Environ. Med.* 52(7):484–488 (1995).
75. **Schwartz, B.S., B.K. Lee, W. Stewart, et al.:** Associations of delta-aminolevulinic acid dehydratase genotype with plant, exposure duration, and blood lead and zinc protoporphyrin levels in Korean lead workers. *Am. J. Epidemiol.* 142(7):738–745 (1995).
76. **Solliway, B.M., A. Schaffer, H. Pratt, et al.:** Effects of exposure to lead on selected biochemical and haematological variables. *Pharmacol. Toxicol.* 78(1):18–22 (1996).
77. **Cordeiro, R., E.C. Lima, P.E.T. Salgado, et al.:** Neurological disturbances in workers with low levels of lead in the blood. II. Neuropsychological disorders. *Rev. Saude Publica* 30(4):358–363 (1996).
78. **Chang, H.R., S.S. Chen, T.J. Chen, et al.:** Lymphocyte beta(2)-adrenergic receptors and plasma catecholamine levels in lead-exposed workers. *Toxicol. Appl. Pharmacol.* 139(1):1–5 (1996).
79. **Lai, J.-S., T.N. Wu, S.-H. Liou, et al.:** A study of the relationship between ambient lead and blood lead among lead battery workers. *Int. Arch. Occup. Environ. Health* 69(4):295–300 (1997).
80. **Yucesoy, B., A. Turhan, S. Mirshahidi, et al.:** Effects of high-level exposure to lead on NK cell activity and T-lymphocyte functions in workers. *Hum. Exper. Toxicol.* 16(6):311–314 (1997).
81. **Schwartz, B.S., B.K. Lee, W. Stewart, et al.:** delta-Aminolevulinic acid dehydratase genotype modifies four hour urinary lead excretion after oral administration of dimercaptosuccinic acid. *Occup. Environ. Med.* 54(4):241–246 (1997).
82. **Kuo, H.W., C.S. Wang, and J.S. Lai:** Semen quality in workers with long-term lead exposure: A preliminary study in Taiwan. *Sci. Total Environ.* 204(3):289–292 (1997).
83. **Ho, S.F., C.T. Sam, and G. Bin Embi:** Lead exposure in the lead-acid storage battery manufacturing and PVC compounding industries. *Occup. Med. (Lond.)* 48(6):369–373 (1998).
84. **Ehrlich, R., T. Robins, E. Jordaan, et al.:** Lead absorption and renal dysfunction in a South African battery factory. *Occup. Environ. Med.* 55(7):453–460 (1998).

85. **Vaglenov, A., E. Carbonell, and R. Marcos:** Biomonitoring of workers exposed to lead. Genotoxic effects, its modulation by polyvitamin treatment and evaluation of the induced radioresistance. *Mutat. Res.* 418(2–3):79–92 (1998).
86. **Williams, G., L. Hall, and J. Addae:** Increase in hair lead, but not blood lead content of occupationally exposed workers. *Environ. Geochem. Health* 20(4):239–243 (1998).
87. **Udeger, U., and N. Basaran:** Effects of lead on neutrophil functions in occupationally exposed workers. *Environ. Toxicol. Pharmacol.* 5(2):113–117 (1998).
88. **Froom, P., E. Kristal-Boneh, J. Benbassat, et al.:** Lead exposure in battery-factory workers is not associated with anemia. *J. Occup. Environ. Med.* 41(2):120–123 (1999).
89. **Caldeira, C., R. Mattos, A. Meyer, et al.:** [Limits in the applicability of urine delta aminolevulinic acid determination as a screening test in the evaluation of occupational lead poisoning.] *Cad. Saude Publica* 16(1):225–230 (2000). [In Portuguese]
90. **Restrepo, H.G., D. Sicard, and M.M. Torres:** DNA damage and repair in cells of lead exposed people. *Am. J. Ind. Med.* 38(3):330–334 (2000).
91. **Roh, Y.M., K. Kim, and H. Kim:** Zinc protoporphyrin IX concentrations between normal adults and the lead-exposed workers measured by HPLC, spectrofluorometer, and hematofluorometer. *Ind. Health* 38(4):372–379 (2000).
92. **Basaran, N., and U. Udeger:** Effects of lead on immune parameters in occupationally exposed workers. *Am. J. Ind. Med.* 38(3):349–354 (2000).
93. **Ratzon, N., P. Froom, E. Leikin, et al.:** Effect of exposure to lead on postural control in workers. *Occup. Environ. Med.* 57(3):201–203 (2000).
94. **Chuang, H.Y., J. Schwartz, S.Y. Tsai, et al.:** Vibration perception thresholds in workers with long term exposure to lead. *Occup. Environ. Med.* 57(9):588–594 (2000).
95. **Duydu, Y., H.S. Suzen, A. Aydin, et al.:** Correlation between lead exposure indicators and sister chromatid exchange (SCE) frequencies in lymphocytes from inorganic lead exposed workers. *Arch. Environ. Contam. Toxicol.* 41(2):241–246 (2001).
96. **Vaglenov, A., A. Creus, S. Laltchev, et al.:** Occupational exposure to lead and induction of genetic damage. *Environ. Health Perspect.* 109(3):295–298 (2001).
97. **Hwang, Y.H., C.W. Chang, K.Y. Chao, et al.:** Using structural equation model to explore occupational lead exposure pathways. *Sci. Total Environ.* 284(1–3):95–108 (2002).
98. **Sonmez, F., O. Donmez, H.M. Sonmez, et al.:** Lead exposure and urinary N-acetyl beta D glucosaminidase activity in adolescent workers in auto repair workshops. *J. Adolesc. Health* 30(3):213–216 (2002).
99. **Mishra, K.P., V.K. Singh, R. Rani, et al.:** Effect of lead exposure on the immune response of some occupationally exposed individuals. *Toxicology* 188(2–3):251–259 (2003).
100. **Palus, J., K. Rydzynski, E. Dziubaltowska, et al.:** Genotoxic effects of occupational exposure to lead and cadmium. *Mutat. Res.* 540(1):19–28 (2003).
101. **Süzen, H., Y. Duydu, A. Aydin, et al.:** Influence of the delta-aminolevulinic acid dehydratase (ALAD) polymorphism on biomarkers of lead exposure in Turkish storage battery manufacturing workers. *Am. J. Ind. Med.* 43(2):165–171 (2003).
102. **Pizent, A., J. Jurasović, and S. Telišman:** Serum calcium, zinc, and copper in relation to biomarkers of lead and cadmium in men. *J. Trace Elem. Med. Biol.* 17(3):199–205 (2003).
103. **Duydu, Y., and H.S. Suzen:** Influence of delta-aminolevulinic acid dehydratase (ALAD) polymorphism on the frequency of sister chromatid exchange (SCE) and the number of high-frequency cells (HFCs) in lymphocytes from lead-exposed workers. *Mutat. Res.* 540(1):79–88 (2003).
104. **Lormphongs, S., K. Miyashita, I. Morioka, et al.:** Lead exposure and blood lead level of workers in a battery manufacturing plant in Thailand. *Ind. Health* 41(4):348–353 (2003).
105. **Gurer-Orhan, H., H.U. Sabir, and H. Ozgunes:** Correlation between clinical indicators of lead poisoning and oxidative stress parameters in controls and lead-exposed workers. *Toxicology* 195(2–3):147–154 (2004).
106. **Chuang, H.Y., K.T. Yu, C.K. Ho, et al.:** Investigations of vitamin D receptor polymorphism affecting workers' susceptibility to lead. *J. Occup. Health* 46(4):316–322 (2004).
107. **Bage, C., A.I. Bozkurt, E.A. Cakmak, et al.:** Blood lead levels of the battery and exhaust workers and their pulmonary function tests. *Int. J. Clin. Pract.* 58(6):568–572 (2004).
108. **Duydu, Y., A. Dur, and H.S. Süzen:** Evaluation of increased proportion of cells with unusually high sister chromatid exchange counts as a cytogenetic biomarker for lead exposure. *Biol. Trace Elem. Res.* 104(2):121–129 (2005).
109. **Chuang, H.Y., K.Y. Chao, and J.D. Wang:** Estimation of burden of lead for offspring of female lead workers: A quality-adjusted life year (QALY) assessment. *J. Toxicol. Environ. Health A* 68(17–18):1485–1496 (2005).
110. **Karakaya, A.E., E. Ozcagli, N. Ertas, et al.:** Assessment of abnormal DNA repair responses and genotoxic effects in lead exposed workers. *Am. J. Ind. Med.* 47(4):358–363 (2005).
111. **Patil, A.J., V.R. Bhagwat, J.A. Patil, et al.:** Effect of lead (Pb) exposure on the activity of superoxide dismutase and catalase in battery manufacturing workers (BMW) of Western Maharashtra (India) with reference to heme biosynthesis. *Int. J. Environ. Res. Public Health* 3(4):329–337 (2006).
112. **Kuruvilla, A., V.V. Pillay, P. Adhikari, et al.:** Clinical manifestations of lead workers of Mangalore, India. *Toxicol. Ind. Health* 22(9):405–413 (2006).
113. **Li, W.F., M.H. Pan, M.C. Chung, et al.:** Lead exposure is associated with decreased serum paraoxonase 1 (PON1) activity and genotypes. *Environ. Health Perspect.* 114(8):1233–1236 (2006).
114. **Engin, A.B., D. Tuzun, and G. Sahin:** Evaluation of pteridine metabolism in battery workers chronically exposed to lead. *Hum. Exp. Toxicol.* 25(7):353–359 (2006).
115. **Chen, Z.J., J.L. Lou, S.J. Chen, et al.:** Evaluating the genotoxic effects of workers exposed to lead using micronucleus assay, comet assay and TCR gene mutation test. *Toxicology* 223(3):219–226 (2006).
116. **Lin, T.A., and J. Tai-yi:** Benchmark dose approach for renal dysfunction in workers exposed to lead. *Environ. Toxicol.* 22(3):229–233 (2007).
117. **Chia, S.E., S.M. Ali, B.L. Lee, et al.:** Association of blood lead and homocysteine levels among lead exposed subjects in Vietnam and Singapore. *Occup. Environ. Med.* 64(10):688–693 (2007).
118. **Chuang, H., W. Cheng, C. Chen, et al.:** A follow-up comparison of blood lead levels between foreign and native workers of battery manufacturing in Taiwan. *Sci. Total Environ.* 394(1):52–56 (2008).
119. **Kasuba, V., R. Rozgaj, M. Milić, et al.:** Evaluation of lead exposure in battery-manufacturing workers with focus on different biomarkers. *J. Appl. Toxicol.* 30(4):321–328 (2009).
120. **Hsu, P.C., H.Y. Chang, Y.L. Guo, et al.:** Effect of smoking on blood lead levels in workers and role of reactive oxygen species in lead-induced sperm chromatin DNA damage. *Fertil. Steril.* 91(4):1096–1103 (2009).
121. **Nsheiwat, Z., E. Kozerb, M. Berkovitch, et al.:** A comparison of lead exposures in industrial "hot spots" in Jordan and Israel. *Toxicol. Environ. Chem.* 92(4):669–682 (2010).
122. **Gao, A., X.-t. Lu, Q.-y. Li, et al.:** Effect of the delta-aminolevulinic acid dehydratase gene polymorphism on renal and neurobehavioral function in workers exposed to lead in China. *Sci. Total Environ.* 408(19):4052–4055 (2010).
123. **Dos Santos, A.C., S. Colacciopo, C.M. Dal Bo, et al.:** Occupational exposure to lead, kidney function tests, and blood pressure. *Am. J. Ind. Med.* 26(5):635–643 (1994).
124. **Chia, S.E., H.P. Chia, C.N. Ong, et al.:** Cumulative blood lead levels and nerve conduction parameters. *Occup. Med. (Lond.)* 46(1):59–64 (1996).

125. Chia, S.E., H.P. Chia, C.N. Ong, et al.: Cumulative blood lead levels and neurobehavioral test performance. *Neurotoxicology* 18(3):793–803 (1997).
126. Sithisarankul, P., B.S. Schwartz, B.K. Lee, et al. Aminolevulinic acid dehydratase genotype mediates plasma levels of the neurotoxin, 5-aminolevulinic acid, in lead-exposed workers. *Am. J. Ind. Med.* 32(1):15–20 (1997).
127. Chia, S.E., K.S. Chia, H.P. Chia, et al.: Three-year follow-up of serial nerve conduction among lead-exposed workers. *Scand. J. Work Environ. Health* 22(5):374–380 (1996).
128. Bergdahl, I.A., M. Sheveleva, A. Schutz, et al.: Plasma and blood lead in humans: Capacity-limited binding to delta-aminolevulinic acid dehydratase and other lead-binding components. *Toxicol. Sci.* 46(2):247–253 (1998).
129. Jakubowski, M., M. Trzcinka-Ochocka, G. Razniewska, et al.: Blood lead levels in industrial workers in Poland. *Int. J. Occup. Med. Environ. Health* 11(1):59–67 (1998).
130. Sun, Y., D. Sun, Z. Zhou, et al.: Osteoporosis in a Chinese population due to occupational exposure to lead. *Am. J. Ind. Med.* 51(6):436–442 (2008).
131. Yeh, J.H., Y.C. Chang, and J.D. Wang: Combined electroneurographic and electromyographic studies in lead workers. *Occup. Environ. Med.* 52(6):415–419 (1995).
132. Wang, J.D., W.T. Soong, K.Y. Chao, et al.: Occupational and environmental lead poisoning: Case study of a battery recycling smelter in Taiwan. *J. Toxicol. Sci.* 23(Suppl 2):241–245 (1998).
133. Corzo, G., and R. Naveda: [Occupational exposure to lead in production units in Maracaibo, Venezuela.] *Invest. Clin.* 39(3):163–173 (1998). [In Spanish]
134. Mehdi, J., F. Al-Imarah, and A. Al-Suhail: Levels of some trace metals and related enzymes in workers at storage-battery factories in Iraq. *East. Mediterr. Health J.* 6(1):66–82 (2000).
135. Suplido, M.L., and C.N. Ong: Lead exposure among small-scale battery recyclers, automobile radiator mechanics, and their children in Manila, the Philippines. *Environ. Res.* 82(3):231–238 (2000).
136. Cardenas-Bustamante, O., M.E. Varona-Urbe, S.M. Nunez-Trujillo et al.: [Correlation of protoporphyrin zinc with lead in blood in workers of the car batteries industry in Bogota, Colombia]. *Salud. Publica Mex.* 43(3):203–210 (2001). [In Spanish]
137. Kumar, P., S.G. Husain, R.C. Murthy, et al.: Neuropsychological studies on lead battery workers. *Vet. Hum. Toxicol.* 44(2):76–78 (2002).
138. Ayatollahi, M.: Study of the impact of blood lead level on humoral immunity in humans. *Toxicol. Ind. Health* 18(1):39–44 (2002).
139. Minozzo, R., E.L. Minozzo, L.I. Deimling, et al.: [Blood lead levels in the battery recycling industry of the metropolitan region of Porto Alegre, RS.] *J. Bras. Patol. Med. Lab.* 44(6):407–412 (2008). [In Portuguese]
140. Jiang, Y.M., L.L. Long, X.Y. Zhu, et al.: Evidence for altered hippocampal volume and brain metabolites in workers occupationally exposed to lead: A study by magnetic resonance imaging and H-1 magnetic resonance spectroscopy. *Toxicol. Lett.* 181(2):118–125 (2008).
141. Peter, A.O.: Total blood and urinary lead levels in battery charging artisans in two metropolitan cities of South West Nigeria. *J. Appl. Biosci.* 14:796–799 (2009).
142. Hwang, Y., K. Chao, C. Chang, et al.: Lip lead as an alternative measure for lead exposure assessment of lead battery assembly workers. *Am. Ind. Hyg. Assoc. J.* 61(6):825–831 (2000).
143. Donguk, P., and P. Namwon: Exposure to lead and its particles size distribution. *J. Occup. Health* 46(3):225–229 (2004).
144. Morales Bonilla, C., and E.A. Mauss: A community-initiated study of blood lead levels of Nicaraguan children living near a battery factory. *Am. J. Public Health* 88(12):1843–1845 (1998).
145. Kaul, B., and H. Mukerjee: Elevated blood lead and erythrocyte protoporphyrin levels of children near a battery-recycling plant in Haina, Dominican Republic. *Int. J. Occup. Environ. Health* 5(4):307–312 (1999).
146. Saraci, M., and K. Ziegler-Skylakakis: Determination of lead in the blood of children in the town of Berat, Albania. *Chemosphere* 39(4):689–696 (1999).
147. Tabaku, A., V. Bizgha, and S.I. Rahlenbeck: Biological monitoring of lead exposure in high risk groups in Berat, Albania. *J. Epidemiol. Community Health* 52(4): 234–236 (1998).
148. Cortes-Maramba, N.P., L.R. Panganiban, E.S. Castillo, et al.: Health and environmental assessment among residents of a community near a battery recycling plant. *J. Phys. IV (Proceedings)* 107:809–821 (2003).
149. Safi, J., A. Fischbein, S. El Haj, et al.: Childhood lead exposure in the Palestinian Authority, Israel, and Jordan: Results from the Middle Eastern Regional Cooperation Project, 1996–2000. *Environ. Health Perspect.* 114(6):917–922 (2006).
150. De Freitas, C.U., E.M. De Capitani, N. Gouveia, et al.: Lead exposure in an urban community: Investigation of risk factors and assessment of the impact of lead abatement measures. *Environ. Res.* 103(3):338–344 (2007).
151. Ahmad, T., A. Mumtaz, D. Ahmad, et al.: Lead exposure in children living around the automobile and battery repair workshops. *Biomedica* 25(July–December):128–132 (2009).