

Санкт-Петербургский политехнический университет Петра
Великого
Институт компьютерных наук и технологий
Высшая школа программной инженерии

Курсовая Работа

по дисциплине
«Математические модели»

Выполнил:
Группа:

Ферапонтов М.В.
гр. 3530904/00104

Проверил:

Воскобойников С. П.

Санкт-Петербург
2023

Содержание

1	Вступление	2
1.1	Постановка задачи	2
2	Разностная схема	3
2.1	Внутренние точки	3
2.2	На левой границе	4
2.3	На правой границе	5
2.4	На нижней границе	6
2.5	На верхней границе	7
2.6	Левый-нижний угол	7
2.7	Левый-верхний угол	8
2.8	Правый-верхний угол	8
2.9	Правый-нижний угол	9
3	Невязка разностной схемы	10
3.1	Невязка во внутренних точках	10
3.2	Невязка на левой границе	12
4	Запись СЛАУ	13
4.1	Запись для внутренних точек	13
4.2	Запись для левой границы	14
4.3	Запись для правой границы	15
4.4	Запись для нижней границы	15
4.5	Запись для верхней границы	16
4.6	Запись для левой нижней граничной точки	17
4.7	Запись для правой нижней граничной точки	17
5	Заключение	19

1 Вступление

1.1 Постановка задачи

Вариант N7. Используя интегро-интерполяционный метод, разработать подпрограмму для моделирования распределения температуры в цилиндре, описываемого математической моделью

$$-\left[\frac{1}{r}\frac{\partial}{\partial r}\left(rk_1(r,z)\frac{\partial u}{\partial r}\right)+\frac{\partial}{\partial z}\left(k_2(r,z)\frac{\partial u}{\partial z}\right)\right]=f(r,z)$$

$$0 \leq c_{11} \leq k_1(r,z) \leq c_{12}, \quad 0 \leq c_{11} \leq k_2(r,z) \leq c_{22}, \quad 0 \leq r \leq R, \quad 0 \leq z \leq L$$

С граничными условиями:

$$\begin{array}{ll} u|_{r=0} - \text{ограничено} & -k_1 \frac{\partial u}{\partial r} \Big|_{r=R} = \chi_2 u|_{r=R} - \varphi_2(z) \\ k_2 \frac{\partial u}{\partial z} \Big|_{z=0} = \chi_3 u|_{z=0} - \varphi_3(r) & u|_{z=L} = \varphi_4(r) \\ \chi_2 \geq 0 & \chi_3 \geq 0 \end{array}$$

Матрица алгебраической системы должна храниться в упакованной форме.

2 Разностная схема

Введем основную сетку:

N_r — число разбиений на $[0, R]$

$$r_0 < r_1 < \dots < r_N$$

$$r_0 = 0, \quad r_N = R$$

$$h_r = \frac{R - 0}{N_r}$$

N_z — число разбиений на $[0, L]$

$$z_0 < z_1 < \dots < z_N$$

$$z_0 = 0, \quad z_N = L$$

$$h_z = \frac{L - 0}{N_z}$$

Введем дополнительную сетку:

$$r_{i-\frac{1}{2}} = \frac{r_i + r_{i-1}}{2} \quad i = 1, \dots, N_r$$

$$h_i = \begin{cases} \frac{h_r}{2}, & i = 0 \\ h_r, & i = 1, 2, \dots, N_r - 1 \\ \frac{h_r}{2}, & i = N_r \end{cases}$$

$$z_{j-\frac{1}{2}} = \frac{z_j + z_{j-1}}{2} \quad j = 1, \dots, N_z$$

$$h_j = \begin{cases} \frac{h_z}{2}, & j = 0 \\ h_z, & j = 1, 2, \dots, N_z - 1 \\ \frac{h_z}{2}, & j = N_z \end{cases}$$

Преобразуем наше начальное уравнение домножив на r

$$- \left[\frac{\partial}{\partial r} \left(rk_1(r, z) \frac{\partial u}{\partial r} \right) + \frac{\partial}{\partial z} \left(rk_2(r, z) \frac{\partial u}{\partial z} \right) \right] = rf(r, z)$$

2.1 Внутренние точки

Проинтегрируем уравнение внутри интервала:

$$- \int_{r_{i-\frac{1}{2}}}^{r_{i+\frac{1}{2}}} \int_{z_{j-\frac{1}{2}}}^{z_{j+\frac{1}{2}}} \left[\frac{\partial}{\partial r} \left(rk_1(r, z) \frac{\partial u}{\partial r} \right) + \frac{\partial}{\partial z} \left(rk_2(r, z) \frac{\partial u}{\partial z} \right) \right] dr dz = \int_{r_{i-\frac{1}{2}}}^{r_{i+\frac{1}{2}}} \int_{z_{j-\frac{1}{2}}}^{z_{j+\frac{1}{2}}} rf(r, z) dr dz$$

Получим:

$$\begin{aligned} & - \left[\int_{z_{j-\frac{1}{2}}}^{z_{j+\frac{1}{2}}} rk_1(r, z) \frac{\partial u}{\partial r} \Big|_{r=r_{i+\frac{1}{2}}} dz - \int_{z_{j-\frac{1}{2}}}^{z_{j+\frac{1}{2}}} rk_1(r, z) \frac{\partial u}{\partial r} \Big|_{r=r_{i-\frac{1}{2}}} dz \right. \\ & \left. + \int_{r_{i-\frac{1}{2}}}^{r_{i+\frac{1}{2}}} rk_2(r, z) \frac{\partial u}{\partial z} \Big|_{z=z_{j+\frac{1}{2}}} dr - \int_{r_{i-\frac{1}{2}}}^{r_{i+\frac{1}{2}}} rk_2(r, z) \frac{\partial u}{\partial z} \Big|_{z=z_{j-\frac{1}{2}}} dr \right] = \int_{r_{i-\frac{1}{2}}}^{r_{i+\frac{1}{2}}} \int_{z_{j-\frac{1}{2}}}^{z_{j+\frac{1}{2}}} rf(r, z) dr dz \end{aligned}$$

Воспользуемся формулами численного дифференцирования:

$$k_1(r, z) \frac{\partial u}{\partial r} \Big|_{r=r_{i-\frac{1}{2}}} \approx k_1(r_{i-\frac{1}{2}}, z) \frac{v_{i,j} - v_{i-1,j}}{h_r}$$

$$k_2(r, z) \frac{\partial u}{\partial r} \Big|_{z=z_{j-\frac{1}{2}}} \approx k_2(r, z_{j-\frac{1}{2}}) \frac{v_{i,j} - v_{i,j-1}}{h_z}$$

Также воспользуемся формулой средних прямоугольников:

$$\int_{r_{i-\frac{1}{2}}}^{r_{i+\frac{1}{2}}} r \varphi(r, z) dr = \bar{h}_i r_i \varphi_i$$

$$\int_{r_{i-\frac{1}{2}}}^{r_{i+\frac{1}{2}}} \int_{z_{j-\frac{1}{2}}}^{z_{j+\frac{1}{2}}} r \varphi(r, z) dr dz = \bar{h}_i \bar{h}_j r_i \varphi_{i,j}$$

В итоге получаем разностную схему внутри интервала:

$$\begin{aligned} & - \left[\bar{h}_j r_{i+\frac{1}{2}} k_1(r_{i+\frac{1}{2}}, z_j) \frac{v_{i+1,j} - v_{i,j}}{h_r} - \bar{h}_j r_{i-\frac{1}{2}} k_1(r_{i-\frac{1}{2}}, z_j) \frac{v_{i,j} - v_{i-1,j}}{h_r} \right. \\ & \left. + \bar{h}_i r_i k_2(r_i, z_{j+\frac{1}{2}}) \frac{v_{i,j+1} - v_{i,j}}{h_z} - \bar{h}_i r_i k_2(r_i, z_{j-\frac{1}{2}}) \frac{v_{i,j} - v_{i,j-1}}{h_z} \right] = \bar{h}_i \bar{h}_j r_i f_{i,j} \end{aligned}$$

Перейдём к частному:

$$\begin{aligned} & - \left[h_z r_{i+\frac{1}{2}} k_1(r_{i+\frac{1}{2}}, z_j) \frac{v_{i+1,j} - v_{i,j}}{h_r} - h_z r_{i-\frac{1}{2}} k_1(r_{i-\frac{1}{2}}, z_j) \frac{v_{i,j} - v_{i-1,j}}{h_r} \right. \\ & \left. + h_r r_i k_2(r_i, z_{j+\frac{1}{2}}) \frac{v_{i,j+1} - v_{i,j}}{h_z} - h_r r_i k_2(r_i, z_{j-\frac{1}{2}}) \frac{v_{i,j} - v_{i,j-1}}{h_z} \right] = h_r h_z r_i f_{i,j} \end{aligned}$$

Теперь найдем значение разностной схемы на углах и границах интервалов

2.2 На левой границе

Проинтегрируем наше уравнение в $i = 0$ и z внутри промежутка

$$- \int_{r_i}^{r_{i+\frac{1}{2}}} \int_{z_{j-\frac{1}{2}}}^{z_{j+\frac{1}{2}}} \left[\frac{\partial}{\partial r} \left(r k_1(r, z) \frac{\partial u}{\partial r} \right) + \frac{\partial}{\partial z} \left(r k_2(r, z) \frac{\partial u}{\partial z} \right) \right] dr dz = \int_{r_i}^{r_{i+\frac{1}{2}}} \int_{z_{j-\frac{1}{2}}}^{z_{j+\frac{1}{2}}} r f(r, z) dr dz$$

Получаем:

$$\begin{aligned}
& - \left[\int_{z_{j-\frac{1}{2}}}^{z_{j+\frac{1}{2}}} rk_1(r, z) \frac{\partial u}{\partial r} \Big|_{r=r_{i+\frac{1}{2}}} dz - \int_{z_{j-\frac{1}{2}}}^{z_{j+\frac{1}{2}}} rk_1(r, z) \frac{\partial u}{\partial r} \Big|_{r=r_i} dz \right. \\
& \left. + \int_{r_i}^{r_{i+\frac{1}{2}}} rk_2(r, z) \frac{\partial u}{\partial z} \Big|_{z=z_{j+\frac{1}{2}}} dr - \int_{r_i}^{r_{i+\frac{1}{2}}} rk_2(r, z) \frac{\partial u}{\partial z} \Big|_{z=z_{j-\frac{1}{2}}} dr \right] = \int_{r_i}^{r_{i+\frac{1}{2}}} \int_{z_{j-\frac{1}{2}}}^{z_{j+\frac{1}{2}}} rf(r, z) dr dz
\end{aligned}$$

Имеем граничное условие:

$$u|_{r=0} - \text{ограничено, т. е. } \frac{\partial u}{\partial r} \Big|_{r=0} = 0$$

$$\int_{r_i}^{r_{i+\frac{1}{2}}} rf dr \approx f_i \int_{r_i}^{r_{i+\frac{1}{2}}} r dr = f_i \frac{r_{i+\frac{1}{2}}^2}{2} = \frac{h_r}{2} f_i \frac{r_{i+\frac{1}{2}}}{2}, \quad i = 0, \quad r_i = 0, r_{i+\frac{1}{2}} = \frac{h_r}{2}$$

Получаем разностную схему:

$$\begin{aligned}
& - \left[h_j r_{i+\frac{1}{2}} k_1(r_{i+\frac{1}{2}}, z_j) \frac{v_{i+1,j} - v_{i,j}}{h_{i+1}} - 0 \right. \\
& \left. + h_i r_i k_2(r_i, z_{j+\frac{1}{2}}) \frac{v_{i,j+1} - v_{i,j}}{h_{j+1}} - h_i r_i k_2(r_i, z_{j-\frac{1}{2}}) \frac{v_{i,j} - v_{i,j-1}}{h_j} \right] = h_i h_j r_i f_{i,j}
\end{aligned}$$

Перейдём к частному:

$$\begin{aligned}
& - \left[h_z r_{\frac{1}{2}} k_1(r_{\frac{1}{2}}, z_j) \frac{v_{1,j} - v_{0,j}}{h_r} - 0 \right. \\
& \left. + \frac{h_r}{2} \frac{r_{\frac{1}{2}}}{2} k_2(r_0, z_{j+\frac{1}{2}}) \frac{v_{0,j+1} - v_{0,j}}{h_z} - \frac{h_r}{2} \frac{r_{\frac{1}{2}}}{2} k_2(r_0, z_{j-\frac{1}{2}}) \frac{v_{0,j} - v_{0,j-1}}{h_z} \right] = \frac{h_r}{2} \frac{r_{\frac{1}{2}}}{2} h_z f_{0,j}
\end{aligned}$$

2.3 На правой границе

Проинтегрируем наше уравнение в $i = N_x$ и z внутри промежутка

$$- \int_{r_{i-\frac{1}{2}}}^{r_i} \int_{z_{j-\frac{1}{2}}}^{z_{j+\frac{1}{2}}} \left[\frac{\partial}{\partial r} \left(rk_1(r, z) \frac{\partial u}{\partial r} \right) + \frac{\partial}{\partial z} \left(rk_2(r, z) \frac{\partial u}{\partial z} \right) \right] dr dz = \int_{r_{i-\frac{1}{2}}}^{r_i} \int_{z_{j-\frac{1}{2}}}^{z_{j+\frac{1}{2}}} rf(r, z) dr dz$$

Получаем:

$$\begin{aligned}
& - \left[\int_{z_{j-\frac{1}{2}}}^{z_{j+\frac{1}{2}}} rk_1(r, z) \frac{\partial u}{\partial r} \Big|_{r=r_i} dz - \int_{z_{j-\frac{1}{2}}}^{z_{j+\frac{1}{2}}} rk_1(r, z) \frac{\partial u}{\partial r} \Big|_{r=r_{i-\frac{1}{2}}} dz \right. \\
& \left. + \int_{r_{i-\frac{1}{2}}}^{r_i} rk_2(r, z) \frac{\partial u}{\partial z} \Big|_{z=z_{j+\frac{1}{2}}} dr - \int_{r_{i-\frac{1}{2}}}^{r_i} rk_2(r, z) \frac{\partial u}{\partial z} \Big|_{z=z_{j-\frac{1}{2}}} dr \right] = \int_{r_{i-\frac{1}{2}}}^{r_i} \int_{z_{j-\frac{1}{2}}}^{z_{j+\frac{1}{2}}} rf(r, z) dr dz
\end{aligned}$$

Имеем граничное условие:

$$-k_1 \frac{\partial u}{\partial r} \Big|_{r=R} = \chi_2 u|_{r=R} - \varphi_2(z)$$

Получаем разностную схему:

$$\begin{aligned}
& - \left[-\hbar_j (\chi_2 v_i - \varphi_2(z)) - \hbar_j r_{i-\frac{1}{2}} k_1(r_{i-\frac{1}{2}}, z_j) \frac{v_{i,j} - v_{i-1,j}}{h_r} \right. \\
& \left. + \hbar_i r_i k_2(r_i, z_{j+\frac{1}{2}}) \frac{v_{i,j+1} - v_{i,j}}{h_z} - \hbar_i r_i k_2(r_i, z_{j-\frac{1}{2}}) \frac{v_{i,j} - v_{i,j-1}}{h_z} \right] = \hbar_i \hbar_j r_i f_{i,j}
\end{aligned}$$

Перейдём к частному:

$$\begin{aligned}
& - \left[-h_z (\chi_2 v_{N_r} - \varphi_2(z)) - h_z r_{N_r-\frac{1}{2}} k_1(r_{N_r-\frac{1}{2}}, z_j) \frac{v_{N_r,j} - v_{N_r-1,j}}{h_r} \right. \\
& \left. + \frac{h_r}{2} r_{N_r} k_2(r_{N_r}, z_{j+\frac{1}{2}}) \frac{v_{N_r,j+1} - v_{N_r,j}}{h_z} - \frac{h_r}{2} r_{N_r} k_2(r_{N_r}, z_{j-\frac{1}{2}}) \frac{v_{N_r,j} - v_{N_r,j-1}}{h_z} \right] = \frac{h_r}{2} r_{N_r} h_z f_{N_r,j}
\end{aligned}$$

2.4 На нижней границе

Проинтегрируем наше уравнение $j = 0$ и i внутри промежутка

$$- \int_{r_{i-\frac{1}{2}}}^{r_{i+\frac{1}{2}}} \int_{z_j}^{z_{j+\frac{1}{2}}} \left[\frac{\partial}{\partial r} \left(rk_1(r, z) \frac{\partial u}{\partial r} \right) + \frac{\partial}{\partial z} \left(rk_2(r, z) \frac{\partial u}{\partial z} \right) \right] dr dz = \int_{r_{i-\frac{1}{2}}}^{r_{i+\frac{1}{2}}} \int_{z_j}^{z_{j+\frac{1}{2}}} rf(r, z) dr dz$$

Получаем:

$$\begin{aligned}
& - \left[\int_{z_j}^{z_{j+\frac{1}{2}}} rk_1(r, z) \frac{\partial u}{\partial r} \Big|_{r=r_{i+\frac{1}{2}}} dz - \int_{z_j}^{z_{j+\frac{1}{2}}} rk_1(r, z) \frac{\partial u}{\partial r} \Big|_{r=r_{i-\frac{1}{2}}} dz \right. \\
& \left. + \int_{r_{i-\frac{1}{2}}}^{r_{i+\frac{1}{2}}} rk_2(r, z) \frac{\partial u}{\partial z} \Big|_{z=z_{j+\frac{1}{2}}} dr - \int_{r_{i-\frac{1}{2}}}^{r_{i+\frac{1}{2}}} rk_2(r, z) \frac{\partial u}{\partial z} \Big|_{z=z_j} dr \right] = \int_{r_{i-\frac{1}{2}}}^{r_{i+\frac{1}{2}}} \int_{z_j}^{z_{j+\frac{1}{2}}} rf(r, z) dr dz
\end{aligned}$$

Имеем граничное условие:

$$k_2 \frac{\partial u}{\partial z} \Big|_{z=0} = \chi_3 u|_{z=0} - \varphi_3(r)$$

Получаем разностную схему:

$$\begin{aligned} & - \left[\hbar_j r_{i+\frac{1}{2}} k_1(r_{i+\frac{1}{2}}, z_j) \frac{v_{i+1,j} - v_{i,j}}{h_r} - \hbar_j r_{i-\frac{1}{2}} k_1(r_{i-\frac{1}{2}}, z_j) \frac{v_{i,j} - v_{i-1,j}}{h_r} \right. \\ & \quad \left. + \hbar_i r_i k_2(r_i, z_{j+\frac{1}{2}}) \frac{v_{i,j+1} - v_{i,j}}{h_z} - \hbar_i (\chi_3 v_i - \varphi_3(r)) \right] = \hbar_i \hbar_j r_i f_{i,j} \end{aligned}$$

Перейдём к частному:

$$\begin{aligned} & - \left[\frac{h_z}{2} r_{i+\frac{1}{2}} k_1(r_{i+\frac{1}{2}}, z_0) \frac{v_{i+1,0} - v_{i,0}}{h_r} - \frac{h_z}{2} r_{i-\frac{1}{2}} k_1(r_{i-\frac{1}{2}}, z_0) \frac{v_{i,0} - v_{i-1,0}}{h_r} \right. \\ & \quad \left. + h_r r_i k_2(r_i, z_{\frac{1}{2}}) \frac{v_{i,1} - v_{i,0}}{h_z} - h_r (\chi_3 v_{i,0} - \varphi_3(r)) \right] = h_r \frac{h_z}{2} r_i f_{i,0} \end{aligned}$$

2.5 На верхней границе

Имеем граничное условие:

$$u|_{z=L} = \varphi_r(r)$$

$$v_{i,N_z} = \varphi(r_i)$$

2.6 Левый-нижний угол

Проинтегрируем наше уравнение в $i = 0$ и $j = 0$ внутри промежутка

$$- \int_{r_i}^{r_{i+\frac{1}{2}}} \int_{z_j}^{z_{j+\frac{1}{2}}} \left[\frac{\partial}{\partial r} \left(r k_1(r, z) \frac{\partial u}{\partial r} \right) + \frac{\partial}{\partial z} \left(r k_2(r, z) \frac{\partial u}{\partial z} \right) \right] dr dz = \int_{r_i}^{r_{i+\frac{1}{2}}} \int_{z_j}^{z_{j+\frac{1}{2}}} r f(r, z) dr dz$$

Получаем:

$$\begin{aligned} & - \left[\int_{z_j}^{z_{j+\frac{1}{2}}} r k_1(r, z) \frac{\partial u}{\partial r} \Big|_{r=r_{i+\frac{1}{2}}} dz - \int_{z_j}^{z_{j+\frac{1}{2}}} r k_1(r, z) \frac{\partial u}{\partial r} \Big|_{r=r_i} dz \right. \\ & \quad \left. + \int_{r_i}^{r_{i+\frac{1}{2}}} r k_2(r, z) \frac{\partial u}{\partial z} \Big|_{z=z_{j+\frac{1}{2}}} dr - \int_{r_i}^{r_{i+\frac{1}{2}}} r k_2(r, z) \frac{\partial u}{\partial z} \Big|_{z=z_j} dr \right] = \int_{r_i}^{r_{i+\frac{1}{2}}} \int_{z_j}^{z_{j+\frac{1}{2}}} r f(r, z) dr dz \end{aligned}$$

Имеем граничное условие:

$$u|_{r=0} - \text{ограничено, т. е. } \frac{\partial u}{\partial r}\bigg|_{r=0} = 0$$

$$\int_{r_i}^{r_{i+\frac{1}{2}}} r f dr \approx f_i \int_{r_i}^{r_{i+\frac{1}{2}}} r dr = f_i \frac{r_{i+\frac{1}{2}}^2}{2} = h_r f_i \frac{r_{i+\frac{1}{2}}}{2}, \quad i = 0, \quad r_i = 0, r_{i+\frac{1}{2}} = \frac{h_r}{2}$$

Также:

$$k_2 \frac{\partial u}{\partial z}\bigg|_{z=0} = \chi_3 u|_{z=0} - \varphi_3(r)$$

Получаем:

$$\begin{aligned} & - \left[\hbar_j r_{i+\frac{1}{2}} k_1(r_{i+\frac{1}{2}}, z_j) \frac{v_{i+1,j} - v_{i,j}}{h_r} - 0 \right. \\ & \left. + \hbar_i \frac{r_{i+\frac{1}{2}}}{2} k_2(r_i, z_{j+\frac{1}{2}}) \frac{v_{i,j+1} - v_{i,j}}{h_z} - \hbar_i \frac{r_{i+\frac{1}{2}}}{2} (\chi_3 u|_{z=0} - \varphi_3(r)) \right] = \hbar_i \hbar_j \frac{r_{i+\frac{1}{2}}}{2} f_{i,j} \end{aligned}$$

Перейдём к частному:

$$\begin{aligned} & - \left[\frac{h_z}{2} r_{\frac{1}{2}} k_1(r_{\frac{1}{2}}, z_0) \frac{v_{1,0} - v_{0,0}}{h_r} - 0 \right. \\ & \left. + \frac{h_r}{2} \frac{r_{\frac{1}{2}}}{2} k_2(r_0, z_{\frac{1}{2}}) \frac{v_{0,1} - v_{0,0}}{h_z} - \frac{h_r}{2} \frac{r_{\frac{1}{2}}}{2} (\chi_3 v_{0,0} - \varphi_3(r)) \right] = \frac{h_r}{2} \frac{h_z}{2} \frac{r_{\frac{1}{2}}}{2} f_{0,0} \end{aligned}$$

2.7 Левый-верхний угол

При $i = 0$ и $z = N_z$ имеем граничное условие:

Имеем граничное условие:

$$u|_{z=L} = \varphi_r(r)$$

$$v_{0,N_z} = \varphi(r_0)$$

2.8 Правый-верхний угол

При $i = N_r$ и $z = N_z$ имеем граничное условие:

Имеем граничное условие:

$$u|_{z=L} = \varphi_r(r)$$

$$v_{N_r,N_z} = \varphi(r_{N_r})$$

2.9 Правый-нижний угол

Проинтегрируем наше уравнение в $i = N_r$ и $j = 0$:

$$- \int_{r_{i-\frac{1}{2}}}^{r_i} \int_{z_j}^{z_{j+\frac{1}{2}}} \left[\frac{\partial}{\partial r} \left(rk_1(r, z) \frac{\partial u}{\partial r} \right) + \frac{\partial}{\partial z} \left(rk_2(r, z) \frac{\partial u}{\partial z} \right) \right] = \int_{r_{i-\frac{1}{2}}}^{r_i} \int_{z_j}^{z_{j+\frac{1}{2}}} rf(r, z)$$

Получаем:

$$\begin{aligned} & - \left[\int_{z_j}^{z_{j+\frac{1}{2}}} rk_1(r, z) \frac{\partial u}{\partial r} \Big|_{r=r_i} dz - \int_{z_j}^{z_{j+\frac{1}{2}}} rk_1(r, z) \frac{\partial u}{\partial r} \Big|_{r=r_{i-\frac{1}{2}}} dz \right. \\ & \left. + \int_{r_{i-\frac{1}{2}}}^{r_i} rk_2(r, z) \frac{\partial u}{\partial z} \Big|_{z=z_{j+\frac{1}{2}}} dr - \int_{r_{i-\frac{1}{2}}}^{r_i} rk_2(r, z) \frac{\partial u}{\partial z} \Big|_{z=z_j} dr \right] = \int_{r_{i-\frac{1}{2}}}^{r_i} \int_{z_j}^{z_{j+\frac{1}{2}}} rf(r, z) dr dz \end{aligned}$$

Имеем граничные условия:

$$\begin{aligned} -k_1 \frac{\partial u}{\partial r} \Big|_{r=R} &= \chi_2 u|_{r=R} - \varphi_2(z) \\ k_2 \frac{\partial u}{\partial z} \Big|_{z=0} &= \chi_3 u|_{z=0} - \varphi_3(r) \end{aligned}$$

Получаем разностную схему:

$$\begin{aligned} & - \left[-\bar{h}_j (\chi_2 u|_{r=R} - \varphi_2(z)) - \bar{h}_j r_{i-\frac{1}{2}} k_1(r_{i-\frac{1}{2}}, z_j) \frac{v_{i,j} - v_{i-1,j}}{h_r} \right. \\ & \left. + \bar{h}_i r_i k_2(r_i, z_{j+\frac{1}{2}}) \frac{v_{i,j+1} - v_{i,j}}{h_z} - \bar{h}_i (\chi_3 v_i - \varphi_3(r)) \right] = \bar{h}_i \bar{h}_j r_i f_{i,j} \end{aligned}$$

Перейдём к частному

$$\begin{aligned} & - \left[-\frac{h_z}{2} (\chi_2 v_{N_r,0} - \varphi_2(z)) - \frac{h_z}{2} r_{N_r-\frac{1}{2}} k_1(r_{N_r-\frac{1}{2}}, z_0) \frac{v_{N_r,0} - v_{N_r-1,0}}{h_r} \right. \\ & \left. + \frac{h_r}{2} r_{N_r} k_2(r_{N_r}, z_{\frac{1}{2}}) \frac{v_{N_r,1} - v_{N_r,0}}{h_z} - \frac{h_r}{2} (\chi_3 v_{N_r,0} - \varphi_3(r)) \right] = \frac{h_r}{2} \frac{h_z}{2} r_{N_r} f_{N_r,0} \end{aligned}$$

3 Невязка разностной схемы

3.1 Невязка во внутренних точках

Запишем для уравнения разностной сетки во внутренних точках невязку:

$$\begin{aligned}
 & - \left[h_z r_{i+\frac{1}{2}} k_1(r_{i+\frac{1}{2}}, z_j) \frac{v_{i+1,j} - v_{i,j}}{h_r} - h_z r_{i-\frac{1}{2}} k_1(r_{i-\frac{1}{2}}, z_j) \frac{v_{i,j} - v_{i-1,j}}{h_r} \right. \\
 & \left. + h_r r_i k_2(r_i, z_{j+\frac{1}{2}}) \frac{v_{i,j+1} - v_{i,j}}{h_z} - h_r r_i k_2(r_i, z_{j-\frac{1}{2}}) \frac{v_{i,j} - v_{i,j-1}}{h_z} \right] = h_r h_z r_i f_{i,j} \\
 \xi_{i,j} &= h_r h_z r_i f_{i,j} + \left[h_z r_{i+\frac{1}{2}} k_1(r_{i+\frac{1}{2}}, z_j) \frac{u_{i+1,j} - u_{i,j}}{h_r} - h_z r_{i-\frac{1}{2}} k_1(r_{i-\frac{1}{2}}, z_j) \frac{u_{i,j} - u_{i-1,j}}{h_r} \right. \\
 & \left. + h_r r_i k_2(r_i, z_{j+\frac{1}{2}}) \frac{u_{i,j+1} - u_{i,j}}{h_z} - h_r r_i k_2(r_i, z_{j-\frac{1}{2}}) \frac{u_{i,j} - u_{i,j-1}}{h_z} \right]
 \end{aligned}$$

Напишем разложение Тейлора для невязки:

$$\begin{aligned}
 u_{i,j-1} &= u(r_i, z_j - h_z) = \left[u - h_z \frac{\partial u}{\partial z} + \frac{h_z^2}{2} \frac{\partial^2 u}{\partial z^2} - \frac{h_z^3}{6} \frac{\partial^3 u}{\partial z^3} + \frac{h_z^4}{24} \frac{\partial^4 u}{\partial z^4} \right]_{i,j} + \mathcal{O}(h_y^5) \\
 \frac{u_{i,j} - u_{i,j-1}}{h_z} &= \left[\frac{\partial u}{\partial z} - \frac{h_z}{2} \frac{\partial^2 u}{\partial z^2} + \frac{h_z^2}{6} \frac{\partial^3 u}{\partial z^3} - \frac{h_z^3}{24} \frac{\partial^4 u}{\partial z^4} \right]_{i,j} + \mathcal{O}(h_y^4) \\
 r_i k_2(r_i, z_{j-\frac{1}{2}}) &= r_i k_2(r_i, z_j - \frac{h_z}{2}) = \left[r k_2 - \frac{h_z}{2} \frac{\partial r k_2}{\partial z} + \frac{h_z^2}{8} \frac{\partial^2 r k_2}{\partial z^2} - \frac{h_z^3}{48} \frac{\partial^3 r k_2}{\partial z^3} \right]_{i,j} + \mathcal{O}(h^4) \\
 r_i k_2(r_i, z_{j-\frac{1}{2}}) \frac{u_{i,j} - u_{i,j-1}}{h_z} &= \left[r k_2 \frac{\partial u}{\partial z} \right]_{i,j} - h_z \left[\frac{1}{2} r k_2 \frac{\partial^2 u}{\partial z^2} + \frac{1}{2} \frac{\partial r k_2}{\partial z} \frac{\partial u}{\partial z} \right]_{i,j} + \\
 &+ h_z^2 \left[\frac{1}{6} r k_2 \frac{\partial^3 u}{\partial z^3} + \frac{1}{4} \frac{\partial r k_2}{\partial z} \frac{\partial^2 u}{\partial z^2} + \frac{1}{8} \frac{\partial^2 r k_2}{\partial z^2} \frac{\partial u}{\partial z} \right]_{i,j} - \\
 &- h_z^3 \left[\frac{1}{24} r k_2 \frac{\partial^4 u}{\partial z^4} - \frac{1}{12} \frac{\partial r k_2}{\partial z} \frac{\partial^3 u}{\partial z^3} + \frac{1}{16} \frac{\partial^2 r k_2}{\partial z^2} \frac{\partial^2 u}{\partial z^2} + \frac{1}{48} \frac{\partial^3 r k_2}{\partial z^3} \frac{\partial u}{\partial z} \right]_{i,j} + \\
 &+ \mathcal{O}(h^4)
 \end{aligned}$$

Для $r_i k_2(r_i, z_{j+\frac{1}{2}})$ можно получить невязку аналогичным способом:

$$\begin{aligned}
 r_i k_2(r_i, z_{j+\frac{1}{2}}) \frac{u_{i,j+1} - u_{i,j}}{h_z} &= \left[r k_2 \frac{\partial u}{\partial z} \right]_{i,j} + h_z \left[\frac{1}{2} r k_2 \frac{\partial^2 u}{\partial z^2} + \frac{1}{2} \frac{\partial r k_2}{\partial z} \frac{\partial u}{\partial z} \right]_{i,j} + \\
 &+ h_z^2 \left[\frac{1}{6} r k_2 \frac{\partial^3 u}{\partial z^3} + \frac{1}{4} \frac{\partial r k_2}{\partial z} \frac{\partial^2 u}{\partial z^2} + \frac{1}{8} \frac{\partial^2 r k_2}{\partial z^2} \frac{\partial u}{\partial z} \right]_{i,j} + \\
 &+ h_z^3 \left[\frac{1}{24} r k_2 \frac{\partial^4 u}{\partial z^4} - \frac{1}{12} \frac{\partial r k_2}{\partial z} \frac{\partial^3 u}{\partial z^3} + \frac{1}{16} \frac{\partial^2 r k_2}{\partial z^2} \frac{\partial^2 u}{\partial z^2} + \frac{1}{48} \frac{\partial^3 r k_2}{\partial z^3} \frac{\partial u}{\partial z} \right]_{i,j} + \\
 &+ \mathcal{O}(h^4)
 \end{aligned}$$

$$u_{i-1,j} = u(r_i - h_r, j) = \left[u - h_r \frac{\partial u}{\partial r} + \frac{h_r^2}{2} \frac{\partial^2 u}{\partial r^2} - \frac{h_r^3}{6} \frac{\partial^3 u}{\partial r^3} + \frac{h_r^4}{24} \frac{\partial^4 u}{\partial r^4} \right]_{i,j} + \mathcal{O}(h^5)$$

$$\frac{u_{i,j} - u_{i-1,j}}{h_r} = \left[\frac{\partial u}{\partial r} - \frac{h_r}{2} \frac{\partial^2 u}{\partial r^2} + \frac{h_r^2}{6} \frac{\partial^3 u}{\partial r^3} - \frac{h_r^3}{24} \frac{\partial^4 u}{\partial r^4} \right]_{i,j} + \mathcal{O}(h^4)$$

$$r_i k_1(r_{i-\frac{1}{2}}, z_j) = r_i k_1(x_i - \frac{h_r}{2}, z_j) = \left[r k_1 - \frac{h_r}{2} \frac{\partial r k_1}{\partial r} + \frac{h_r^2}{8} \frac{\partial^2 r k_1}{\partial r^2} - \frac{h_r^3}{48} \frac{\partial^3 r k_1}{\partial r^3} \right]_{i,j} + \mathcal{O}(h^4)$$

$$\begin{aligned} r_i k_1(r_{i-\frac{1}{2}}, z_j) \frac{u_{i,j} - u_{i-1,j}}{h_r} &= \left[r k_1 \frac{\partial u}{\partial r} \right]_{i,j} - h_r \left[\frac{1}{2} r k_1 \frac{\partial^2 u}{\partial r^2} + \frac{1}{2} \frac{\partial r k_1}{\partial r} \frac{\partial u}{\partial r} \right]_{i,j} + \\ &+ h_r^2 \left[\frac{1}{6} r k_1 \frac{\partial^3 u}{\partial r^3} + \frac{1}{4} \frac{\partial r k_1}{\partial r} \frac{\partial^2 u}{\partial r^2} + \frac{1}{8} \frac{\partial^2 r k_1}{\partial r^2} \frac{\partial u}{\partial r} \right]_{i,j} - \\ &- h_r^3 \left[\frac{1}{24} r k_1 \frac{\partial^4 u}{\partial r^4} - \frac{1}{12} \frac{\partial r k_1}{\partial r} \frac{\partial^3 u}{\partial r^3} + \frac{1}{16} \frac{\partial^2 r k_1}{\partial r^2} \frac{\partial^2 u}{\partial r^2} + \frac{1}{48} \frac{\partial^3 r k_1}{\partial r^3} \frac{\partial u}{\partial r} \right]_{i,j} + \\ &+ \mathcal{O}(h^4) \end{aligned}$$

Для $r_i k_1(x_{i=\frac{1}{2}}, z_j) \frac{u_{i+1,j} - u_{i,j}}{h_r}$ можно получить невязку аналогично:

$$\begin{aligned} r_i k_1(x_{i+\frac{1}{2}}, z_j) \frac{u_{i+1,j} - u_{i,j}}{h_r} &= \left[r k_1 \frac{\partial u}{\partial r} \right]_{i,j} + h_r \left[\frac{1}{2} r k_1 \frac{\partial^2 u}{\partial r^2} + \frac{1}{2} \frac{\partial r k_1}{\partial r} \frac{\partial u}{\partial r} \right]_{i,j} + \\ &+ h_r^2 \left[\frac{1}{6} r k_1 \frac{\partial^3 u}{\partial r^3} + \frac{1}{4} \frac{\partial r k_1}{\partial r} \frac{\partial^2 u}{\partial r^2} + \frac{1}{8} \frac{\partial^2 r k_1}{\partial r^2} \frac{\partial u}{\partial r} \right]_{i,j} + \\ &+ h_r^3 \left[\frac{1}{24} r k_1 \frac{\partial^4 u}{\partial r^4} - \frac{1}{12} \frac{\partial r k_1}{\partial r} \frac{\partial^3 u}{\partial r^3} + \frac{1}{16} \frac{\partial^2 r k_1}{\partial r^2} \frac{\partial^2 u}{\partial r^2} + \frac{1}{48} \frac{\partial^3 r k_1}{\partial r^3} \frac{\partial u}{\partial r} \right]_{i,j} + \\ &+ \mathcal{O}(h^4) \end{aligned}$$

Тогда:

$$\begin{aligned} r_i k_1(x_{i+\frac{1}{2}}, z_j) \frac{u_{i+1,j} - u_{i,j}}{h_r} - r_i k_1(r_{i-\frac{1}{2}}, z_j) \frac{u_{i,j} - u_{i-1,j}}{h_r} &= h_r h_z \left(\left[r k_1 \frac{\partial^2 u}{\partial r^2} + \frac{\partial r k_1}{\partial r} \frac{\partial u}{\partial r} \right]_{i,j} + \right. \\ &+ h_r^2 \left[\frac{1}{12} r k_1 \frac{\partial^4 u}{\partial r^4} - \frac{1}{6} \frac{\partial r k_1}{\partial r} \frac{\partial^3 u}{\partial r^3} + \frac{1}{8} \frac{\partial^2 r k_1}{\partial r^2} \frac{\partial^2 u}{\partial r^2} + \frac{1}{24} \frac{\partial^3 r k_1}{\partial r^3} \frac{\partial u}{\partial r} \right] + \mathcal{O}(h_r^4) \Big) \end{aligned}$$

$$\begin{aligned} r_i k_2(r_i, z_{j+\frac{1}{2}}) \frac{u_{i,j+1} - u_{i,j}}{h_z} - r_i k_2(r_i, z_{j-\frac{1}{2}}) \frac{u_{i,j} - u_{i,j-1}}{h_z} &= h_r h_z \left(\left[r k_2 \frac{\partial^2 u}{\partial z^2} + \frac{\partial r k_2}{\partial z} \frac{\partial u}{\partial z} \right]_{i,j} + \right. \\ &+ h_z^2 \left[\frac{1}{12} r k_2 \frac{\partial^4 u}{\partial z^4} - \frac{1}{6} \frac{\partial r k_2}{\partial z} \frac{\partial^3 u}{\partial z^3} + \frac{1}{8} \frac{\partial^2 r k_2}{\partial z^2} \frac{\partial^2 u}{\partial z^2} + \frac{1}{24} \frac{\partial^3 r k_2}{\partial z^3} \frac{\partial u}{\partial z} \right]_{i,j} + \mathcal{O}(h_z^4) \Big) \end{aligned}$$

Будем искать невязку в следующем виде:

$$\tilde{\xi}_{i,j} = \frac{\xi_{i,j}}{h_r h_z}$$

Тогда:

$$\begin{aligned} \tilde{\xi}_{i,j} = & r_i f_{i,j} + \left[rk_1 \frac{\partial^2 u}{\partial r^2} + \frac{\partial rk_1}{\partial r} \frac{\partial u}{\partial r} \right]_{i,j} \left[rk_2 \frac{\partial^2 u}{\partial z^2} + \frac{\partial rk_2}{\partial z} \frac{\partial u}{\partial z} \right]_{i,j} + \\ & + h_r^2 \left[\frac{1}{12} rk_1 \frac{\partial^4 u}{\partial r^4} - \frac{1}{6} \frac{\partial rk_1}{\partial r} \frac{\partial^3 u}{\partial r^3} + \frac{1}{8} \frac{\partial^2 rk_1}{\partial r^2} \frac{\partial^2 u}{\partial r^2} + \frac{1}{24} \frac{\partial^3 rk_1}{\partial r^3} \frac{\partial u}{\partial r} \right] + \mathcal{O}(h_r^3) + \\ & + h_z^2 \left[\frac{1}{12} rk_2 \frac{\partial^4 u}{\partial z^4} - \frac{1}{6} \frac{\partial rk_2}{\partial z} \frac{\partial^3 u}{\partial z^3} + \frac{1}{8} \frac{\partial^2 rk_2}{\partial z^2} \frac{\partial^2 u}{\partial z^2} + \frac{1}{24} \frac{\partial^3 rk_2}{\partial z^3} \frac{\partial u}{\partial z} \right]_{i,j} + \mathcal{O}(h_z^3) \end{aligned}$$

Можно заметить, что:

$$\begin{aligned} \left[rk_1 \frac{\partial^2 u}{\partial r^2} + \frac{\partial rk_1}{\partial r} \frac{\partial u}{\partial r} \right]_{i,j} &= \frac{\partial}{\partial r} \left(k_1 \frac{\partial u}{\partial r} \right) \\ \left[rk_2 \frac{\partial^2 u}{\partial z^2} + \frac{\partial rk_2}{\partial z} \frac{\partial u}{\partial z} \right]_{i,j} &= \frac{\partial}{\partial z} \left(k_2 \frac{\partial u}{\partial z} \right) \end{aligned}$$

Тем самым:

$$r_i f_{i,j} + \frac{\partial}{\partial r} \left(k_1 \frac{\partial u}{\partial r} \right) + \frac{\partial}{\partial z} \left(k_2 \frac{\partial u}{\partial z} \right) = 0$$

Получаем:

$$\begin{aligned} \tilde{\xi}_{i,j} = & h_r^2 \left[\frac{1}{12} rk_1 \frac{\partial^4 u}{\partial r^4} - \frac{1}{6} \frac{\partial rk_1}{\partial r} \frac{\partial^3 u}{\partial r^3} + \frac{1}{8} \frac{\partial^2 rk_1}{\partial r^2} \frac{\partial^2 u}{\partial r^2} + \frac{1}{24} \frac{\partial^3 rk_1}{\partial r^3} \frac{\partial u}{\partial r} \right] + \mathcal{O}(h_r^3) + \\ & + h_z^2 \left[\frac{1}{12} rk_2 \frac{\partial^4 u}{\partial z^4} - \frac{1}{6} \frac{\partial rk_2}{\partial z} \frac{\partial^3 u}{\partial z^3} + \frac{1}{8} \frac{\partial^2 rk_2}{\partial z^2} \frac{\partial^2 u}{\partial z^2} + \frac{1}{24} \frac{\partial^3 rk_2}{\partial z^3} \frac{\partial u}{\partial z} \right]_{i,j} + \mathcal{O}(h_z^3) \end{aligned}$$

Порядок аппроксимации $p_r = 2 - 0$, $p_z = 2 - 0$.

3.2 Невязка на левой границе

4 Запись СЛАУ

Перейдём к одноиндексной записи

$$m = j(N_r + 1) + i$$

Индексы изменяются в следующих границах:

$$0 \leq i \leq N_r$$

$$0 \leq j \leq N_z$$

Тогда имеем:

$$0 \leq m < (N_r + 1)(N_z + 1)$$

4.1 Запись для внутренних точек

Перепишем наше уравнение с использованием нового индеса для $i \in (0, N_r)$ и $j \in (0, N_z)$:

$$\begin{aligned} & - \left[h_z r_{i+\frac{1}{2}} k_1(r_{i+\frac{1}{2}}, z_j) \frac{v_{i+1,j} - v_{i,j}}{h_r} - h_z r_{i-\frac{1}{2}} k_1(r_{i-\frac{1}{2}}, z_j) \frac{v_{i,j} - v_{i-1,j}}{h_r} \right. \\ & \left. + h_r r_i k_2(r_i, z_{j+\frac{1}{2}}) \frac{v_{i,j+1} - v_{i,j}}{h_z} - h_r r_i k_2(r_i, z_{j-\frac{1}{2}}) \frac{v_{i,j} - v_{i,j-1}}{h_z} \right] = h_r h_z r_i f_{i,j} \\ & - \frac{h_r}{h_z} r_i k_2(r_i, z_{j-\frac{1}{2}}) v_{i,j-1} - \frac{h_z}{h_r} r_{i-\frac{1}{2}} k_1(r_{i-\frac{1}{2}}, z_j) v_{i-1,j} + \\ & + \left[\frac{h_z}{h_r} r_{i+\frac{1}{2}} k_1(r_{i+\frac{1}{2}}, z_j) + \frac{h_z}{h_r} r_{i-\frac{1}{2}} k_1(r_{i-\frac{1}{2}}, z_j) + \frac{h_r}{h_z} r_i k_2(r_i, z_{j+\frac{1}{2}}) + \frac{h_r}{h_z} r_i k_2(r_i, z_{j-\frac{1}{2}}) \right] v_{i,j} \\ & - \frac{h_z}{h_r} r_{i+\frac{1}{2}} k_1(r_{i+\frac{1}{2}}, z_j) v_{i+1,j} - \frac{h_r}{h_z} r_i k_2(r_i, z_{j+1}) v_{i,j+\frac{1}{2}} = h_r h_z r_i f_{i,j} \end{aligned}$$

Введём новые обозначения:

$$a_m w_{m-L} + b_m w_{m-1} + c_m w_m + d_m w_{m+1} + e_m w_{m+L} = g_m$$

где коэффициенты равны:

$$a_m = -\frac{h_r}{h_z} r_i k_2(r_i, z_{j-\frac{1}{2}})$$

$$b_m = -\frac{h_z}{h_r} r_{i-\frac{1}{2}} k_1(r_{i-\frac{1}{2}}, z_j)$$

$$c_m = \frac{h_z}{h_r} r_{i+\frac{1}{2}} k_1(r_{i+\frac{1}{2}}, z_j) + \frac{h_z}{h_r} r_{i-\frac{1}{2}} k_1(r_{i-\frac{1}{2}}, z_j) + \frac{h_r}{h_z} r_i k_2(r_i, z_{j+\frac{1}{2}}) + \frac{h_r}{h_z} r_i k_2(r_i, z_{j-\frac{1}{2}})$$

$$d_m = -\frac{h_z}{h_r} r_{i+\frac{1}{2}} k_1(r_{i+\frac{1}{2}}, z_j)$$

$$e_m = -\frac{h_r}{h_z} r_i k_2(r_i, z_{j+1})$$

$$g_m = h_r h_z r_i f_{i,j}$$

4.2 Запись для левой границы

Перепишем наше уравнение с использованием нового индекса для $i = 0$ и $j \in (0, N_z)$:

$$\begin{aligned} & - \left[h_z r_{\frac{1}{2}} k_1(r_{\frac{1}{2}}, z_j) \frac{v_{1,j} - v_{0,j}}{h_r} - 0 \right. \\ & \left. + \frac{h_r}{2} \frac{r_{\frac{1}{2}}}{2} k_2(r_0, z_{j+\frac{1}{2}}) \frac{v_{0,j+1} - v_{0,j}}{h_z} - \frac{h_r}{2} \frac{r_{\frac{1}{2}}}{2} k_2(r_0, z_{j-\frac{1}{2}}) \frac{v_{0,j} - v_{0,j-1}}{h_z} \right] = \frac{h_r}{2} \frac{r_{\frac{1}{2}}}{2} h_z f_{0,j} \\ & - \frac{h_r}{2h_z} \frac{r_{\frac{1}{2}}}{2} k_2(r_0, z_{j-\frac{1}{2}}) v_{0,j-1} + \\ & + \left[\frac{h_z}{h_r} r_{\frac{1}{2}} k_1(r_{\frac{1}{2}}, z_j) + \frac{h_r}{2h_z} \frac{r_{\frac{1}{2}}}{2} k_2(r_0, z_{j+\frac{1}{2}}) + \frac{h_r}{2h_z} \frac{r_{\frac{1}{2}}}{2} k_2(r_0, z_{j-\frac{1}{2}}) \right] v_{0,j} \\ & - \frac{h_z}{h_r} r_{\frac{1}{2}} k_1(r_{\frac{1}{2}}, z_j) v_{1,j} - \frac{h_r}{2h_z} \frac{r_{\frac{1}{2}}}{2} k_2(r_0, z_{j+\frac{1}{2}}) v_{0,j+1} = \frac{h_r}{2} \frac{r_{\frac{1}{2}}}{2} h_z f_{0,j} \end{aligned}$$

Введём новые обозначения:

$$a_m w_{m-L} + c_m w_m + d_m w_{m+1} + e_m w_{m+L} = g_m$$

где коэффициенты равны:

$$a_m = -\frac{h_r}{2h_z} \frac{r_{\frac{1}{2}}}{2} k_2(r_0, z_{j-\frac{1}{2}})$$

$$c_m = \frac{h_z}{h_r} r_{\frac{1}{2}} k_1(r_{\frac{1}{2}}, z_j) + \frac{h_r}{2h_z} \frac{r_{\frac{1}{2}}}{2} k_2(r_0, z_{j+\frac{1}{2}}) + \frac{h_r}{2h_z} \frac{r_{\frac{1}{2}}}{2} k_2(r_0, z_{j-\frac{1}{2}})$$

$$d_m = -\frac{h_z}{h_r} r_{\frac{1}{2}} k_1(r_{\frac{1}{2}}, z_j)$$

$$e_m = -\frac{h_r}{2h_z} \frac{r_{\frac{1}{2}}}{2} k_2(r_0, z_{j+\frac{1}{2}})$$

$$g_m = \frac{h_r}{2} \frac{r_{\frac{1}{2}}}{2} h_z f_{0,j}$$

4.3 Запись для правой границы

Перепишем наше уравнение с использованием нового индекса для $i = N_r$ и $j \in (0, N_z)$:

$$\begin{aligned}
& - \left[-h_z(\chi_2 v_{N_r} - \varphi_2(z)) - h_z r_{N_r-\frac{1}{2}} k_1(r_{N_r-\frac{1}{2}}, z_j) \frac{v_{N_r,j} - v_{N_r-1,j}}{h_r} \right. \\
& + \frac{h_r}{2} r_{N_r} k_2(r_{N_r}, z_{j+\frac{1}{2}}) \frac{v_{N_r,j+1} - v_{N_r,j}}{h_z} - \frac{h_r}{2} r_{N_r} k_2(r_{N_r}, z_{j-\frac{1}{2}}) \frac{v_{N_r,j} - v_{N_r,j-1}}{h_z} \left. \right] = \frac{h_r}{2} r_{N_r} h_z f_{N_r,j} \\
& - \frac{h_r}{2h_z} r_{N_r} k_2(r_{N_r}, z_{j-\frac{1}{2}}) v_{N_r,j-1} - \frac{h_z}{h_r} r_{N_r-\frac{1}{2}} k_1(r_{N_r-\frac{1}{2}}, z_j) v_{N_r-1,j} \\
& + \left[h_z \chi_2 + \frac{h_z}{h_r} r_{N_r-\frac{1}{2}} k_1(r_{N_r-\frac{1}{2}}, z_j) + \frac{h_r}{2h_z} r_{N_r} k_2(r_{N_r}, z_{j+\frac{1}{2}}) + \frac{h_r}{2h_z} r_{N_r} k_2(r_{N_r}, z_{j-\frac{1}{2}}) \right] v_{N_r,j} \\
& - \frac{h_r}{2h_z} r_{N_r} k_2(r_{N_r}, z_{j+\frac{1}{2}}) v_{N_r,j+1} = \frac{h_r}{2} r_{N_r} h_z f_{N_r,j} + h_z \phi_2(z)
\end{aligned}$$

Введём новые обозначения:

$$a_m w_{m-L} + b_m w_{m-1} + c_m w_m + e_m w_{m+L} = g_m$$

где коэффициенты:

$$a_m = -\frac{h_r}{2h_z} r_{N_r} k_2(r_{N_r}, z_{j-\frac{1}{2}})$$

$$b_m = -\frac{h_z}{h_r} r_{N_r-\frac{1}{2}} k_1(r_{N_r-\frac{1}{2}}, z_j)$$

$$c_m = h_z \chi_2 + \frac{h_z}{h_r} r_{N_r-\frac{1}{2}} k_1(r_{N_r-\frac{1}{2}}, z_j) + \frac{h_r}{2h_z} r_{N_r} k_2(r_{N_r}, z_{j+\frac{1}{2}}) + \frac{h_r}{2h_z} r_{N_r} k_2(r_{N_r}, z_{j-\frac{1}{2}})$$

$$e_m = -\frac{h_r}{2h_z} r_{N_r} k_2(r_{N_r}, z_{j+\frac{1}{2}})$$

$$g_m = \frac{h_r}{2} r_{N_r} h_z f_{N_r,j} + h_z \varphi_2(z)$$

4.4 Запись для нижней границы

Перепишем наше уравнение с использованием нового индекса для $i \in (0, N_r)$ и $j = 0$:

$$\begin{aligned}
& - \left[\frac{h_z}{2} r_{i+\frac{1}{2}} k_1(r_{i+\frac{1}{2}}, z_0) \frac{v_{i+1,0} - v_{i,0}}{h_r} - \frac{h_z}{2} r_{i-\frac{1}{2}} k_1(r_{i-\frac{1}{2}}, z_0) \frac{v_{i,0} - v_{i-1,0}}{h_r} \right. \\
& \left. + h_r r_i k_2(r_i, z_{\frac{1}{2}}) \frac{v_{i,1} - v_{i,0}}{h_z} - h_r (\chi_3 v_{i,0} - \varphi_3(r)) \right] = h_r \frac{h_z}{2} r_i f_{i,0}
\end{aligned}$$

$$\begin{aligned}
& -\frac{h_z}{2h_r}v_{i-\frac{1}{2}}k_1(r_{i-\frac{1}{2}}, z_0)v_{i-1,0} \\
& + \left[h_r\chi_3 + \frac{h_z}{2h_r}r_{i+\frac{1}{2}}k_1(r_{i+\frac{1}{2}}, z_0) + \frac{h_z}{2h_r}r_{i-\frac{1}{2}}k_1(r_{i-\frac{1}{2}}, z_0) + \frac{h_r}{h_z}r_ik_2(r_i, z_{\frac{1}{2}}) \right] v_{i,0} \\
& - \frac{h_z}{2h_r}r_{i+\frac{1}{2}}k_1(r_{i+\frac{1}{2}}, z_0)v_{i+1,0} - \frac{h_r}{h_z}r_ik_2(r_i, z_{\frac{1}{2}})v_{i,1} = h_r\frac{h_z}{2}r_if_{i,0} + h_r\varphi(r)
\end{aligned}$$

Введём новые обозначения:

$$b_m w_{m-1} + c_m w_m + d_m w_{m+1} + e_m w_{m+L} = g_m$$

где коэффициенты равны:

$$b_m = -\frac{h_z}{2h_r}r_{i-\frac{1}{2}}k_1(r_{i-\frac{1}{2}}, z_0)$$

$$c_m = h_r\chi_3 + \frac{h_z}{2h_r}r_{i+\frac{1}{2}}k_1(r_{i+\frac{1}{2}}, z_0) + \frac{h_z}{2h_r}r_{i-\frac{1}{2}}k_1(r_{i-\frac{1}{2}}, z_0) + \frac{h_r}{h_z}r_ik_2(r_i, z_{\frac{1}{2}})$$

$$d_m = -\frac{h_z}{2h_r}r_{i+\frac{1}{2}}k_1(r_{i+\frac{1}{2}}, z_0)$$

$$e_m = -\frac{h_r}{h_z}r_ik_2(r_i, z_{\frac{1}{2}})$$

$$g_m = h_r\frac{h_z}{2}r_if_{i,0} + h_r\varphi_3(r)$$

4.5 Запись для верхней границы

Перепишем наше уравнение с использованием нового индекса для $i \in [0, N_r]$ и $j = N_z$:

$$v_{i,N_z} = \varphi(r_i)$$

Перейдём к новым обозначениям:

$$c_m w_m = \varphi_m$$

где:

$$c_m = 1, \quad \varphi_m = \varphi_2(r_i)$$

4.6 Запись для левой нижней граничной точки

Перепишем наше уравнение с использованием нового индекса для $i = 0$ и $j = 0$:

$$\begin{aligned}
& - \left[\frac{h_z}{2} r_{\frac{1}{2}} k_1(r_{\frac{1}{2}}, z_0) \frac{v_{1,0} - v_{0,0}}{h_r} - 0 \right. \\
& \left. + \frac{h_r}{2} \frac{r_{\frac{1}{2}}}{2} k_2(r_0, z_{\frac{1}{2}}) \frac{v_{0,1} - v_{0,0}}{h_z} - \frac{h_r}{2} \frac{r_{\frac{1}{2}}}{2} (\chi_3 v_{0,0} - \varphi_3(r)) \right] = \frac{h_r}{2} \frac{h_z}{2} \frac{r_{\frac{1}{2}}}{2} f_{0,0} \\
& \left[\frac{h_z}{2h_r} r_{\frac{1}{2}} k_1(r_{\frac{1}{2}}, z_0) + \frac{h_r}{2h_z} \frac{r_{\frac{1}{2}}}{2} k_2(r_0, z_{\frac{1}{2}}) + \frac{h_r}{2} \chi_3 \right] v_{0,0} - \\
& - \frac{h_z}{2h_r} r_{\frac{1}{2}} k_1(r_{\frac{1}{2}}, z_0) v_{1,0} - \frac{h_r}{2h_z} \frac{r_{\frac{1}{2}}}{2} k_2(r_0, z_{\frac{1}{2}}) v_{0,1} = \frac{h_r}{2} \frac{h_z}{2} \frac{r_{\frac{1}{2}}}{2} f_{0,0} + \frac{h_r}{2} \frac{r_{\frac{1}{2}}}{2} \varphi_3(r_0)
\end{aligned}$$

Введём новые обозначения:

$$c_m w_m + d_m w_{m+1} + e_m w_{m+L} = g_m$$

где коэффициенты ранвы:

$$\begin{aligned}
c_m &= \frac{h_z}{2h_r} r_{\frac{1}{2}} k_1(r_{\frac{1}{2}}, z_0) + \frac{h_r}{2h_z} \frac{r_{\frac{1}{2}}}{2} k_2(r_0, z_{\frac{1}{2}}) + \frac{h_r}{2} \chi_3 \\
d_m &= -\frac{h_z}{2h_r} r_{\frac{1}{2}} k_1(r_{\frac{1}{2}}, z_0) \\
e_m &= -\frac{h_r}{2h_z} \frac{r_{\frac{1}{2}}}{2} k_2(r_0, z_{\frac{1}{2}}) \\
g_m &= \frac{h_r}{2} \frac{h_z}{2} \frac{r_{\frac{1}{2}}}{2} f_{0,0} + \frac{h_r}{2} \frac{r_{\frac{1}{2}}}{2} \varphi_3(r_0)
\end{aligned}$$

4.7 Запись для правой нижней граничной точки

Перепишем наше уравнение с использованием нового индекса для $i = N_r$ и $j = 0$:

$$\begin{aligned}
& - \left[-\frac{h_z}{2} (\chi_2 v_{N_r,0} - \varphi_2(z)) - \frac{h_z}{2} r_{N_r - \frac{1}{2}} k_1(r_{N_r - \frac{1}{2}}, z_0) \frac{v_{N_r,0} - v_{N_r-1,0}}{h_r} \right. \\
& \left. + \frac{h_r}{2} r_{N_r} k_2(r_{N_r}, z_{\frac{1}{2}}) \frac{v_{N_r,1} - v_{N_r,0}}{h_z} - \frac{h_r}{2} (\chi_3 v_{N_r,0} - \varphi_3(r)) \right] = \frac{h_r}{2} \frac{h_z}{2} r_{N_r} f_{N_r,0}
\end{aligned}$$

$$\begin{aligned}
& -\frac{h_z}{2h_r}r_{N_r-\frac{1}{2}}k_1(r_{N_r-\frac{1}{2}},z_0)v_{N_r-1,0}+ \\
& +\left[\frac{h_z}{2}\chi_2+\frac{h_z}{2h_r}r_{N_r-\frac{1}{2}}k_1(r_{N_r-\frac{1}{2}},z_0)+\frac{h_r}{2h_z}r_{N_r}k_2(r_{N_r,z_{\frac{1}{2}}})+\frac{h_r}{2}\chi_3\right]v_{N_r,0}- \\
& -\frac{h_r}{2h_z}r_{N_r}k_2(r_{N_r},z_{\frac{1}{2}})v_{N_r,1}=\frac{h_r}{2}\frac{h_z}{2}r_{N_r}f_{N_r,0}+\frac{h_z}{2}\varphi_2(z)+\frac{h_r}{2}\varphi_3(r)
\end{aligned}$$

Введём новые обозначения:

$$b_m w_{m-1} + c_m w_m + e_m w_{m+L} = g_m$$

где коэффициенты равны:

$$b_m = -\frac{h_z}{2h_r}r_{N_r-\frac{1}{2}}k_1(r_{N_r-\frac{1}{2}},z_0)$$

$$c_m = \frac{h_z}{2}\chi_2 + \frac{h_z}{2h_r}r_{N_r-\frac{1}{2}}k_1(r_{N_r-\frac{1}{2}},z_0) + \frac{h_r}{2h_z}r_{N_r}k_2(r_{N_r,z_{\frac{1}{2}}}) + \frac{h_r}{2}\chi_3$$

$$e_m = -\frac{h_r}{2h_z}r_{N_r}k_2(r_{N_r},z_{\frac{1}{2}})$$

$$g_m = \frac{h_r}{2}\frac{h_z}{2}r_{N_r}f_{N_r,0} + \frac{h_z}{2}\varphi_2(z) + \frac{h_r}{2}\varphi_3(r)$$

5 Заключение