

eNotes: The Conic Section

CBCO

1 Introduction

A conic section is an intersection of a cone with a plane.

In Figure 1.1, the cones were plotted with two different methods.




Figure 1.1: (a) Cone = $f(x,y,z)$ (b) Cone = $f(\rho,\phi,z)$

Open Cone01.py. Study the codes and run the file.

The equation in cylindrical coordinate systems is expressed as follows.

$$(1.1) \quad \rho = \frac{b_{ase}(height - z)}{height}$$

Note that ϕ is absent. It means that for any value of phi eConeCyl equation holds.

The geometric relationship between Cartesian and cylindrical coordinate are expressed as follows.

$$(1.2) \quad x = \rho \cos(\phi)$$

$$(1.3) \quad y = \rho \sin(\phi)$$

$$(1.4) \quad z = z$$

The equation of cone can be found at <https://mathworld.wolfram.com/Cone.html>

$$(1.5) \quad \delta = 2 \operatorname{atan}\left(\frac{b_{ase}}{h_{eight}}\right)$$

$$(1.6) \quad z = u$$

$$(1.7) \quad x = \frac{b_{ase}(h_{eight} - u) \cos(\phi)}{h_{eight}}$$

$$(1.8) \quad y = \frac{b_{ase}(h_{eight} - u) \sin(\phi)}{h_{eight}}$$

The equations above is the conversion of cylindrical coordinate to cartesian systems.

$$(1.9) \quad x = \frac{b_{ase}(h_{eight} - z) \cos(\phi)}{h_{eight}}$$

$$(1.10) \quad y = \frac{b_{ase}(h_{eight} - z) \sin(\phi)}{h_{eight}}$$

$$(1.11) \quad x = \frac{b_{ase}(h_{eight} - z) \cos(\phi)}{h_{eight}} = \rho \cos(\phi)$$

$$(1.12) \quad y = \frac{b_{ase}(h_{eight} - z) \sin(\phi)}{h_{eight}} = \rho \sin(\phi)$$

The symbolic math equation of a cone in Cartesian system is expressed as follows.

$$(1.13) \quad x^2 + y^2 = \frac{b_{ase}^2 (h_{eigh}t - z)^2}{h_{eigh}t^2}$$

The base radius and height of the cone are given as follows.

$$(1.14) \quad b_{ase} = 2$$

$$(1.15) \quad h_{eigh}t = 8$$

The equation of the cone is substituted with numerical parameters given above as follows.

$$(1.16) \quad x^2 + y^2 = \frac{(z - 8)^2}{16}$$

2 Surface Plot Algorithms of a Cone

The mathematical expressions for surface plots in Cartesian coordinate system are expressed as follows.

$$(2.1) \quad z = f(x, y) \quad \text{where } z \text{ is dependent variable and } x \text{ and } y \text{ are independent variables}$$

$$(2.2) \quad x = f(z, y) \quad \text{where } x \text{ is dependent variable and } z \text{ and } y \text{ are independent variables}$$

$$(2.3) \quad y = f(x, z) \quad \text{where } y \text{ is dependent variable and } x \text{ and } z \text{ are independent variables}$$

The mathematical expression for surface plots in cylindrical coordinate system are expressed as follows.

$$(2.4) \quad \phi = f(\rho, z) \quad \text{where } \phi \text{ is the dependent variable and } \rho \text{ and } z \text{ are independent variables}$$

$$(2.5) \quad \rho = f(\phi, z) \quad \text{where } \rho \text{ is the dependent variable and } \phi \text{ and } z \text{ are independent variables}$$

$$(2.6) \quad z = f(\phi, \rho) \quad \text{where } z \text{ is the dependent variable and } \phi \text{ and } \rho \text{ are independent variables}$$

The surface plots of a cone in Figure 1 (a) were generated by cone01.py codes. The dependent variable is z and the independent variables are x and y . Thus $z = f(x,y)$. The algorithm of plot 1 is directly Cartesian meshgrid. Hence, $mx, my = \text{meshgrid}(x,y)$. The meshgrid of z is $mz=f(mx,my)$.

The algorithm of plot 2 is cylindrical meshgrid that is converted to Cartesian meshgrid. The surface equation in cylindrical coordinate system is $z = f(\phi, \rho)$. Hence, cylindrical meshgrids is $m\phi, m\rho = \text{meshgrid}(\phi, \rho)$. Converting cylindrical meshgrids to Cartesian, $mx = m\rho\cos(m\phi)$ and $my = m\rho\sin(m\phi)$ by virtue of geometric relationship between Cartesian and Cylindrical coordinate systems. The meshgrid z is $mz = f(mx, my)$.

3 Intersection of a Cone and of a Plane

The equation of a cone in cylindrical system is expressed as follows.

$$(3.1) \quad \rho = \frac{b_{ase} (height - z)}{height}$$

The variable ϕ is absent. It means that for any value of ϕ the equation holds.

To make the ϕ and ρ as independent variables, the dependent variable is z . Thus,

$$(3.2) \quad z = \frac{height (-\rho + b_{ase})}{b_{ase}}$$

The equations of a plane are expressed as follows.

$$(3.3) \quad Ax + By + Cz + K = 0$$

$$(3.4) \quad z = f(x, y) = -\frac{2x}{5} + \frac{8y}{5} + 5$$

The plot of a cone is shown in Figure 3.1 (a) and the plot of a plane is shown in Figure 3.1 (b).




Figure 3.1: A plane overlapping with cone.

The plot of the cone in Figure 3.1 (a) and the Plot of the plane in Figure 3.1 (b) were plotted together in Figure 3.1 (c). There exists locus of points that are common to both cone and plane surfaces. The locus of points defines a curve line. The curve line is an elliptical as shown in Figure 3.1 (d).

The surface equations are expressed as $z = f(x,y)$ or $x=f(z,y)$ or $y=f(x,z)$. The rhs has the independent variables. The lhs has the dependent variable. Consider the surface plots of a cone and a plane with z as the dependent variable and x and y as the independent variables. Thus,

$$(3.5) \quad z_c = f(x_c, y_c) \quad \text{cone equation}$$

$$(3.6) \quad z_p = f(x_p, y_p) \quad \text{plane equation}$$

A line in 3D space where the cone and the plane intersected could be drawn and plotted. The locus of points must be the common points of the cone and the plane. Hence, $x=x_c=x_p$; $y=y_c=y_p$; and $z=z_c=z_p$

The equations of the cone with x_c a dependent variable and y_c and z_c as independent variable is expressed as follows.

$$(3.7) \quad x_c = f(y_c, z_c)$$

The equations of the plane with y_p a dependent variable and x_p and z_p as independent variable is expressed as follows.

$$(3.8) \quad y_p = f(x_p, z_p)$$

Substituting y_p in y_c ,

$$(3.9) \quad x_c = f(f(x_p, z_p), z_c)$$

Since the locus of points of intersection must be the same for both cone and plane, thus,

$$(3.10) \quad x = f(f(x, z), z)$$

Solving for x ,

$$(3.11) \quad x = f(z)$$

In like process,

$$(3.12) \quad y = f(z)$$

Given a set of values of z , the x and y are computed. The values of x , y , and z forms the locus of points common to both the cone and plane.

The relationships of objects, equations and type of variables for plot algorithm are tabulated as follows.

object	no of equations	no. of dependant variable	no. of independent variable
surface	1	1	2
line	2	2	1
point	0	0	3

4 Conic Sections

The Figure 3 illustrate the mirrored cones that define conic section.

The equation of upper cone expressed in terms Cartesian coordinate system is shown as follows.

$$(4.1) \quad x^2 + y^2 = \frac{4(z - 25)^2}{25}$$

The equation of the disk in terms of Cartesian coordinate system is expressed as follows.

$$(4.2) \quad z = 12$$

The intersection of the upper cone with a disk is shown in Figure 3

The disk plane intersecting with a cone generates a locus of points of circle.

The mirrored cones with a plane Cut perpendicular to the height.




Figure 4.1: Cones section that defined a circle.

The disk could be tilted and the locus of intersecting points forms an ellipse.

The equation of the tilted disk is expressed as follows.

$$(4.3) \quad -0.609710760849692x + 0.7926239891046z = 9.5114878692552$$

The plane cut is at an angle.




Figure 4.2: Conic section cut by disk plane at an angle generated an ellipse an elliptical locus of points.

When the tilt is in parallel with the slope of the side of the cone, the locus of points form a parabola.

The equation of the disk tilted in parallel with the slope of the lateral side of the cone. The cone and the tilted disk is shown in Figure 5.

$$(4.4) \quad -0.970142500145332y - 0.242535625036333z = -7.76114000116266$$

The a plane Cut parallel to the lateral surface generated a parabola.




Figure 4.3: A Conic section that generated a parabola.

When the disk plane is in parallel with the vertical axis of the cone, the locus of points forms a hyperbola. Note the intersection cut across both the upper and lower cones.

The equation of the disk titled in parallel with the cone vertical axis is shown as follows.

$$(4.5) \quad 1.0y = 5.0$$

The plane Cut parallel to the vertical axis of the cones.




Figure 4.4: A Conic section that generated a hyperbola.

5 Exercises

1. Create your own inverted and upright cones. The tips of the cones rest at origin. Formulate the cylindrical and Cartesian equations of the inverted and upright cones. (75)
2. Create your own plane disk perpendicular to the z axis and cut the up-right cone below the origin. Choose your own distance value below the origin. Formulate the disk equation. (5)

3. Formulate the two equations that define the locus of intersection points of the upright cone and the disk. (5)
4. What is the range of disk tilt angle such that the intersection forms circular and/or elliptical curve lines. (5)
5. What is the disk angle of tilt such that the intersection forms a parabolic curve line. (5)
6. What is the disk angle of tilt such that the intersection forms a hyperbolic curve lines. (5)

References

- [1] Daniel Braun Tim Hoffmann (TeXstudio) Pascal Brachet (Texmaker) Luc Buant (QCodeEdit) Joel Amblard (html conversion) Benito van der Zander, Jan Sundermeyer. Textstudio 2.12.10. *Source Forge*, Copyright © 2018. download from <https://sourceforge.net/projects/textstudio/>.
- [2] Karl Berry. Latex2e: An unofficial reference manual, Copyright © 2022, Karl Berry, Latex2e.
- [3] Matthew Brett pv (avatar) Ralf Gommers rkern Travis Oliphant Tyler reddy Burovski, Evgeni. Scipy 1.1.0. *Scipy Developrs. Created using Sphinx 1.6.6* <http://www.sphinx-doc.org/en/master/>, <https://pypi.org/project/scipy/>, 2018. download from <https://pypi.org/project/scipy/#files>.
- [4] cdavid certik Charles R. Harris Jarrod Milman jtaylor Mathew Brett Matti Picus Ralf Gommers rkern Travis Oliphant Haldane, Allan. Numpy 1.15.4. *Numpy Developrs. Created using Sphinx 1.6.6* [urlhttp=//www.sphinx-doc.org/en/master/](http://www.sphinx-doc.org/en/master/), 2018. download from <https://pypi.org/project/numpy/#files>.
- [5] John Hunter. Matplotlib. *MatPlotLib Development Team*, 2002-2012, 2012-2018. downloaded from <https://matplotlib.org/>.
- [6] Donald Knuth. Miktex. *Chrissitian Schenk*. [https=//miktex.org/](https://miktex.org/), 2018.

- [7] Pierre Raybaut. Spyder 3.2.8, the scientific python development environment. *The Spyder Project Contributors, Licensed under the terms of the MIT License*, 2018. Download from <https://github.com/spyder-ide/spyder>.
- [8] Guido van Rossum. Python 3.6.5. *Python Software Foundation*. <https://www.python.org/>, 2018.
- [9] SymPy Development Team. SymPy 1.3. *Github*. <https://github.com/sympy/sympy>, 2018.