

NAME

kh – compute the Henry constant in water or heavy water.

LIBRARY

iapws library (*libiapws*, *-liapws*)

SYNOPSIS

```

Fortran      pure subroutine kh(T,gas,heavywater,k)
               real(dp), intent(in), contiguous :: T(:)
               character(len=*), intent(in) :: gas
               integer(int32), intent(in) :: heavywater
               real(dp), intent(out), contiguous :: k(:)

C            void iapws_g704_kh(double *T,char *gas,int heavywater,double *k,int size_gas,size_t
               size_T)

Python       kh(T:np.ndarray,gas:str,heavywater:bool=False)->Union[np.ndarray,float]:

```

DESCRIPTION

Computes the Henry constant kH in MPa for a given temperature T in water or heavy water.

Arguments:

<i>T(:)</i>	Temperature in K.
<i>gas</i>	Gas.
<i>heavywater</i>	Flag if D2O (1) is used or H2O (0).
<i>k(:)</i>	Henry constant in MPa.

RETURN VALUE

k is filled with NaN if gas is not found.

EXAMPLES

Fortran:

```

use iapws
real(dp) :: T(1), kH(1)
T(1) = 25.0_dp + 273.15_dp
call kh(T, "O2", 0, kH)

```

C:

```

#include "iapws.h"
double T = 25.0 + 273.15;
double kH;
iapws_g704_kh(&T, "O2", 0, &kH, strlen(gas), 1);

```

Python:

```

import numpy as np
import pyiapws
T = np.asarray([25.0])

```

```
kh = pyiapws.kh(T+273.15, "02", False)
print(kh)
```

SEE ALSO

iapws(3) **iapws_cli(1)** **iapws_kh(3)** **ciaaw(3)**, **codata(3)**,