

NAME

kh – compute the Henry constant in water or heavy water.

LIBRARY

iapws library (*libiapws*, *-liapws*)

SYNOPSIS

```

Fortran      pure subroutine kh(T,gas,heavywater,k)
              real(dp), intent(in), contiguous :: T(:)
              character(len=*), intent(in) :: gas
              integer(int32), intent(in) :: heavywater
              real(dp), intent(out), contiguous :: k(:)

C            void iapws_g704_kh(double *T,char *gas,int  heavywater,double *k,int  size_gas,size_t
              size_T)

Python      kh(T:np.ndarray,gas:str,heavywater:bool=False)->Union[np.ndarray,float]:

```

DESCRIPTION

Computes the Henry constant kH in MPa for a given temperature T in water or heavy water.

Arguments:

<i>T</i> (:)	Temperature in K.
<i>gas</i>	Gas.
<i>heavywater</i>	Flag if D2O (1) is used or H2O (0).
<i>k</i> (:)	Henry constant in MPa.

RETURN VALUE

k is filled with NaN if gas is not found.

EXAMPLES

Fortran:

```

use iapws
real(dp) :: T(1), k(1)
T(1) = 25.0_dp + 273.15_dp
call kh(T, "O2", 0, k)

```

C:

```

#include "iapws.h"
double T = 25.0 + 273.15;
char *gas = "O2";
double k;
iapws_g704_kh(&T, gas, 0, &k, strlen(gas), 1);

```

Python:

```

import numpy as np
import pyiapws

```

```
T = np.asarray([25.0])
k = pyiapws.kh(T+273.15, "02", False)
print(k)
```

SEE ALSO

**iapws(1) iapws_lib(3) iapws_psat(3) iapws_Tsat(3) iapws_kh(3) iapws_kd(3) iapws_kw(3) ciaaw(3),
codata(3),**