

Task History

Initiating Search

February 23, 2025, 9:08 PM

Substances:

Filtered By:



Structure Match: As Drawn

Search Tasks

Task		Search Type	View
Returned Substance Results + Filters (2,302)		Substances	View Results
Exported: Retr	ieved Related Reaction Results + Filters (11)	Reactions	View Results
Filtered By:			
Substance Role:	Reagent		
Catalyst:	Manganese(1+), tricarbonyl[(1,2,3,4,5,6-η)-methoxybenzene]-, tetrafluoroborate(1-), Manganese(1+), tricarbonyl[<i>N</i> -[2- (diphenylphosphino-κ <i>P</i>)ethyl]-2-pyridinemethanamine-κ <i>N</i> ¹ ,κ <i>N</i> ²]-, bromide (1:1), (<i>OC</i> -6-44)-, Manganese, [4,8-bis[bis(1-methylethyl)phosphino-κ <i>P</i>]-2,10-bis(1,1-dimethylethyl)-6 <i>H</i> -dipyrido[1,2- <i>c</i> .2',1'- <i>e</i>]imidazol-6-ylidene-κ <i>C</i>]dicarbonylhydro-, (<i>OC</i> -6-42)-[2-[Bis(1-methylethyl)phosphino-κ <i>P</i>]- <i>N</i> -[2-[bis(1-methylethyl)phosphino-κ <i>P</i>]ethyl]ethanaminato-κ <i>N</i>]dicarbonylmanganese		
Document	Journal		
Type: Language:	English		

Copyright © 2025 American Chemical Society (ACS). All Rights Reserved.

Internal use only. Redistribution is subject to the terms of your CAS SciFinder License Agreement and CAS information Use Policies.



Reactions (3)

View in CAS SciFinder

Steps: 1



➤ Suppliers (76)

31-116-CAS-23235270 Steps: 1

Reagents: Sodium *tert*-butoxide, Deuterium
 Catalysts: (*OC*-6-42)-[2-[Bis(1-methylethyl)phosphino-κ*P*]-*N*-[2-[bis(1-methylethyl)phosphino-κ*P*]ethyl]ethanamine-κ*N*] bromodicarbonylmanganese


Solvents: Toluene; rt; 24 h, rt → 150 °C
Experimental Protocols

Carbon monoxide and hydrogen (syngas) as a C1-building block for selective catalytic methylation

By: Kaithal, Akash; et al

Chemical Science (2021), 12(3), 976-982.

Scheme 2 (1 Reaction)



□ Suppliers (17)

■ Suppliers (119)

Steps: 1

31-116-CAS-23235686

1.1 Reagents: Sodium *tert*-butoxide, Deuterium
 Catalysts: (*OC*-6-42)-[2-[Bis(1-methylethyl)phosphino-κ*P*]-*N*-[2-[bis(1-methylethyl)phosphino-κ*P*]ethyl]ethanamine-κ*N*]

bromodicarbonylmanganese

Solvents: Toluene; rt; 24 h, rt → 150 °C


Experimental Protocols

Carbon monoxide and hydrogen (syngas) as a C1-building block for selective catalytic methylation

By: Kaithal, Akash; et al

Chemical Science (2021), 12(3), 976-982.

Scheme 3 (1 Reaction) Steps: 1



Suppliers (119)

CAS SciFinder® Page 4

31-116-CAS-23234681 Steps: 1		Carbon monoxide and hydrogen (syngas) as a C1-building	
 1.1 Reagents: Sodium tert-butoxide, Deuterium Catalysts: (OC-6-42)-[2-[Bis(1-methylethyl)phosp [2-[bis(1-methylethyl)phosphino-κP]ethyl]ethana bromodicarbonylmanganese Solvents: Toluene; rt; 24 h, 10 bar, rt → 150 °C Experimental Protocols 		block for selective catalytic methylation By: Kaithal, Akash; et al Chemical Science (2021), 12(3), 976-982.	

Copyright © 2025 American Chemical Society (ACS). All Rights Reserved.

Internal use only. Redistribution is subject to the terms of your CAS SciFinder License Agreement and CAS information Use Policies.