

Hip pain

Description

409 records of patients with scrotal pain, collected in summer 2003

Created

End of summer 2003 (~ September 2003)

Last update

Summer 2004 (~ August 2004)

Special features

Data cleaned and reviewed by ...

Clinical attributes

#	Code	Name	Possible values
1	NUMBER	Study ID number	
2	AGE	Age (in years)	<i>Real-valued, discretized to:</i> LT10Y < 10 years GE10Y_LE13Y ≥ 10 years and ≤ 13 years GT13Y > 13 years
3	HX_AS	Prior hx of acute scrotal pain	YES, NO
4	NAUSEA	Nausea with current presentation	YES, NO
5	VOMIT	Vomiting with current presentation	YES, NO
6	NAUSEA_OR_VOMIT	Nausea and/or vomiting with current presentation (VOMIT=YES may include NAUSEA=YES, ie. Vomiting overrides Nausea)	NONE, NAUSEA, VOMIT
7	ABD_PAIN	Abdominal pain with current presentation	YES, NO
8	ONSET	Rapidity of onset	GRADUAL, SUDDEN
9	PAIN_TYPE	Type of pain	INTERMITTENT, CONSTANT
10	DURATION	Duration of pain (in hours)	<i>Real-valued, discretized to:</i> LT6H < 6 hours GE6H_LE24H ≥ 6 hours and ≤ 24 hours LT24H > 24 hours
11	ASSOC_UT_SX	Associated dysuria, frequency or urgency	YES, NO
12	HX_TRAUMA	History of minor trauma	YES, NO
13	HX_SPORTS	History of pain during sport	YES, NO
14	HX_SEX	History of prior sexual activity	YES, NO
15	PMHX_UTI	Previous UTI's	YES, NO
16	PMHX_SURG	Previous urologic surgery	YES, NO
17	PMHX_UT_ABN	Previous abnormality of the urinary tract	YES, NO
18	PMHX_URETH_INSTR	Previous urethral instrumentation	YES, NO
19	TEMP	Current temperature (in Celsius)	<i>Real-valued, discretized to:</i> LT38 < 38 °C GE38 ≥ 38 °C

20	LOCATION	Location of pain	<p><i>In initial coding:</i> LEFT_SCROTUM, RIGHT_SCROTUM, BILATERAL_SCROTUM, LEFT_GROIN, RIGHT_GROIN, BILATERAL_GROIN</p> <p><i>In final coding changed to:</i> SINGLE_TESTICLE, BILATERAL_TESTICLE, SINGLE_GROIN, BILATERAL_GROIN</p> <p>LEFT_SCROTUM □ SINGLE_TESTICLE RIGHT_SCROTUM □ SINGLE_TESTICLE BILATERAL_SCROTUM □ BILATERAL_TESTICLE LEFT_GROIN □ SINGLE_GROIN RIGHT_GROIN □ SINGLE_GROIN BILATERAL_GROIN □ BILATERAL_GROIN</p>
21	SITE_TENDER	Site of tenderness on exam	NOT_TENDER, ENTIRE_TESTICLE, UPPER_POLE, POSTERIOR, CANAL_GROIN, OSC, TENDERNESS_NOT_SPECIFIED
22	SITE_SWELLING	Site of swelling on exam	NONE, TESTICLE, OSC, CANAL_GROIN
23	COLOR	Color of scrotum on exam	NORMAL, ERYTHEMA, BLUE_DOT
24	TEST_LIE	Orientation of testicle on exam	<p><i>In initial coding:</i> NORMAL, TRANSVERSE, ELEVATED, TRANSVERSE_ELEVATED</p> <p><i>In final coding changed to:</i> NORMAL, ABNORMAL</p> <p>NORMAL □ NORMAL TRANSVERSE □ ABNORMAL ELEVATED □ ABNORMAL TRANSVERSE_ELEVATED □ ABNORMAL</p>
25	CORD_PALP	Presence of cord on exam	NORMAL, ABNORMAL
26	CREM_REF	Presence of cremasteric reflex	YES, NO
27	TRANSILLUM	Transillumination	YES, NO
28	URINE_WBC	Urine WBC on microscopy	<p><i>In initial coding:</i> NEG_0_5, 5_10, 10_20, 20_50, 50_</p> <p><i>In final coding changed to:</i> NEG_0_5, POS_5_</p> <p>NEG_0_5 □ NEG_0_5 5_10 □ POS_5_ 10_20 □ POS_5_ 20_50 □ POS_5_ 50_ □ POS_5_</p> <p><i>Another suggestion for final coding:</i> NEG_0_10, POS_10_</p> <p>NEG_0_5 □ NEG_0_10 5_10 □ NEG_0_10 10_20 □ POS_10_ 20_50 □ POS_10_ 50_ □ POS_10_</p>
29	URINE_BACT	Bacteria on urine microscopy	NEG, POS

30	URINE_RBC	Urine red cells on microscopy	<p><i>In initial coding:</i> NEG_0_5, 5_10, 10_20, 20_50, 50_</p> <p><i>In final coding changed to:</i> NEG_0_5, POS_5_</p> <p>NEG_0_5 □ NEG_0_5 5_10 □ POS_5_ 10_20 □ POS_5_ 20_50 □ POS_5_ 50_ □ POS_5_</p> <p><i>Another suggestion for final coding:</i> NEG_0_10, POS_10_</p> <p>NEG_0_5 □ NEG_0_10 5_10 □ NEG_0_10 10_20 □ POS_10_ 20_50 □ POS_10_ 50_ □ POS_10_</p>
31	TRIAGE	Final triage category	DISCHARGE, CLINIC, CONSULT