

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/228946002>

Tower of Hanoi Problem with Arbitrary Number of Pegs and Present a Solution

Article in World Applied Sciences Journal · January 2010

CITATIONS

0

READS

1,730

3 authors:



[Sohrab Kordrostami](#)

Islamic Azad University

94 PUBLICATIONS 761 CITATIONS

[SEE PROFILE](#)

[Ramin Ahmadzadeh](#)

University of Maragheh

2 PUBLICATIONS 7 CITATIONS

[SEE PROFILE](#)



[Armin Ghane](#)

Islamic Azad University of Lahijan

14 PUBLICATIONS 22 CITATIONS

[SEE PROFILE](#)

Tower of Hanoi Problem with Arbitrary Number of Pegs and Present a Solution

S. Kordrostami, R. Ahmadzadeh and A. Ghane

Department of Mathematics, Islamic Azad University, Lahijan Branch, Iran

Abstract: Suppose that we have m pegs and n disks of distinct sizes such that initially disks are stacked on the first peg ordered by size with the smallest at the top and the largest at the bottom. In this Paper we want to find a solution for transition disks to one of another pegs using the well-known movements of Tower of Hanoi such that there are no solution better than it yet.

Key words: Tower of Hanoi % Recursive formula % Pegs

INTRODUCTION

In the well-known Tower of Hanoi problem, composed over a hundred years ago by Lucas [1], a player is given 3 pegs and a certain number n of disks of distinct sizes and is required to transfer them from one peg to another.

Initially all disks are stacked (composing a tower) on the first peg (the source) ordered by size, with the smallest at the top and the largest at the bottom. The goal is to transfer them to the third peg (the destination), moving only topmost disks and never placing a disk on top of a smaller one. The known recursive algorithm, which may be easily shown to be optimal, takes $h_n = 2^n - 1$ steps to accomplish the task. Since there is not much mathematical mystery left about the original game, its lovers developed various versions of it [2-6]. In this paper we want to solve Tower of Hanoi problem with arbitrary number of pegs.

Tower of Hanoi with M Pegs: Suppose that we have n disks of distinct sizes d_1, d_2, \dots, d_n such that $d_1 > d_2 > \dots > d_n$ and m pegs ($m > 3$). Initially disks are stacked on the first peg ordered by size (Figure 1). The goal is to transfer all disks to one of another pegs.

Algorithm of Solution: For transfer of disks from peg 1 to one of another pegs such as peg m , at the first d_1 must be placed in the bottom of the peg m . It is possible when all other disks be stacked in another $m-2$ pegs. Then d_2 must be transferred to peg m , top of d_1 . But it is possible when d_2 be in one peg alone, for example peg 2 and all disks except d_1 and d_2 be stacked in other $m-3$ pegs. In this manner for transition of d_{m-2} to peg m top of d_1, d_2, \dots, d_{m-3} disks must have been a position as following (Figure 2).




Fig. 1: The board of the Tower of Hanoi problem with m pegs




Fig. 2: Steps of transition of disks

That means $n - (m - 2)$ remainder disks must be stacked on one peg such as peg $m - 1$ in Figure, 2. certainly. In fact we want to find a recursive formula using this idea for solving this problem.

If we show the number of motions for transition of n disks to the peg m with H_n^m , it is obvious that if $n < m$ then $H_n^m = 2n - 1$.

Now with this introductions we want to find H_n^m for $n \geq m$, using following steps:

Step 1: Transition of disks d_{m-3}, \dots, d_n with $H_{n-(m-2)}^m$ motions from peg 1 to peg $m - 1$.

Step 2: Transition of disks d_2, d_3, \dots, d_{m-2} 2 from peg 1 to pegs 2, 3, ..., $m-2$, with $m-3$ motions.

Step 3: Transition of disks $d_1, d_2, d_3, \dots, d_{m-2}$ to peg m with $m-2$ motions.

Step 4: Transition of disks d_{m-3}, \dots, d_n with $H_{n-(m-2)}^m$ motions from peg $m-1$ to peg m .

Therefore for $n \geq m$, we have:

$$H_n^m = H_{n-(m-2)}^m + (m-3) + (m-2) + H_{n-(m-2)}^m$$

$$\Rightarrow H_n^m = 2H_{n-(m-2)}^m + 2m-5$$

Upper relation is a non-homogeneous recursive formula of degree $m-2$ and we want to solve it with primal conditions:

$$H_0^m = 0, H_1^m = 1, H_2^m = 3, \dots, H_{m-3}^m = 2(m-3) - 1$$

We must solve this recursive formula in $m-2$ cases:

Case 1: $n = (m-2)k$

Case 2: $n = (m-2)k+1$

Case 3: $n = (m-2)k+2$

!

Case m-2: $n = (m-2)k + (m-3)$

In all upper cases k is a non-negative integer number.

Studing of Case 1:

In this case $n = (m-2)k$ and we have:

$$H_n^m = 2H_{n-(m-2)}^m + 2m-5$$

$$= 2(2H_{n-2(m-2)}^m + 2m-5) + 2m-5 = 2^2 H_{n-2(m-2)}^m + 3(2m-5)$$

$$\vdots$$

$$= 2^k H_{n-k(m-2)}^m + (2^k - 1)(2m-5) = 2^k H_0^m + (2^k - 1)(2m-5)$$

$$\Rightarrow H_n^m = (2^{\frac{n}{m-2}} - 1)(2m-5)$$

Studing of Case 2:

In this case $n = (m-2)k+1$ and we have:

$$H_n^m = 2^k H_1^m + (2^k - 1)(2m-5) = 2^k + (2^k - 1)(2m-5)$$

$$\Rightarrow H_n^m = 2^{\frac{n-1}{m-2}} + (2^{\frac{n-1}{m-2}} - 1)(2m-5)$$

Studing of Case 3:

In this case $n = (m-2)k+2$ and we have:

$$H_n^m = 2^k H_2^m + (2^k - 1)(2m-5) = (2^k \times 3) + (2^k - 1)(2m-5)$$

$$\Rightarrow H_n^m = 3 \times 2^{\frac{n-2}{m-2}} + (2^{\frac{n-2}{m-2}} - 1)(2m-5)$$

Studing of Case m-2:

In this case $n = (m-2)k + (m-3)$ and we have:

$$H_n^m = 2^k H_{m-3}^m + (2^k - 1)(2m-5) = 2^k (2(m-3) - 1) + (2^k - 1)(2m-5)$$

$$\Rightarrow H_n^m = (2(m-3) - 1)2^{\frac{n-(m-3)}{m-2}} + (2^{\frac{n-(m-3)}{m-2}} - 1)(2m-5)$$

Therefore in the abstract we have:

If $n = (m-2)k$ then

$$H_n^m = (2^{\frac{n}{m-2}} - 1)(2m-5)$$

If $n = (m-2)k+1$ then

$$H_n^m = 2^{\frac{n-1}{m-2}} + (2^{\frac{n-1}{m-2}} - 1)(2m-5)$$

If $n = (m-2)k+2$ then

$$H_n^m = 3 \times 2^{\frac{n-2}{m-2}} + (2^{\frac{n-2}{m-2}} - 1)(2m-5)$$

!

If $n = (m-2)k + (m-3)$ then

$$H_n^m = (2(m-3) - 1)2^{\frac{n-(m-3)}{m-2}} + (2^{\frac{n-(m-3)}{m-2}} - 1)(2m-5)$$

CONCLUSION

We found a solution for transfer of n disks using the well-known movements of Tower of Hanoi with m pegs from one peg to one of another pegs. We did not find optimal solution, but there are no solution better than this solution yet.

In general case, H_n^m can be obtained as following:

$$H_n^m = \begin{cases} (2m-5)(2^{\frac{n}{m-2}} - 1) & i = 0 \\ (2i-1)2^{\frac{n-i}{m-2}} + (2m-5)(2^{\frac{n-i}{m-2}} - 1) & i \neq 0 \end{cases}$$

Such that i is remainder of division of n by $m-2$ ($i = n \bmod (m-2)$).

REFERENCES

1. Lucas, E., 1893. *Recreations Mathematiques*, Volume 3, Gauthier-Villars, Paris.
2. A.Sapir, A., 2004. The tower of Hanoi with forbidden moves, *The Computer J.*, 47(1): 20-24.
3. Stockmeyer, P.K., 1994. Variations on the four-post Tower of Hanoi puzzle, *Congr*, 102: 3-12.
4. Kordrostami, S., R. Ahmadzadeh, A. Ghane and S. Pourjafar, 2007. Review of Tower of Hanoi and Present New Formula, *J. Applied Mathematics*, Islamic Azad University of Lahijan, 4(14): 41-47.
5. Szegedy, M., 1999. In how many steps the k peg version of the Tower of Hanoi game can be solved ? STACS 99 (Trier), *Lecture Notes in computer Science* 1563, Springer, Berlin, pp: 356-361.
6. Klavzar, S., U. Milutinovic and C. Petr, 2002. On the Frame-Stewart algorithm for the multi-peg Tower of Hanoi Problem *Discrete Appl. Math.*, 120: 141-157.