

Banach algebras

Contents

1. Introduction	4
2. Semi-normed Abelian groups	4
3. Semi-normed rings	5
4. Banach rings	5
5. Semi-normed modules	7
Bibliography	9

1. Introduction

This section concerns the theory of Banach algebras. Our references are [Ber12] and [BGR84].

In this chapter, all rings are assumed to be commutative.

2. Semi-normed Abelian groups

Definition 2.1. Let A be an Abelian group. A *semi-norm* on A is a function $\|\bullet\| : A \rightarrow [0, \infty]$ satisfying

- (1) $\|0\| = 0$;
- (2) $\|f - g\| \leq \|f\| + \|g\|$ for all $f, g \in A$.

A semi-norm $\|\bullet\|$ on A is a *norm* if moreover the following conditions is satisfied:

- (0) if $\|f\| = 0$ for some $f \in A$, then $f = 0$.

A semi-norm $\|\bullet\|$ on A is *non-Archimedean* or *ultra-metric* if Condition (2) can be replaced by

- (2') $\|f - g\| \leq \max\{\|f\|, \|g\|\}$ for all $f, g \in A$.

Definition 2.2. A *semi-normed Abelian group* (resp. *normed Abelian group*) is a pair $(A, \|\bullet\|)$ consisting of an Abelian group A and a semi-norm (resp. norm) $\|\bullet\|$ on A . When $\|\bullet\|$ is clear from the context, we also say A is a semi-normed Abelian group (resp. normed Abelian group).

Definition 2.3. Let $(A, \|\bullet\|_A)$ be a semi-normed Abelian group and $B \subseteq A$ be a subgroup. Then we define the *quotient semi-norm* $\|\bullet\|_{A/B}$ on A/B as follows:

$$\|a + B\|_{A/B} := \inf\{\|a + b\|_A : b \in B\}$$

for all $a + B \in A/B$.

We define the *subgroup semi-norm* on B as follows:

$$\|b\|_B = \|b\|_A$$

for all $b \in B$.

Definition 2.4. Let A be an Abelian group and $\|\bullet\|, \|\bullet\|'$ be two seminorms on A . We say $\|\bullet\|$ and $\|\bullet\|'$ are *equivalent* if there is a constant $C > 0$ such that

$$C^{-1}\|f\| \leq \|f\|' \leq C\|f\|$$

for all $f \in A$.

Definition 2.5. Let $(A, \|\bullet\|_A), (B, \|\bullet\|_B)$ be semi-normed Abelian groups. A homomorphism $\varphi : A \rightarrow B$ is said to be

- (1) *bounded* if there is a constant $C > 0$ such that $\|\varphi(f)\|_B \leq C\|f\|_A$ for any $f \in A$;
- (2) *admissible* if the quotient semi-norm on $A/\ker \varphi$ is equivalent to the subspace semi-norm on $\text{Im } \varphi$.

Observe that an admissible homomorphism is always bounded.

Next we study the topology defined by a semi-norm.

Lemma 2.6. Let $(A, \|\bullet\|)$ be a semi-normed Abelian group. Define

$$d(a, b) = \|a - b\|$$

for $a, b \in A$. Then $\|\bullet\|$ is a pseudo-metric on A . This pseudo-metric is a metric if and only if $\|\bullet\|$ is a norm.

PROOF. This is clear from the definitions. \square

We always endow A with the topology induced by the psuedo-metric d .

3. Semi-normed rings

Definition 3.1. Let A be a ring. A *semi-norm* $\|\bullet\|$ on A is a semi-norm $\|\bullet\|$ on the underlying additive group satisfying the following extra properties:

- (3) $\|1\| = 1$;
- (4) for any $f, g \in A$, $\|fg\| \leq \|f\|\|g\|$.

A semi-norm $\|\bullet\|$ on A is called *power-multiplicative* if $\|f\|^n = \|f^n\|$ for all $f \in A$ and $n \in \mathbb{N}$.

A semi-norm $\|\bullet\|$ on A is called *multiplicative* if $\|fg\| = \|f\|\|g\|$ for all $f, g \in A$.

Definition 3.2. A *semi-normed ring* (resp. *normed ring*) is a pair $(A, \|\bullet\|)$ consisting of a ring A and a semi-norm (resp. norm) $\|\bullet\|$ on A . When $\|\bullet\|$ is clear from the context, we also say A is a semi-normed ring (resp. normed ring).

Definition 3.3. Let A be a ring. A *semi-valuation* on A is a multiplicative semi-norm on A . A semi-valuation on A is a *valuation* on A if its underlying semi-norm of Abelian groups is a norm.

Definition 3.4. A *semi-valued ring* (resp. *valued ring*) is a pair $(A, \|\bullet\|)$ consisting of a ring A and a semi-valuation (resp. valuation) $\|\bullet\|$ on A . When $\|\bullet\|$ is clear from the context, we also say A is a semi-valued ring (resp. valued ring).

A semi-valued ring (resp. valued ring) $(A, \|\bullet\|)$ is called a *semi-valued field* (resp. *valued field*) if A is a field.

4. Banach rings

Definition 4.1. A *Banach ring* is a normed ring that is complete with respect to the metric defined in [Lemma 2.6](#).

Proposition 4.2. Let $(A, \|\bullet\|)$ be a Banach ring and $f \in A$. Assume that $\|f\| < 1$, then $1 - f$ is invertible.

PROOF. Define

$$g = \sum_{i=0}^{\infty} f^i.$$

From our assumption, the series converges and $g \in A$. It is elementary to check that g is the inverse of $1 - f$. \square

Example 4.3. The ring \mathbb{C} with its usual norm $|\bullet|$ is a Banach ring. In fact, $(\mathbb{C}, |\bullet|)$ is a complete valued field.

Example 4.4. For any Banach ring $(A, \|\bullet\|)$, any $n \in \mathbb{N}$ and any $r = (r_1, \dots, r_n) \in \mathbb{R}_{>0}^n$, we define $A\langle r^{-1}z \rangle = A\langle r_1^{-1}z_1, \dots, r_n^{-1}z_n \rangle_{r_1, \dots, r_n}$ as the subring of $A[[z_1, \dots, z_n]]$ consisting of formal power series

$$f = \sum_{\alpha \in \mathbb{N}^n} a_{\alpha} z^{\alpha}, \quad a_{\alpha} \in A$$

such that

$$\|f\|_r := \sum_{\alpha \in \mathbb{N}^n} \|a_{\alpha}\| r^{\alpha} < \infty.$$

We will verify in [Proposition 4.5](#) that $(A\langle r^{-1}z \rangle, \|\bullet\|_r)$ is a Banach ring.

When $r = (1, \dots, 1)$, we omit r^{-1} from our notations.

Proposition 4.5. In the setting of [Example 4.4](#), $(A\langle r^{-1}z \rangle, \|\bullet\|_r)$ is a Banach ring.

PROOF. By induction, we may assume that $n = 1$.

It is obvious that $\|\bullet\|_r$ is a norm on the underlying Abelian group. To see that $\|\bullet\|_r$ is a norm on the ring $A\langle r^{-1}z \rangle$, we need to verify the condition in [Definition 3.1](#). Condition (3) in [Definition 3.1](#) is obvious. Let us consider Condition (4). Let

$$f = \sum_{i=0}^{\infty} a_i z^i, \quad g = \sum_{j=0}^{\infty} b_j z^j$$

be two elements in $A\langle r^{-1}z \rangle$. Then

$$fg = \sum_{k=0}^{\infty} \left(\sum_{i+j=k} a_i b_j \right) z^k.$$

We compute

$$\|fg\|_r = \sum_{k=0}^{\infty} \left\| \sum_{i+j=k} a_i b_j \right\|_r r^k \leq \sum_{k=0}^{\infty} \left(\sum_{i+j=k} \|a_i\| \cdot \|b_j\| \right) r^k = \|f\|_r \cdot \|g\|_r.$$

It remains to verify that $A\langle r^{-1}z \rangle$ is complete.

For this purpose, take a Cauchy sequence

$$f^b = \sum_{i=0}^{\infty} a_i^b z^i \in A\langle r^{-1}z \rangle$$

for $b \in \mathbb{N}$. Then for each i , the coefficients $(a_i^b)_b$ is a Cauchy sequence in A . Let a_i be the limit of a_i^b as $b \rightarrow \infty$ and set

$$f = \sum_{i=0}^{\infty} a_i z^i \in A[[z]].$$

We need to show that $f \in A\langle r^{-1}z \rangle$ and $f^b \rightarrow f$.

Fix a constant $\epsilon > 0$. There is $m = m(\epsilon) > 0$ such that for all $j \geq m$ and all $k \geq 0$, we have

$$\sum_{i=0}^{\infty} \|a_i^{j+k} - a_i^j\| r^i < \epsilon/2.$$

In particular, for any $s > 0$, we have

$$\sum_{i=0}^s \|a_i - a_i^j\| r^i \leq \sum_{i=0}^s \|a_i - a_i^{j+k}\| r^i + \sum_{i=0}^s \|a_i^j - a_i^{j+k}\| r^i \leq \sum_{i=0}^s \|a_i - a_i^{j+k}\| r^i + \epsilon/2.$$

When k is large enough, we can guarantee that

$$\sum_{i=0}^s \|a_i - a_i^{j+k}\| r^i < \epsilon/2.$$

So

$$\sum_{i=0}^s \|a_i - a_i^j\| r^i \leq \epsilon.$$

Let $s \rightarrow \infty$, we find

$$\|f - f^j\|_r \leq \sum_{i=0}^{\infty} \|a_i - a_i^j\| r^i \leq \epsilon.$$

In particular, $\|f\|_r < \infty$ and $f^j \rightarrow f$ as $j \rightarrow \infty$. \square

5. Semi-normed modules

Definition 5.1. Let $(A, \|\bullet\|_A)$ be a normed ring. A *semi-normed A -module* (resp. *normed A -module*) is a pair $(M, \|\bullet\|_M)$ consisting of a A -module M and a semi-norm (resp. norm) on the underlying Abelian group of M such that there is a constant $C > 0$ such that

$$\|fm\|_M \leq C\|f\|_A\|m\|_M$$

for all $f \in A$ and $m \in M$. When $\|\bullet\|_M$ is clear from the context, we say M is a semi-normed A -module (resp. normed A -module).

A *Banach A -module* is a normed A -module which is complete with respect to the metric [Lemma 2.6](#).

Bibliography

- [Ber12] V. G. Berkovich. Spectral theory and analytic geometry over non-Archimedean fields. 33. American Mathematical Soc., 2012.
- [BGR84] S. Bosch, U. Güntzer, and R. Remmert. Non-Archimedean analysis. Vol. 261. Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. A systematic approach to rigid analytic geometry. Springer-Verlag, Berlin, 1984, pp. xii+436. URL: <https://doi.org/10.1007/978-3-642-52229-1>.